


1993
Distribution of
Submerged Aquatic Vegetation
in the Chesapeake Bay



Virginia Institute of Marine Science
School of Marine Science
The College of William & Mary



Distribution of Submerged Aquatic Vegetation in the Chesapeake Bay and Tributaries and Chincoteague Bay - 1993

by

Robert J. Orth, Judith F. Nowak, Gary F. Anderson,
and Jennifer R. Whiting

College of William and Mary
School of Marine Science
Virginia Institute of Marine Science
Gloucester Point, VA 23062

Funded by: U.S. Environmental Protection Agency (Grant CB003909-02)

National Oceanographic Atmospheric Administration
(Grant No. NA370Z0360-01)

College of William and Mary, School of Marine Science,
Virginia Institute of Marine Science

Maryland Department of Natural Resources
(C350-94-007)

U.S. Fish and Wildlife Service (14-16-0005-93-9038)

Allied-Signal Foundation

Final Report Submitted to: U.S. Environmental Protection Agency
Chesapeake Bay Program Office
Annapolis, MD 21403

December 1994

Cover Photograph: John Flood, a participant in the Citizens' SAV Survey, holds "doubler" blue crabs (*Callinectes sapidus*) from a large bed of Redhead grass (*Potamogeton perfoliatus*) in Round Bay, at the mouth of Asquith Creek, Severn River, Maryland, 1994. This bed appeared for the first time in 1994. (Photography courtesy of Bob Gilbert of *The Capital* newspaper, Annapolis, Maryland.)

CONTENTS

	Page
TABLES	v
FIGURES	vi
EXECUTIVE SUMMARY	ix
ACKNOWLEDGEMENTS	xv
SAV SPECIES	1
METHODS	2
Introduction	2
Aerial photography	2
Mapping process	5
SAV Perimeter Digitization and Quality Assurance Procedures	10
Calculation of 1993 SAV Areas	12
Organizational procedures for analysis and discussion	12
Ground surveys and other data bases	18
RESULTS	20
Data presentation	20
1993 Summary	20
Upper Bay Zone	20
Middle Bay Zone	44
Lower Bay Zone	45
Chincoteague Bay	45
Discussion of sections arranged within zones	46
Upper Bay Zone	46
Susquehanna Flats (Section 1)	46
Upper Eastern Shore (Section 2)	46
Upper Western Shore (Section 3)	46
Chester River (Section 4)	49
Middle Bay Zone	49
Central Western Shore (Section 5)	49
Eastern Bay (Section 6)	53
Choptank River (Section 7)	53

CONTENTS (continued)

	Page
RESULTS (continued)	
Patuxent River (Section 8)	53
Middle Western Shore (Section 9)	59
Lower Potomac River (Section 10)	59
Upper Potomac River (Section 11)	59
Middle Eastern Shore (Section 12)	67
Mid-Bay Island Complex (Section 13)	67
Lower Bay Zone	67
Lower Eastern Shore (Section 14)	67
Reedville Region (Section 15).	71
Rappahannock River Complex (Section 16).	71
New Point Comfort Region (Section 17)	74
Mobjack Bay Complex (Section 18).	74
York River (Section 19)	74
Lower Western Shore (Section 20)	78
James River (Section 21)	78
Chincoteague Bay.	78
LITERATURE CITED	84
APPENDICES	87
A. Species of submerged aquatic plants found in Chesapeake Bay and tributaries	89
B. Latitude and longitude coordinate points defining the 21 Chesapeake Bay sections and Chincoteague Bay	93
C. USGS 7.5 minute quadrangles for Chesapeake Bay and Chincoteague Bay showing distribution, abundance, and ground truthing of SAV in 1993	101
Map Legend.	102
D. Number of square meters of SAV for individual beds and totals for density categories for each USGS 7.5 minute quadrangle in 1993	211
E. 1993 SAV ground survey data listed by USGS 7.5 minute quadrangle and by 1993 SAV bed	243
Key to Appendix E	244

TABLES

Number	Page
1 Guidelines followed during acquisition of aerial photographs	4
2 List of USGS 7.5 minute quadrangles for Chesapeake Bay and Chincoteague Bay SAV study areas with corresponding code numbers	7
3 Area descriptions for each of the 21 sections of the Chesapeake Bay SAV study area.	13
4 Total area of SAV in hectares by USGS 7.5 minute quadrangles for 1992 and 1993	21
5 Number of hectares of SAV in 1992 and 1993 for the 21 sections and three zones of Chesapeake Bay, and for Chincoteague Bay	27
6 Number of square meters of SAV in 1993 for each USGS 7.5 minute quadrangle of the 21 sections of Chesapeake Bay and of Chincoteague Bay	28
7 Number of square meters of SAV in 1993 by density class for the 21 sections of Chesapeake Bay and for Chincoteague Bay	39

FIGURES

Number	Page	
1	Map of Chesapeake Bay and tributaries with Upper, Middle, and Lower zones, and of Chincoteague Bay, with locations of all SAV beds in 1993	x
2	A comparison of the total hectares of SAV for the Upper, Middle, and Lower zones of the Chesapeake Bay in 1992 and 1993	xi
3	Number of hectares SAV per density class in 1993 by section and zone of the Chesapeake Bay.	xii
4	Map of Chesapeake Bay, its tributaries, and of Chincoteague Bay, with approximate locations of flight lines for 1993 SAV photography.	3
5	Location of USGS 7.5 minute quadrangles in Chesapeake Bay, its tributaries, and in Chincoteague Bay with corresponding code numbers.	6
6	Crown density scale used for estimating density of SAV beds from aerial photography.	11
7	Location of Chincoteague Bay and Chesapeake Bay with Upper, Middle, and Lower zones and 21 sections used for delineation of SAV distribution patterns	16
8	Distribution of SAV in the Susquehanna Flats (Section 1) in 1993	47
9	Distribution of SAV in the Upper Eastern Shore (Section 2) in 1993	48
10	Distribution of SAV in the Upper Western Shore (Section 3) in 1993	50
11	Distribution of SAV in the Chester River (Section 4) in 1993	51
12	Distribution of SAV in the Central Western Shore (Section 5) in 1993	52
13	Distribution of SAV in the Eastern Bay (Section 6) in 1993	54
14	Distribution of SAV in the Choptank River (Section 7) in 1993	55

FIGURES (continued)

Number	Page
15	Detail of Figure 14 showing distribution of SAV in the Choptank River 56
16	Distribution of SAV in the Patuxent River (Section 8) in 1993 57
17	Detail of Figure 16 showing distribution of SAV in the Patuxent River 58
18	Distribution of SAV in the Middle Western Shore (Section 9) in 1993 60
19	Distribution of SAV in the Lower Potomac River (Section 10) in 1993 61
20	Detail of Figure 19 showing distribution of SAV in the Lower Potomac River 62
21	Detail of Figure 19 showing distribution of SAV in the Lower Potomac River 63
22	Distribution of SAV in the Upper Potomac River (Section 11) in 1993 64
23	Detail of Figure 22 showing distribution of SAV in the Upper Potomac River. 65
24	Detail of Figure 22 showing distribution of SAV in the Upper Potomac River. 66
25	Distribution of SAV in the Middle Eastern Shore (Section 12) in 1993 68
26	Distribution of SAV in the Mid-Bay Island Complex (Section 13) in 1993 69
27	Distribution of SAV in the Lower Eastern Shore (Section 14) in 1993 70
28	Distribution of SAV in the Reedville Region (Section 15) in 1993. 72
29	Distribution of SAV in the Rappahannock River Complex (Section 16) in 1993 . . . 73
30	Distribution of SAV in the New Point Comfort Region (Section 17) in 1993 75
31	Distribution of SAV in the Mobjack Bay Complex (Section 18) in 1993 76
32	Distribution of SAV in the York River (Section 19) in 1993 77
33	Distribution of SAV in the Lower Western Shore (Section 20) in 1993. 79
34	Distribution of SAV in the James River (Section 21) in 1993 80

FIGURES (continued)

Number	Page
35 Detail of Figure 34 showing the distribution of SAV in the James River	81
36 Distribution of SAV in the Chincoteague Bay in 1993	82

EXECUTIVE SUMMARY

The distribution of submerged aquatic vegetation, principally rooted vascular macrophytes, in the Chesapeake Bay, its tributaries, and Chincoteague Bay, was mapped from black and white aerial photographs taken during May to October 1993 at a scale of 1:24,000. SAV bed perimeter information was digitized and stored in a computerized data base. Ground truth information was obtained from the U. S. Fish and Wildlife Service; USGS National Center; Harford Community College, Maryland; Maryland-National Capital Parks and Planning Commission, Patuxent River Park; and the College of William and Mary, School of Marine Science, Virginia Institute of Marine Science. Citizen support via the U. S. Fish and Wildlife Service and the Chesapeake Bay Foundation provided additional ground truth information.

In 1993, the Chesapeake Bay had 29,589 hectares of SAV, compared to 28,591 hectares in 1992, with 2,700 hectares (9.0%), 13,901 hectares (47.0%), and 13,018 hectares (44.0%) occurring in the Upper, Middle, and Lower Bay zones, respectively (Figures 1, 2, and 3). SAV generally increased in each of the three zones from 1992. Decreases in some sections (e.g. Mid-Bay Island Complex, Upper Eastern Shore, and Western Shore) were offset by larger increases in other sections (e.g. Choptank River, Eastern Bay, and Chester River sections). SAV increased in abundance in all sections in the Lower Bay zone.

In 1993, in the Upper Bay zone, 66.5% (1,777 hectares) of the SAV was located in the Susquehanna Flats (Section 1). Overall abundance and density of SAV was similar to the 1992 level (1,792 hectares). In the Flats, 85.4% of all SAV beds were classified as very sparse in 1993 (0-10% coverage), while 7.7% of beds were classified as dense (70-100% coverage) (Figure 3). *Myriophyllum spicatum*, *Heteranthera dubia*, *Vallisneria americana*, *Hydrilla verticillata*, *Ceratophyllum demersum*, and *Najas guadalupensis* were the six species reported. In the Upper Eastern Shore (Section 2) there were 184 hectares of SAV in 1993 (99 hectares less than in 1992) located principally in the Elk and lower Sassafras rivers, with *M. spicatum* and *V. americana* found most frequently, especially in the Elk River. Much of the difference from 1992 in the Upper Eastern Shore section was recorded from the Elk River. The Upper Western Shore (Section 3) had 80 hectares of SAV compared to 186 hectares recorded in 1992. SAV was reported from the Gunpowder River area including Dundee Creek; the lower Spesutie Narrows; the Middle and Magothy rivers; and Romney and Delph creeks. SAV was noticeably reduced in Saltpeter Creek from 1992 and was absent from Seneca Creek in 1993. SAV was mapped in the Magothy River for the first time since it was last reported in 1978. *Myriophyllum spicatum*, *Elodea canadensis*, and *Zannichellia palustris* were frequently cited. In the Chester River (Section 4) SAV abundance (629 hectares) was up 374 hectares from 1992. SAV was most abundant adjacent to Eastern Neck, Eastern Neck Island, and in the lower Chester River. *Ruppia maritima* was most commonly cited.

In 1993, 39.3% (5,467 hectares) of the SAV in the Middle Bay zone was found in the Mid-Bay Island Complex (Section 13) which includes the broad shoal area between Smith and Tangier Islands. This is a decrease of 527 hectares over 1992. In this zone, 21.4% (2972 hectares) of the SAV was present in the Middle Eastern Shore (Section 12), primarily in the Barren Island-Honga River area; the Big

UPPER

MIDDLE

LOWER




Figure 1. Map of Chesapeake Bay and tributaries with Upper, Middle, and Lower zones, and of Chincoteague Bay, with locations of all SAV beds in 1993 (SAV is shown in red).

Hectares of SAV in Each Zone of the Chesapeake Bay, 1992-93




Figure 2. A comparison of the total hectares of SAV for the Upper, Middle, and Lower zones of Chesapeake Bay in 1992 and 1993. (Refer to Figures 1 and 7 for zone locations.)

Hectares of SAV in 1993 by Section




Figure 3. Number of hectares SAV per density class in 1993 by section and zone of Chesapeake Bay. (Refer to Figure 7, Table 3, and Appendix B for section locations and boundaries.)

and Little Annessex rivers; and the lower section of the Manokin River, with *R. maritima* reported most frequently. SAV was much less abundant in the Barren Island area. No SAV was mapped from the Central Western Shore (Section 5) and Middle Western Shore (Section 9). SAV in the Patuxent River (Section 8) increased from 0 hectares in 1992 to 10 hectares in 1993. Citizens' surveys reported *Z. palustris* and *R. maritima* at numerous locations in the South and Severn rivers. Eleven species were reported in the Patuxent River.

The Middle Bay zone also includes the entire Potomac River, where 2,820 hectares of SAV were present in 1993. SAV was concentrated in two distinct regions: 1) the Upper Potomac River (Section 11) with 2,363 hectares; and 2) the upper portion of the Lower Potomac River (Section 10) with 458 hectares, including Nanjemoy Creek and Port Tobacco River. Although the total abundance of SAV in the Upper Potomac section decreased from 1992 by 99 hectares, there was a notable increase in SAV in the Alexandria quadrangle. In particular, a large shoal area in the middle Potomac River, just above the large bed around the Woodrow Wilson Bridge, which had been previously unvegetated, supported sparse SAV for the first time since this survey was initiated. Ground truth data was reported by USGS and Citizens' surveys with 9 species reported: *M. spicatum*, *V. americana*, *H. verticillata*, *H. dubia*, *C. demersum*, *Najas minor*, *Potamogeton pectinatus*, *Potamogeton crispus*, and *N. guadalupensis*. SAV in the lower Potomac River also decreased (113 hectares) in 1993. Ground truth data was reported by Citizens' and VIMS surveys with 4 species cited: *Z. palustris*, *P. crispus*, *R. maritima*, and *M. spicatum*. SAV continued to increase in the Eastern Bay and Choptank River sections from 1992. SAV in the Eastern Bay (Section 6) increased 183 hectares from 1992 to a total of 737 hectares in 1993, while in the Choptank River (Section 7) it increased 809 hectares from 1992 to a total of 1,894 hectares in 1993. Most of the increase in the Eastern Bay occurred in Cox Creek, Crab Alley Bay, around Parson Island, and Piney Neck Point. In the Choptank River section, SAV beds were most abundant in Harris and Broad creeks and in Trippe Bay. Two species were reported from Section 6, with *R. maritima* most commonly cited. Three species were reported from Section 7, with *R. maritima* most commonly cited.

Distribution and abundance of SAV in 1993, in the Lower Bay zone, were similar to 1992. In this zone, 48.4% (6,299 hectares) of the SAV was found in the Lower Eastern Shore (Section 14) around the Fox, Cedar, Webb, and Halfmoon islands; and the mouths of major creeks (i.e. Cherrystone Inlet; Hungars, Mattawoman, Nassawadox, Occohannock, Craddock, Pungoteague, Nandua, and Onancock creeks; and Beasley Bay). Along the western shore of the Chesapeake Bay, SAV was abundant in Mobjack Bay (Section 18) (14.2% of SAV in the Lower Bay zone), in the lower York River (Section 19) (6.5% of SAV in the Lower Bay zone), and in the Lower Western Shore (Section 20), specifically Back River and the Drum Island Flats area adjacent to Plum Tree Island (15.9% of SAV in the Lower Bay zone). Sparse SAV was documented for a segment of the south shore of the York River, downstream from Yorktown, for the first time in over twenty years (Orth and Gordon, 1975). There were 813 hectares of SAV mapped in the Reedville Region (Section 15) in 1993, a 4.5% increase over 1992. There were 431 hectares of SAV identified in 1993 in the New Point Comfort Region (Section 17), compared to 396 hectares in 1992. SAV abundance was up 23.1% from 1992 in both the Piankatank and Rappahannock rivers (Section 16). The James River (Section 21) had 4 hectares of SAV in 1993. *Zostera marina* and *R. maritima* were the abundant species in this zone.

SAV in the Chincoteague Bay section increased in distribution with 3,576 hectares mapped in 1993 compared to 3,324 hectares in 1992. Most of the SAV in Chincoteague and Sinepuxent bays, which consisted mainly of *Z. marina* and *R. maritima*, was located along the eastern side of the bay behind Assateague Island. Some small beds, consisting mainly of *R. maritima*, were located along the eastern side of Isle of Wight and Assawoman bays.

ACKNOWLEDGEMENTS

We would like to gratefully acknowledge all federal and state agencies who financially supported this project: the U. S. Environmental Protection Agency (USEPA); the U. S. Fish and Wildlife Service (USFWS); the Virginia Department of Environmental Quality's and MD-DNR's Coastal Resources Management Program of the National Oceanographic and Atmospheric Administration (NOAA), under the Coastal Zone Management Act of 1972 as amended by the Office of Ocean Coastal Resource Management (OCRM); the Maryland Department of Natural Resources (MD-DNR); and the Virginia Institute of Marine Science, School of Marine Science, College of William and Mary (VIMS). Also, we thank Allied - Signal, Inc. for providing supplemental funding for this work.

Acknowledgement would not be complete without commendation for the groups which provided ground truthing of SAV beds which was used in conjunction with interpretation of the 1993 photography. USFWS with the Chesapeake Bay Foundation (CBF) organized citizens to report locations and species composition of grass beds around the bay. The U.S. Geological Survey (USGS), the USFWS, and Stan Kollar of Harford Community College (HCC), Maryland, provided ground truth information for certain specific regions of the Maryland portion of the Bay. Patuxent River Park staff provided ground truth data for the Patuxent River. Ken Moore, Curtis Harper, Jill Goodman, Susan Bogardy, and James Fishman of VIMS provided ground truth information for the lower bay.

The production of this report required the dedication of numerous scientists, technicians, artists, photographers, and others. The following people deserve a note of thanks: Rich Batiuk and Carin Bisland, USEPA-Chesapeake Bay Program Office; Ed Pendleton and Kathryn Reshetiloff, USFWS; Vincent Pito, MD-DNR; and Christina Pompa, CBF.

We are especially grateful to the dedicated VIMS personnel who contributed greatly to the production of this report: Leah Nagey, Krisna Davis, Carol Hayes, and Susan Bogardy for their tremendous assistance and perseverance in digitizing the SAV maps, editing the digital data files, mapping ground truth information, and for their constant, careful efforts to maintain high quality control; Ellen F. McLean for assistance in edgematching ARC/INFO SAV maps and for producing SAV section plots for this report; Wanda Cohen, Kay Stubblefield, and Sylvia Motley of the VIMS Publications Center for report production services; and Pat Hall for computer services.

Air Photographics, Inc. conducted the aerial photographic missions and was responsible for the high quality aerial photographs.

SAV SPECIES

The term "submerged aquatic vegetation" for the purpose of this report encompasses 20 taxa from 10 vascular macrophyte families and 3 taxa from 1 freshwater macrophytic algal family, the Characeae, but excludes all other algae, both benthic and planktonic, which occur in Chesapeake Bay and its tributaries (Appendix A). Although these other algae do constitute a portion of the SAV biomass in Chesapeake Bay and its tributaries (Humm, 1979), this study has not attempted to identify, delineate, or discuss the algal component of the vegetation nor its relative importance in the flora, except for the Characeae. This is the case, for example, with the benthic marine algae, including many macrophytes, which sometimes co-occur in the same beds as vascular plants, even as epiphytes on vascular plants.

Ten species of submerged aquatic vegetation are commonly found in the Chesapeake Bay and its tributaries. *Zostera marina* (eelgrass) is dominant in the lower reaches of the bay. *Myriophyllum spicatum* (Eurasian watermilfoil), *Potamogeton pectinatus* (sago pondweed), *Potamogeton perfoliatus* (redhead grass), *Zannichellia palustris* (horned pondweed), *Vallisneria americana* (wild celery), *Elodea canadensis* (common elodea), *Ceratophyllum demersum* (coontail), and *Najas guadalupensis* (southern naiad) are less tolerant of high salinities and are found in the middle and upper reaches of the bay (Stevenson and Confer, 1978; Orth et al., 1979; Orth and Moore, 1981, 1983). *Ruppia maritima* (widgeon grass) is tolerant of a wide range of salinities and is found from the bay mouth to the Susquehanna Flats. Approximately 13 other species are only occasionally found. When present, these species occur primarily in the middle and upper reaches of the bay and the tidal rivers (Appendix A). *Hydrilla verticillata* (hydrilla), a recently introduced species, presently dominates SAV beds in the tidal freshwater reaches of the Potomac River. It has also been reported again in 1993 in the Susquehanna River and Flats where its growth has not been as widespread as in the Potomac River (Kollar, pers. comm.).

Zostera marina and *R. maritima* are the dominant species reported from Chincoteague Bay.

METHODS

INTRODUCTION

Black and white aerial photography at a scale of 1:24,000 was the principal source of information used to assess distribution and abundance of SAV in Chesapeake Bay, its tributaries, and Chincoteague Bay in 1993. There were 1,500 photographs from 137 flight lines which were carefully examined to identify all SAV beds visible on the photography. Outlines of SAV beds were subsequently drawn onto USGS 7.5 minute quadrangles and then digitized, which provided a digital data base for analysis of bed areas and locations. Ground survey information collected in 1993 was tabulated, placed onto the same 7.5 minute quadrangles, and entered into the SAV digital data base.

AERIAL PHOTOGRAPHY

The 1993 SAV aerial photography was obtained by Air Photographics (Martinsburg, West Virginia) using a Wild RC-20 camera with a 153 mm (6 inch) focal length Aviogon lens and Agfa Pan 200 film. The camera was mounted in the bottom fuselage of Air Photographics' Piper Aztec, a twin engine reconnaissance aircraft. Photography was acquired at an altitude of approximately 12,000 feet, which yielded 1:24,000 scale photographs.

Flight lines to obtain the photography were predetermined by Air Photographics to include all areas known to have SAV, as well as those areas which could potentially have SAV (i.e. all areas where water depths were less than 2 m at mean low water). There were 137 flight lines covering 1,764 miles of shoreline and yielding 1,500 exposures. Flight lines included land features that were necessary to establish control points for accurate mapping (Figure 4). Sections of the upper Rappahannock, upper York, and most of the James rivers were not photographed for analysis because of prior determination of the continued absence of SAV in these areas.

Flight lines were prioritized by sections. Flights were timed to occur during the peak growing season of species known to occur in the sections. In addition, specific areas with significant SAV coverage were given priority. Dates of photography for each quadrangle are noted on each map in Appendix C.

General guidelines followed during acquisition of aerial photography (Table 1) address tidal stage, plant growth, sun elevation, water and atmospheric transparency, turbidity, wind, sensor operation, and plotting. Adherence to these guidelines assured acquisition of photography under nearly optimal conditions for detection of SAV, thus insuring accurate photointerpretation.




Figure 4. Map of Chesapeake Bay, its tributaries, and of Chincoteague Bay with approximate locations of flight lines for 1993 SAV photography.

TABLE 1**Guidelines Followed During Acquisition of Aerial Photographs.**

1. Tidal Stage - Photography was acquired at low tide, +/- 0-1.5 ft., as predicted by the National Ocean Survey tables.
2. Plant Growth - Imagery was acquired when growth stages ensured maximum delineation of SAV, and when phenologic stage overlap was greatest.
3. Sun Angle - Photography was acquired when surface reflection from sun glint did not cover more than 30 percent of frame. Sun angle was generally between 20° and 40° to minimize water surface glitter. At least 60 percent line overlap and 20 percent side lap was used to minimize image degradation due to sun glint.
4. Turbidity - Photography was acquired when clarity of water ensured complete delineation of grass beds. This was visually determined from the airplane to insure that SAV could be seen by the observer.
5. Wind - Photography was acquired during periods of no or low wind. Off-shore winds were preferred over on-shore winds when wind conditions could not be avoided.
6. Atmospherics - Photography was acquired during periods of no or low haze and/or clouds below aircraft. There could be no more than scattered or thin broken clouds, or thin overcast above aircraft, to ensure maximum SAV to bottom contrast.
7. Sensor Operation - Photography was acquired in the vertical mode with less than 5 degrees tilt. Scale/altitude/film/focal length combination permitted resolution and identification of one square meter area of SAV (at the surface).
8. Plotting - Each flight line included sufficient identifiable land area to assure accurate plotting of grass beds.

Deviation from any of these guidelines required prior approval by VIMS staff. Quality assurance and calibration procedures were consistently followed. The altimeter was calibrated annually by the Federal Aviation Administration. Camera settings were selected by automatic exposure control. Sun angle was measured with a sensor on the plane. Flight lines were plotted on 1:250,000 scale maps to allow for overlap of photography. To minimize image degradation due to sun glint, the camera was equipped with a computer controlled intervalometer which established 60% line overlap and 20% sidelap. An automatic bubble level held the camera to within one degree tilt. The scale/altitude/film/focal length combination was coordinated so that SAV patches of one square meter could be resolved. Wind speed was monitored hourly. Under normal operating conditions, flights were usually conducted under wind speeds less than 10 mph. Above this speed, wind-generated waves stir bottom sediments which can easily obscure SAV beds in less than one hour. The pilot used experiential knowledge to determine what acceptable level of turbidity would allow complete delineation of SAV beds. During optimum flight conditions the pilot was able to distinguish bottom features such as SAV or algae at low tide. Excessively turbid conditions precluded photography. Determination of optimum cloud cover level was based on pilot experience. Records of this parameter were kept in a flight notebook. Every attempt was made to acquire photographs when there was no cloud cover below 12,000 feet. Cloud cover did not exceed 5% of the area covered by the camera frame. A thin haze layer above 12,000 feet was generally acceptable. Experience with the Chesapeake Bay has shown that optimal atmospheric conditions generally occur two to three days following passage of a cold front, when winds have shifted from north-northwest to south and have moderated to less than 10 mph. Within the guidelines given for prioritizing and executing the photography, the flights were planned to coincide with these atmospheric conditions where possible.

All film was processed by Air Photographics. A 9 inch x 9 inch black and white contact print was produced for each exposed frame. Each photograph was labeled with the date of acquisition as well as flight line number. Film and photographs were stored under appropriate environmental conditions to prevent degradation.

MAPPING PROCESS

For this analysis USGS 7.5 minute quadrangle maps were utilized for mapping SAV beds from aerial photography, for digitizing the SAV beds, and for compiling SAV bed area measurements. Figure 5 gives locations of 179 quadrangles in the study area which includes all regions with potential for SAV growth. Most quadrangles are sequentially numbered for efficient access to data. The name corresponding to each quadrangle in Figure 5 is listed in Table 2. Identification and delineation of SAV beds by photointerpretation utilized all available information including: knowledge of aquatic grass signatures on film, distribution of SAV in 1993 from aerial photography, 1993 ground truth information, and aerial site surveys. USGS 7.5 minute quadrangle maps (1:24,000 scale) printed by




Figure 5. Location of USGS 7.5 minute quadrangles in Chesapeake Bay, its tributaries, and in Chincoteague Bay with corresponding code numbers. (See Table 2 for quad names.)

TABLE 2

List of USGS 7.5 Minute Quadrangles for Chesapeake Bay and Chincoteague Bay SAV Study Areas with Corresponding Code Numbers. (See Figure 5 for Location of Quadrangles. ARC/INFO Generated 7.5 Minute Quadrangles with SAV Beds and Ground truthing Are Reproduced in Appendix C.)

001. Conowingo Dam, Md.-Pa.	034. Alexandria, Va.-D.C.-Md.
002. Aberdeen, Md.	035. Deale, Md.
003. Havre de Grace, Md.	036. Claiborne, Md.
004. North East, Md.	037. St. Michaels, Md.
005. Elkton, Md.-Del.	038. Easton, Md.
006. White Marsh, Md.	039. Fort Belvoir, Va.-Md.
007. Edgewood, Md.	040. Mt. Vernon, Md.-Va.
008. Perryman, Md.	041. Lower Marlboro, Md.
009. Spesutie, Md.	042. North Beach, Md.
010. Earleville, Md.	043. Tilghman, Md.
011. Cecilton, Md.	044. Oxford, Md.
012. Baltimore East, Md.	045. Trappe, Md.
013. Middle River, Md.	046. Preston, Md.
014. Gunpowder Neck, Md.	047. Quantico, Va.-Md.
015. Hanesville, Md.	048. Indian Head, Va.-Md.
016. Betterton, Md.	049. Benedict, Md.
017. Galena, Md.	050. Prince Frederick, Md.
018. Curtis Bay, Md.	051. Hudson, Md.
019. Sparrows Point, Md.	052. Church Creek, Md.
020. Swan Point, Md.	053. Cambridge, Md.
021. Rock Hall, Md.	054. East New Market, Md.
022. Chestertown, Md.	055. Widewater, Va.-Md.
023. Round Bay, Md.	056. Nanjemoy, Md.
024. Gibson Island, Md.	057. Mathias Point, Md.-Va.
025. Love Point, Md.	058. Popes Creek, Md.
026. Langford Creek, Md.	059. Mechanicsville, Md.
027. Centreville, Md.	060. Broomes Island, Md.
028. Washington West, Md.-D.C.-Va.	061. Cove Point, Md.
029. Washington East, D.C.-Md.	062. Taylors Island, Md.
030. South River, Md.	063. Golden Hill, Md.
031. Annapolis, Md.	064. Passapatanzy, Md.-Va.
032. Kent Island, Md.	065. King George, Va.-Md.
033. Queenstown, Md.	066. Dahlgren, Va.-Md.

TABLE 2 (continued)

067. Colonial Beach North, Md.-Va.	106. Reedville, Va.
068. Rock Point, Md.	107. Tangier Island, Va.
069. Leonardtown, Md.	108. Chesconessex, Va.
070. Hollywood, Md.	109. Parksley, Va.
071. Solomons Island, Md.	110. Urbanna, Va.
072. Barren Island, Md.	111. Irvington, Va.
073. Honga, Md.	112. Fleets Bay, Va.
074. Wingate, Md.	113. Nandua Creek, Va.
075. Nanticoke, Md.	114. Pungoteague, Va.
076. Colonial Beach South, Va.-Md.	115. West Point, Va.
077. Stratford Hall, Va.-Md.	116. Saluda, Va.
078. St. Clements Island, Va.-Md.	117. Wilton, Va.
079. Piney Point, Md.-Va.	118. Deltaville, Va.
080. St. Marys City, Md.	119. Jamesville, Va.
081. Point No Point, Md.	120. Toano, Va.
082. Richland Point, Md.	121. Gressitt, Va.
083. Bloodsworth Island, Md.	122. Ware Neck, Va.
084. Deal Island, Md.	123. Mathews, Va.
085. Monie, Md.	124. Franktown, Va.
086. Champlain, Va.	125. Westover, Va.
087. Machodoc, Va.	126. Charles City, Va.
088. Kinsale, Va.-Md.	127. Brandon, Va.
089. St. George Island, Va.-Md.	128. Norge, Va.
090. Point Lookout, Md.	129. Williamsburg, Va.
091. Kedges Straits, Md.	130. Clay Bank, Va.
092. Terrapin Sand Point, Md.	131. Achilles, Va.
093. Marion, Md.	132. New Point Comfort, Va.
094. Mount Landing, Va.	133. Cape Charles, Va.
095. Tappahannock, Va.	134. Cheriton, Va.
096. Lottsburg, Va.	135. Savedge, Va.
097. Heathsville, Va.-Md.	136. Claremont, Va.
098. Burgess, Va.-Md.	137. Surry, Va.
099. Ewell, Md.-Va.	138. Hog Island, Va.
100. Great Fox Island, Va.-Md.	139. Yorktown, Va.
101. Crisfield, Md.-Va.	140. Poquoson West, Va.
102. Saxis, Va.-Md.	141. Poquoson East, Va.
103. Dunnsville, Va.	142. Elliotts Creek, Va.
104. Morattico, Va.	143. Townsend, Va.
105. Lively, Va.	144. Bacons Castle, Va.

TABLE 2 (concluded)

- | | |
|------------------------------|-------------------------------------|
| 145. Mulberry Island, Va. | 163. Mardela Springs, Md. |
| 146. Newport News North, Va. | 164. Wetipquin, Md. |
| 147. Hampton, Va. | 165. Selbyville, Md. |
| 148. Benns Church, Va. | 166. Assawoman Bay, Md. |
| 149. Newport News South, Va. | 167. Berlin, Md |
| 150. Norfolk North, Va. | 168. Ocean City, Md. |
| 151. Little Creek, Va. | 169. Public Landing, Md. |
| 152. Cape Henry, Va. | 170. Tingles Island, Md. |
| 153. Chuckatuck, Va. | 171. Girdle Tree, Md.-Va. |
| 154. Bowers Hill, Va. | 172. Boxiron, Md.-Va. |
| 155. Norfolk South, Va. | 173. Whittington Point, Md.-Va. |
| 156. Kempsville, Va. | 174. Chincoteague West, Va. |
| 157. Princess Anne, Va. | 175. Chincoteague East, Va. |
| 158. Wye Mills, Md. | 176. Anacostia, D.C.-Md. |
| 159. Bristol, Md. | 177. East of New Point Comfort, Va. |
| 160. Fowling Creek, Md. | 178. Bethel Beach, Va. |
| 161. Port Tobacco, Md. | 179. Goose Island, Va. |
| 162. Charlotte Hall, Md. | |

SAV

the Mid-Continent Mapping Center of the National Cartographic Information Center on stable transparent mylar were used as base maps. Distortion-free, identical copies of these base maps were made at the same scale on stable transparent mylar using a contact diazo process.


SAV beds from the 1993 aerial photographs were then mapped onto these diazo mylar copies of USGS 7.5 minute quadrangles. Delineation of each SAV bed was facilitated by superimposing the photographic print with the appropriate diazo mylar quadrangle on a light table. SAV bed boundaries were then traced directly onto the diazo mylar quadrangle with a pencil. Where minor scale differences were evident between a photograph and a quadrangle, or where significant shoreline erosion or accretion had occurred since USGS publication of a map, either a best fit was obtained or shoreline changes were noted on the quadrangle.

In addition to delineating SAV bed boundaries, an estimate of SAV density within each bed was made by visually comparing each bed to an enlarged Crown Density Scale similar to those developed for estimating forest tree crown cover from aerial photography (Figure 6). Bed density was categorized into one of four classes based on a subjective comparison with the density scale. These were: 1, very sparse (<10% coverage); 2, sparse (10-40%); 3, moderate (40-70%); or 4, dense (70-100%). Either the entire bed or subsections within the bed were assigned a bed density number (1 to 4) corresponding to the above density classes. Some beds were subsectioned to delineate where variations in SAV density occurred. Additionally, each distinct SAV unit (bed or bed subsection) was assigned an identifying two letter designation unique to its map. Subsections were further identified as contiguous beds by the addition of two letters unique to that sequence. These contiguous bed identifications aid the tracking and analysis of single natural bed units that were subsectioned due to variation in SAV density. Coupled with the appropriate SAV map number and year of photography, these two letter designations uniquely identify each SAV bed in the data base.

SAV PERIMETER DIGITIZATION AND QUALITY ASSURANCE PROCEDURES

The perimeters of all SAV beds mapped from the aerial photography onto the diazo mylar copies were digitized in ARC/INFO, using an Altek Model 41 tablet, with a resolution of .001 inches (.00254 cm) and an accuracy of .005 inches (.0127 cm). The beds for each quadrangle were digitized twice into two separate ARC/INFO coverages. Each coverage was plotted at an exact scale of 1:24,000 on translucent plotter paper and overlaid on the original mylar for visual checking. In instances where the digitized SAV bed boundaries did not correspond to the original, the bed was re-digitized. Once the SAV outlines on both coverages passed visual inspection, a bed-by-bed comparison of the areas (sq. meters) was made as an additional quality assurance check. Beds were rejected and re-digitized if they were larger than 1 hectare and there was a difference of greater than 0.1% area between the two coverages. The bed-by-bed comparison was useful in identifying instances where SAV beds were incorrectly labelled, thus eliminating coding errors.

DENSITY CLASS



-1-
VERY SPARSE
0-10%

-2-
SPARSE
10-40%

-3-
MODERATE
40-70%

-4-
DENSE
70-100%

PERCENT CROWN COVER

Figure 6. Crown density scale used for estimating density of SAV beds from aerial photography. (Rows of squares with black and white patterns represent three different arrangements of vegetated cover for a given percentage.)

SAV

Prior to each digitization session, the Altek instrument was checked manually against a digitizing standard. This was accomplished by first securing a diazo mylar quadrangle with SAV polygons to the digitizing tablet. Then the mylar standard was secured to the same quadrangle and digitized. The digitized area of each standard was compared to the known area of the standard. If a variation between the known and the mean of the observed areas exceeded 1.0%, the maps were redigitized. In addition, visual checks were made with respect to the absolute location of the digitizing standard as secured to the map.

Maximum accuracy was maintained by exclusively using mylar quadrangles and standards rather than paper ones, which can change scale as a function of changes in air temperature and humidity in the digitizer room.

Standard operating procedures (SOPs) were developed to facilitate orderly and efficient processing of the 1993 SAV maps and the SAV computer files produced from them, and to comply with the need for consistency, quality assurance, and quality control. SOPs developed include: a detailed procedure for digitization of SAV maps; a digitizer log in which all operations were recorded and dated, which was used to guide and record editing operations; and a flow chart used to track progress of all operations.

CALCULATION OF 1993 SAV AREAS

The SAV coverages in UTM ARC/INFO Zone 18 format were used to calculate area in square meters for all SAV beds. These areas are reported as USGS 7.5 minute quadrangle section, and zone totals in the tables in the Results section. Section and zone totals were calculated by using an overlay operation of the polygons on the SAV beds in ARC/INFO. The definition of the sections used in this analysis are provided in Table 3.

ORGANIZATIONAL PROCEDURES FOR ANALYSIS AND DISCUSSION

Discussion of the distribution of SAV in Chesapeake Bay and tributaries has been organized into three zones as established by Orth and Moore (1982) and modified by Orth et al., (1989) (Figure 7). The Lower Bay zone is the area from the entrance of the bay to a line originating from Smith Point at the mouth of the Potomac River, extending to approximately 3 nautical miles south of Tangier Island, then extending to just below the Little Annemessex River mouth. From this line north to the Chesapeake Bay Bridge at Kent Island is the area referred to as the Middle Bay zone. The area between the Chesapeake Bay Bridge and the Susquehanna Flats is referred to as the Upper Bay zone.

TABLE 3**Area Descriptions for Each of the 21 Sections of the Chesapeake Bay SAV Study Area.**

- Section 1. Susquehanna Flats - all areas between and including Spesutie Island and Turkey Point at the mouth of the Elk River to include the Northeast River.
- Section 2. Upper Eastern Shore - all areas in the Elk, Bohemia, and Sassafras rivers, and areas on the eastern shore above the Swan Point quadrangle.
- Section 3. Upper Western Shore - all areas south of Spesutie Island and north of the Chesapeake Bay Bridge to include the Bush, Gunpowder, Middle, Patapsco, and Magothy rivers.
- Section 4. Chester River - includes all of the Chester River, Eastern Neck, and areas north of the Chesapeake Bay Bridge on Kent Island extending to north of Swan Point.
- Section 5. Central Western Shore - all areas south of the Chesapeake Bay Bridge and north of Holland Point on Herring Bay to include the Severn, Rhode, South, and West rivers and Herring Bay.
- Section 6. Eastern Bay - all areas south of the Chesapeake Bay Bridge on Kent Island and north of Tilghman Island from Green Marsh Point to include the Wye, East, and Miles rivers, Crab Alley and Prospect bays, and Poplar, Jefferson, and Coaches islands.
- Section 7. Choptank River - all areas south of Tilghman Island from Green Marsh Point and north of Taylor Island to include the Choptank and Little Choptank rivers.
- Section 8. Patuxent River - all areas in the Patuxent River.
- Section 9. Middle Western Shore - all areas south of Holland Point at Herring Bay and north of Point Lookout on the Potomac River not including the mouth of the Patuxent River.

(continue on next page)

TABLE 3 (continued)

- Section 10. Lower Potomac River - all areas between the mouth of the Potomac River to a line extending from Maryland Point on the north shore, just above Nanjemoy Creek, to Somerset Beach on the south shore.
- Section 11. Upper Potomac River - all areas upstream of the Lower Potomac River Section to Chain Bridge at Washington D.C.
- Section 12.** Middle Eastern Shore - all areas south of Taylor Island and north of a line bisecting Cedar Island to include the Big and Little Annemessex, Honga, Nanticoke, Wicomico, and Manokin rivers, and Fishing Bay.
- Section 13.** Mid-Bay Island Complex - all areas in and adjacent to Bloodsworth, South Marsh, Smith, and Tangier islands.
- Section 14.** Lower Eastern Shore - all areas south of a line bisecting Cedar Island and located just above the Maryland-Virginia border to Fisherman's Island.
- Section 15. Reedville Region - includes the area between Windmill Point on the Rappahannock River and Smith Point at the mouth of the Potomac River.
- Section 16. Rappahannock River Complex - includes the entire Rappahannock and Piankatank rivers, and the Milford Haven area.
- Section 17. New Point Comfort Region - includes the area from New Point Comfort Lighthouse north to Garden Creek just south of Milford Haven.
- Section 18.** Mobjack Bay Complex - includes the East, North, Ware, and Severn rivers, the north shore of Mobjack Bay from New Point Comfort Lighthouse to the North River, and north of a line bisecting the large shoal area around the Guinea Marshes.

(continue on next page)

TABLE 3 (concluded)

Section 19.** York River - all areas of the York River from north of the Poropotank River to the mouth, including south of a line bisecting the large shoal area around the Guinea Marshes and the north shore of Goodwin Island.

Section 20.** Lower Western Shore - includes all areas south of Goodwin Island to Lynnhaven Inlet, including Broad Bay but not including the James River.

Section 21. James River - all areas in the James River including the Chickahominy River.

** Sections 12, 13, 14, 18, 19, and 20 were given new boundaries for the 1987 report (Orth et al., 1989) which also changed the delineation of the three zones. These new boundaries were retained for the 1989, 1990, 1991, and 1992 reports (Orth and Nowak, 1990; Orth et al., 1991; Orth et al., 1992, Orth et al., 1993) and for this report. (Refer to Figure 7 and Appendix B for boundary locations.)

UPPER

MIDDLE

LOWER




Figure 7. Location of Chincoteague Bay and Chesapeake Bay with Upper, Middle, and Lower zones and 21 sections used for delineation of SAV distribution patterns. (See Table 3 and Appendix B for exact boundary positions.)

The salinity within each zone roughly coincides with the major salinity zones of estuaries: polyhaline (18-25 ‰), Lower zone; mesohaline (5-18 ‰), Middle zone; oligohaline (0.5-5 ‰), Upper zone. Although the major rivers and smaller tributaries of Chesapeake Bay have their own salinity regimes, the distribution of SAV in each river is discussed within the zone where it connects to the bay.

In addition, 21 sections of the bay are identified (Figure 7, Table 3) for a more detailed discussion of SAV distribution. These sections, which were first delineated for the 1984 SAV survey (Orth et al., 1985) and slightly modified for the 1987 SAV survey (Orth et al., 1989), denote relatively distinct parts of Chesapeake Bay and its tributaries that are readily identifiable. The section boundaries used for analysis and discussion of the 1993 SAV distribution and abundance data were used for the 1987, 1989, 1990, 1991 and 1992 reports (Orth et al., 1989; Orth and Nowak, 1990, Orth et al., 1991, Orth et al., 1992, Orth et al., 1993). Sections 1 through 4 are located in the Upper Bay zone; sections 5 through 13 are in the Middle Bay zone; and sections 14 through 21 are in the Lower Bay zone. SAV distribution in Chincoteague Bay is presented and discussed separately from Chesapeake Bay. Appendix B gives the latitude and longitude, in decimal degrees, of the boundary points of each Chesapeake Bay section and of Chincoteague Bay.

GROUND SURVEYS AND OTHER DATA BASES

Ground surveys were accomplished by cooperative efforts from a number of agencies and individuals. Although not all areas of the bay were surveyed, the data did provide valuable supplemental information. The surveys confirmed the existence of some SAV beds mapped from the 1993 aerial photography, as well as SAV beds not visible from the photography. The surveys also provided species data for many of the SAV beds. Ground survey information supplied to VIMS researchers was included on the SAV distribution and abundance maps reproduced in Appendix C. Each survey was designated by a unique symbol to identify the different methods of sampling. In most cases the symbols on the SAV maps (Appendix C) have been enlarged and offset from the actual sampling point to avoid confusion with the mapped SAV bed. Where species information was available, it was included on the map. Because of space limitations on the maps reproduced in Appendix C, occasionally one or more survey points were combined where the information was duplicated. All ground survey data supplied to VIMS are tabulated in Appendix E.

In Maryland, ground survey data were obtained in 1993 by VIMS, the USGS National Center, U.S. Fish and Wildlife Service, Patuxent River Park staff, and by the Citizens' volunteer survey. The USFWS contributed ground survey data for the Magothy and Chester rivers. The USGS National Center provided ground truth data for the Potomac River. Patuxent River ground survey data were obtained by the Maryland-National Capital Parks and Planning Commission Patuxent River Park staff. The Citizens' volunteer survey, under the guidance of the USFWS and the Chesapeake Bay Foundation (CBF), identified SAV locations and SAV species when possible throughout various areas of the Chesapeake and Chincoteague bays. Volunteers, who were recruited through press releases, newsletters, and personal letters, were provided with a SAV identification guide, reduced 1992 SAV maps to aid in the location of SAV beds, and data sheets for reporting visits to numerous sites around the bays. USFWS staff mapped the data on copies of 1992 SAV distribution maps (USGS 7.5 minute quads with 1992 SAV beds). These maps were supplied to VIMS SAV researchers and transferred to the 1993 SAV distribution maps reproduced in Appendix C. Data from the USFWS, Patuxent River Park staff, and the Citizens' surveys were compiled and tabulated by USFWS. This table became the basis of the much expanded table published in Appendix E.

One 1993 SAV project being conducted on the Susquehanna Flats by Stan Kollar of Harford Community College, Maryland, provided data in the form of species presence by percentage.

For those areas in Virginia waters where aerial photographic evidence of SAV beds was inconclusive, photoverification was accomplished by ground truth surveys. Observations were principally made from small boats and by divers snorkeling over areas indicated from the photographs. In the York, Piankatank, and Rappahannock rivers, where VIMS researchers had transplanted SAV (principally eelgrass), transplant sites were also examined carefully by divers for any extant SAV. VIMS

scientists also surveyed a number of sites in the Chesapeake Bay as part of an intensive quantitative SAV study (VIMS, unpublished data). Data for Virginia waters were also collected by the Citizens' volunteer survey (compiled by the USFWS). In addition, a great deal of ground survey information could be extrapolated from earlier studies (Orth et al., 1979; Orth and Moore, 1982). SAV beds in the lower bay contain primarily one or two species and most areas have not undergone wide fluctuations in distribution and abundance since the first bay-wide survey in 1978.

Ground survey data from all sources reported here were added to the USFWS table and each SAV siting was cross-referenced with its associated 1993 SAV bed location. This expanded ground survey table is presented in Appendix E.

RESULTS

DATA PRESENTATION

SAV distribution data are presented by quadrangle (Table 4), by section and zone (Table 5), by quadrangles within a section (Table 6), and by density class for each section (Table 7). Quadrangle maps annotated with all SAV beds are presented in Appendix C, while individual bed areas for each quadrangle are given in Appendix D. Appendix E tabulates all ground truth data for 1993. The 1993 SAV distribution data and species occurrences are first discussed relative to the Upper, Middle, and Lower Bay zones. The 21 sections of the Chesapeake Bay, and Chincoteague Bay, are then discussed individually, and the data compared to results from the 1992 survey of SAV distribution and abundance (Orth, et al., 1993). SAV is plotted for each section and for Chincoteague Bay in Figures 8 through 36. SAV beds are plotted in red, and bold, black lines represent section boundaries. USGS 7.5 minute quadrangles are represented by a grid of numbered rectangles (refer to Table 2 for quadrangle names listed by map number). Specific names of rivers, creeks, or points of land which are not found on the section plots, are on the quadrangle maps for that section.

1993 SUMMARY

In 1993, the Chesapeake Bay had 29,589 hectares of SAV, compared to 28,591 hectares in 1992, with 2,700 hectares (9.0%), 13,901 hectares (47.0%), and 13,018 hectares (44.0%) occurring in the Upper, Middle, and Lower Bay zones, respectively (Figures 1, 2, and 3). SAV generally increased in each of the three zones from 1992. Decreases in some sections (e.g. Mid-Bay Island Complex, Upper Eastern Shore, and Western Shore) were offset by larger increases in other sections (e.g. Choptank River, Eastern Bay, and Chester River sections). SAV increased in abundance in all sections in the Lower Bay zone.

Upper Bay Zone

In 1993, in the Upper Bay zone, 66.5% (1,777 hectares) of the SAV was located in the Susquehanna Flats (Section 1). Overall abundance and density of SAV was similar to the 1992 level (1,792 hectares). In the Flats, 85.4% of all SAV beds were classified as very sparse in 1993 (0-10% coverage), while 7.7% of beds were classified as dense (70-100% coverage) (Table 7; Figure 3). *Myriophyllum spicatum*, *Heteranthera dubia*, *V. americana*, *H. verticillata*, *C. demersum*, and *N. guadalupensis* were the six species reported. In the Upper Eastern Shore (Section 2) there were 184 hectares of SAV in 1993 (99 hectares less than in 1992) located principally in the Elk and lower Sassafras rivers, with *M. spicatum* and *V. americana* found most frequently, especially in the Elk River. Much of the difference from 1992 in the Upper Eastern Shore section was recorded from the

TABLE 4

Total Area of SAV in Hectares by USGS 7.5 Minute Quadrangles for 1992 and 1993.

QUADRANGLE	1992	1993
001. Conowingo Dam, Md.-Pa.	0	0
002. Aberdeen, Md.	14.98	8.21
003. Havre de Grace, Md.	1,745.68	1,734.74
004. North East, Md.	126.21	46.18
005. Elkton, Md.-Del.	0	0
006. White Marsh, Md.	0	0
007. Edgewood, Md.	.43	0
008. Perryman, Md.	8.78	8.03
009. Spesutie, Md.	45.08	46.32
010. Earleville, Md.	116.16	53.43
011. Cecilton, Md.	0	0
012. Baltimore East, Md.	0	0
013. Middle River, Md.	16.07	5.47
014. Gunpowder Neck, Md.	155.87	47.81
015. Hanesville, Md.	26.19	4.24
016. Betterton, Md.	2.47	68.99
017. Galena, Md.	2.98	4.48
018. Curtis Bay, Md.	0	0
019. Sparrows Point, Md.	#	0
020. Swan Point, Md.	5.39	17.62
021. Rock Hall, Md.	12.34	32.63
022. Chestertown, Md.	0	0
023. Round Bay, Md.	#	#
024. Gibson Island, Md.	#	13.21
025. Love Point, Md.	0	0
026. Langford Creek, Md.	220.66	518.08
027. Centreville, Md.	0	0
028. Washington West, Md.-D.C.	9.92	25.22
029. Washington East, D.C.-Md.	0	.75
030. South River, Md.	#	#
031. Annapolis, Md.	0	#
032. Kent Island, Md.	69.59	154.37
033. Queenstown, Md.	87.40	181.41

(continue on next page)

TABLE 4 (continued)

QUADRANGLE	1992	1993
034. Alexandria, Va.-D.C.-Md.	318.29	336.46
035. Deale, Md.	#	0
036. Claiborne, Md.	231.64	426.61
037. St. Michaels, Md.	243.63	270.70
038. Easton, Md.	0	0
039. Fort Belvoir, Va.-Md.	133.72	111.39
040. Mt. Vernon, Va.-Md.	254.57	236.45
041. Lower Marlboro, Md.	#	8.78
042. North Beach, Md.	0	0
043. Tilghman, Md.	222.47	393.98
044. Oxford, Md.	115.79	444.33
045. Trappe, Md.	0	#
046. Preston, Md.	#	0
047. Quantico, Va.-Md.	594.92	599.93
048. Indian Head, Md.- Va.	336.04	346.05
049. Benedict, Md.	0	#
050. Prince Frederick, Md.	-	-
051. Hudson, Md.	515.96	567.35
052. Church Creek, Md.	105.79	149.08
053. Cambridge, Md.	5.66	4.02
054. East New Market, Md.	0	0
055. Widewater, Va.-Md.	730.95	623.44
056. Nanjemoy, Md.	168.32	89.01
057. Mathias Point, Md.-Va.	292.05	252.84
058. Popes Creek, Md.	1.30	1.52
059. Mechanicsville, Md.	0	0
060. Broomes Island, Md.	#	#
061. Cove Point, Md.	#	#
062. Taylors Island, Md.	62.39	100.19
063. Golden Hill, Md.	29.23	65.10
064. Passapatanzy, Md.-Va.	12.24	6.60
065. King George, Va.-Md.	74.45	79.07
066. Dahlgren, Va.-Md.	33.98	28.57
067. Colonial Beach North, Va.	47.76	49.80
068. Rock Point, Md.	0	#

(continue on next page)

TABLE 4 (continued)

QUADRANGLE	1992	1993
069. Leonardtown, Md.	0	0
070. Hollywood, Md.	0	#
071. Solomons Island, Md.	#	.99
072. Barren Island, Md.	433.61	205.83
073. Honga, Md.	1,326.88	1,340.42
074. Wingate, Md.	480.81	540.89
075. Nanticoke, Md.	0	0
076. Colonial Beach South, Va.	0	#
077. Stratford Hall, Va.-Md.	0	0
078. St. Clements Island, Va.	#	#
079. Piney Point, Md.-Va.	0	0
080. St. Mary's City, Md.	8.81	12.26
081. Point No Point, Md.	-	-
082. Richland Point, Md.	45.90	41.06
083. Bloodsworth Island, Md.	1,024.10	863.08
084. Deal Island, Md.	68.75	77.09
085. Monie, Md.	0	7.09
086. Champlain, Va.	-	-
087. Machodoc, Va.	0	0
088. Kinsale, Va.-Md.	0	0
089. St. George Island, Md.-Va	3.08	4.26
090. Point Lookout, Md.	0	0
091. Kedges Straits, Md.	971.21	903.44
092. Terrapin Sand Point, Md.	267.81	231.38
093. Marion, Md.	278.43	293.94
094. Mount Landing, Va	-	-
095. Tappahannock, Va	-	-
096. Lottsburg, Va.	0	0
097. Heathsville, Va.-Md.	0	0
098. Burgess, Va.-Md.	#	0
099. Ewell, Md.-Va.	2,543.16	2,376.65
100. Great Fox Island, Md.-Va.	1,504.94	1,483.11
101. Crisfield, Md.-Va.	321.69	339.28
102. Saxis, Va.-Md.	2.86	2.10
103. Dunnsville, Va.	-	-

(continue on next page)

TABLE 4 (continued)

QUADRANGLE	1992	1993
104. Morattico, Va.	0	0
105. Lively, Va.	0	0
106. Reedville, Va.	302.51	304.16
107. Tangier Island, Va.	601.73	571.70
108. Chesconessex, Va.	1,042.80	1,103.95
109. Parksley, Va.	461.99	510.57
110. Urbanna, Va.	11.25	0
111. Irvington, Va.	165.60	159.94
112. Fleets Bay, Va.	475.89	508.92
113. Nandua Creek, Va.	473.91	467.61
114. Pungoteague, Va.	949.27	1,008.32
115. West Point, Va.	-	-
116. Saluda, Va.	0	0
117. Wilton, Va.	18.18	44.09
118. Deltaville, Va.	142.86	216.84
119. Jamesville, Va.	634.02	683.82
120. Toano, Va.	-	-
121. Gressitt, Va.	-	-
122. Ware Neck, Va.	318.37	313.09
123. Mathews, Va.	326.70	395.88
124. Franktown, Va.	718.67	767.84
125. Westover, Va.	-	-
126. Charles City, Va.	-	-
127. Brandon, Va.	-	-
128. Norge, Va.	-	-
129. Williamsburg, Va.	-	-
130. Clay Bank, Va.	#	0
131. Achilles, Va.	1,040.46	1,057.83
132. New Point Comfort, Va.	1,486.00	1,503.15
133. Cape Charles, Va.	361.03	465.78
134. Cheriton, Va.	87.25	96.82
135. Savedge, Va.	-	-
136. Claremont, Va.	-	-
137. Surry, Va.	-	-

(continue on next page)

TABLE 4 (continued)

QUADRANGLE	1992	1993
138. Hog Island, Va.	-	-
139. Yorktown, Va.	1.16	2.52
140. Poquoson West, Va.	582.94	618.48
141. Poquoson East, Va.	1,161.06	1,181.72
142. Elliotts Creek, Va.	111.96	113.63
143. Townsend, Va.	0	-
144. Bacons Castle, Va.	-	-
145. Mulberry Island, Va.	-	-
146. Newport News North, Va.	-	-
147. Hampton, Va.	380.63	366.99
148. Benns Church, Va.	-	-
149. Newport News South, Va.	-	0
150. Norfolk North, Va.	-	-
151. Little Creek, Va.	0	0
152. Cape Henry, Va.	19.55	21.22
153. Chuckatuck, Va.	-	-
154. Bowers Hill, Va.	-	-
155. Norfolk South, Va.	-	-
156. Kempsville, Va.	-	-
157. Princess Anne, Va.	0	-
158. Wye Mills, Md.	0	0
159. Bristol, Md.	#	#
160. Fowling Creek, Md.	0	-
161. Port Tobacco, Md.	12.60	12.21
162. Charlotte Hall, Md.	0	4.60
163. Mardela Springs, Md.	0	0
164. Wetipquin, Md.	0	0
165. Selbyville, Md.	0	0
166. Assawoman Bay, Md.	7.94	20.35
167. Berlin, Md.	10.69	15.36
168. Ocean City, Md.	23.57	33.06
169. Public Landing, Md.	0	0
170. Tingles Island, Md.	1,180.30	1,189.95
171. Girdle Tree, Md.-Va.	0	0

(continue on next page)

TABLE 4 (concluded)

QUADRANGLE	1992	1993
172. Boxiron, Md.-Va.	771.61	817.41
173. Whittington Point, Md.-Va	399.10	451.65
174. Chincoteague West, Va.	6.27	13.92
175. Chincoteague East, Va.	924.70	1,034.71
176. Anacostia, D.C.-Md.	0	0
177. East of New Point Comfort, Va.	8.67	18.55
178. Bethel Beach, Va.	0*	5.75
179. Goose Island, Va.	214.79	177.38
TOTAL SAV - Chesapeake Bay	28,591.23	29,588.68
TOTAL SAV - Chincoteague Bay	3,324.18	3,576.42

- = Indicates quadrangle not photographed and assumed to have no SAV.
- 0 = Indicates quadrangle photographed and no SAV noted.
- 0* = This quadrangle was newly published and was not available to this year's mapping. SAV beds located on this quadrangle were mapped on the overlapping portion of the adjoining quadrangle.
- # = SAV detected by ground truthing only.

Table 5

Number of Hectares of SAV in 1992 and 1993 for the 21 Sections and Three Zones of Chesapeake Bay and for Chincoteague Bay.

ZONE	SECTION	AREA (HECTARES)	
		1992	1993
Upper	1. Susquehanna Flats	1,791.97	1,776.52
	2. Upper Eastern Shore	282.96	184.46
	3. Upper Western Shore	185.97	80.13
	4. Chester River	<u>255.16</u>	<u>628.82</u>
	Zone Total	2,516.06	2,669.93
Middle	5. Central Western Shore	0.00	0.00
	6. Eastern Bay	553.93	737.25
	7. Choptank River	1,085.39	1,894.30
	8. Patuxent River	0.00	9.77
	9. Middle Western Shore	0.00	0.00
	10. Lower Potomac River	571.03	457.78
	11. Upper Potomac River	2,461.96	2,362.65
	12. Middle Eastern Shore	3,046.93	2,972.14
	13. Mid-Bay Island Complex	<u>5,993.93</u>	<u>5,466.78</u>
	Zone Total	13,713.17	13,900.67
Lower	14. Lower Eastern Shore	5,920.17	6,298.94
	15. Reedville	778.40	813.09
	16. Rappahannock River Complex	586.84	722.18
	17. New Point Comfort Region	395.91	430.94
	18. Mobjack Bay Complex	1,818.03	1,842.66
	19. York River	830.08	840.62
	20. Lower Western Shore	2,029.07	2,065.64
21. James River	<u>3.50</u>	<u>4.01</u>	
	Zone Total	12,362.00	13,018.08
Total SAV for Chesapeake Bay		28,591.23	29,588.68
Total SAV for Chincoteague Bay		3,324.18	3,576.42

TABLE 6

Number of Square Meters of SAV in 1993 for Each USGS 7.5 Minute Quadrangle of the 21 Sections of Chesapeake Bay and of Chincoteague Bay. (Map Code Numbers from Table 2 in Parentheses.)

SECTION	QUADRANGLE	AREA
Susquehanna Flats - 1	Conowingo Dam, Md.-Pa. (1)	0.00
	Aberdeen, Md. (2)	82,134.82
	Havre de Grace, Md. (3)	17,347,404.14
	North East, Md. (4)	0.00
	Elkton, Md.-Del. (5)	0.00
	Perryman (8)	11,748.72
	Spesutie, Md. (9)	323,927.89
	Earleville, Md. (10)	0.00
		17,765,216 sq. m
		1,776.52 hectares
	4,389.79 acres	
Upper Eastern Shore - 2	North East, Md. (4)	461,782.17
	Elkton, Md.-Del. (5)	0.00
	Perryman, Md. (8)	0.00
	Spesutie, Md. (9)	71,319.88
	Earleville, Md. (10)	534,344.23
	Cecilton, Md. (11)	0.00
	Gunpowder Neck, Md. (14)	0.00
	Hanesville, Md. (15)	42,409.63
	Betterton, Md. (16)	689,938.61
	Galena, Md. (17)	44,811.70
	Swan Point, Md. (20)	0.00
	Rock Hall, Md. (21)	0.00
	Chestertown, Md. (22)	0.00
	1,844,606 sq. m	
	184.46 hectares	
	455.80 acres	
Upper Western Shore - 3	White Marsh, Md. (6)	0.00

(continue on next page)

Table 6 (continued)

SECTION	QUADRANGLE	AREA
Upper Western Shore - 3 (continued)	Edgewood (7)	0.00
	Perryman, Md. (8)	68,574.17
	Spesutie, Md. (9)	67,955.40
	Baltimore East, Md. (12)	0.00
	Middle River, Md. (13)	54,652.32
	Gunpowder Neck, Md. (14)	478,077.09
	Hanesville, Md. (15)	0.00
	Curtis Bay, Md. (18)	0.00
	Sparrows Point, Md. (19)	0.00
	Swan Point, Md. (20)	0.00
	Round Bay, Md. (23)	0.00
	Gibson Island, Md. (24)	132,052.85
	Love Point, Md. (25)	<u>0.00</u>
		801,312 sq. m
	80.13 hectares	
	198.00 acres	
Chester River - 4	Betterton, Md. (16)	0.00
	Galena, Md. (17)	0.00
	Swan Point, Md. (20)	176,178.84
	Rock Hall, Md. (21)	326,330.99
	Chestertown, Md. (22)	0.00
	Love Point, Md. (25)	0.00
	Langford Creek, Md. (26)	5,180,752.93
	Centreville, Md. (27)	0.00
	Kent Island, Md. (32)	0.00
	Queenstown, Md. (33)	<u>604,969.92</u>
	6,288,233 sq.	
	628.82 hectares	
	1,553.82 acres	
Central Western Shore -5	Curtis Bay, Md. (18)	0.00
	Round Bay, Md. (23)	0.00
	Gibson Island, Md. (24)	0.00
	Love Point, Md. (25)	0.00

(continue on next page)

Table 6 (continued)

SECTION	QUADRANGLE	AREA
Central Western Shore -5 (continued)	South River, Md. (30)	0.00
	Annapolis, Md. (31)	0.00
	Kent Island, Md. (32)	0.00
	Deale, Md. (35)	0.00
	North Beach, Md. (42)	0.00
		<hr/>
		0.00 sq. m
		0.00 hectares
		0.00 acres
Eastern Bay - 6	Centreville, Md. (27)	0.00
	Annapolis, Md. (31)	0.00
	Kent Island, Md. (32)	1,543,673.84
	Queenstown, Md. (33)	1,209,126.25
	Claiborne, Md. (36)	2,842,965.13
	St. Michaels, Md. (37)	1,776,699.18
	Easton, Md. (38)	0.00
	Tilghman, Md. (43)	0.00
	Oxford, Md. (44)	0.00
	Wye Mills, Md. (158)	0.00
		<hr/>
		7,372,464 sq. m
		737.25 hectares
		1,821.74 acres
Choptank River - 7	Centreville, Md. (27)	0.00
	Claiborne, Md. (36)	1,423,106.87
	St. Michaels, Md. (37)	930,320.53
	Easton, Md. (38)	0.00
	Tilghman, Md. (43)	3,939,834.99
	Oxford, Md. (44)	4,443,327.36
	Trappe, Md. (45)	0.00
	Preston, Md. (46)	0.00
	Hudson, Md. (51)	5,673,502.74
	Church Creek, Md. (52)	1,490,785.18
Cambridge, Md. (53)	40,219.67	
East New Market, Md. (54)	0.00	

(continue on next page)

Table 6 (continued)

SECTION	QUADRANGLE	AREA
Choptank River 7- (continued)	Taylor's Island, Md. (62)	1,001,937.99
	Golden Hill, Md. (63)	0.00
	Nanticoke, Md. (75)	0.00
	Wye Mills, Md. (158)	0.00
	Fowling Creek, Md. (160)	0.00
		<hr/>
		18,943,035 sq. m
		1,894.30 hectares
		4,680.83 acres
Patuxent River - 8	Deale, Md. (35)	0.00
	Lower Marlboro, Md. (41)	87,761.58
	North Beach, Md. (42)	0.00
	Benedict, Md. (49)	0.00
	Prince Frederick, Md. (50)	0.00
	Mechanicsville, Md. (59)	0.00
	Broomes Island, Md. (60)	0.00
	Cove Point, Md. (61)	0.00
	Leonardtown, Md. (69)	0.00
	Hollywood, Md. (70)	0.00
	Solomons Island, Md. (71)	9,946.00
	Bristol, Md. (159)	0.00
	Charlotte Hall, Md. (162)	0.00
		<hr/>
		97,708 sq. m
		9.77 hectares
		24.14 acres
Middle Western Shore - 9	North Beach, Md. (42)	0.00
	Prince Frederick, Md. (50)	0.00
	Hudson, Md. (51)	0.00
	Broomes Island, Md. (60)	0.00
	Cove Point, Md. (61)	0.00
	Taylor's Island, Md. (62)	0.00
	Solomons Island, Md. (71)	0.00
	Barren Island, Md. (72)	0.00
St. Mary's City, Md. (80)	0.00	

(continue on next page)

Table 6 (continued)

SECTION	QUADRANGLE	AREA
Middle Western Shore -9 (continued)	Point No Point, Md. (81)	0.00
	Richland Point, Md. (82)	0.00
	Point Lookout, Md. (90)	<u>0.00</u>
		0.00 sq. m 0.00 hectares 0.00 acres
Lower Potomac River - 10	Nanjemoy, Md. (56)	890,060.10
	Mathias Point, Md.-Va. (57)	2,528,396.64
	Popes Creek, Md. (58)	15,239.88
	Mechanicsville, Md. (59)	0.00
	King George, Va.-Md. (65)	149,134.80
	Dahlgren, Va.-Md. (66)	285,711.40
	Colonial Beach North, Va.-Md. (67)	498,035.53
	Rock Point, Md. (68)	0.00
	Leonardtwn, Md. (69)	0.00
	Hollywood, Md. (70)	0.00
	Solomons Island, Md. (71)	00.0
	Colonial Beach South, Va.-Md. (76)	0.00
	Stratford Hall, Va.-Md. (77)	0.00
	St. Clements Island, Va.-Md. (78)	0.00
	Piney Point, Md.-Va. (79)	0.00
	St. Mary's City, Md. (80)	122,586.69
	Champlain, Va. (86)	0.00
	Machodoc, Va. (87)	0.00
	Kinsale, Va.-Md. (88)	0.00
	St. George Island, Md.-Va. (89)	42,625.36
	Point Lookout, Md. (90)	0.00
	Lottsburg, Va. (96)	0.00
	Heathsville, Va.-Md. (97)	0.00
Burgess, Va.-Md. (98)	0.00	
Port Tobacco, Md. (161)	0.00	
Charlotte Hall, Md. (162)	<u>46,044.19</u>	
	4,577,835 sq. m 457.78 hectares 1,131.18 acres	

(continue on next page)

Table 6 (continued)

SECTION	QUADRANGLE	AREA
Upper Potomac River - 11	Washington West, Md.-D.C.-Va. (28)	252,155.37
	Washington East, D.C.-Md. (29)	7,488.02
	Alexandria, VA.-D.C.-Md. (34)	3,364,573.07
	Fort Belvoir, Va.-Md. (39)	1,113,940.17
	Mt. Vernon, Va.-Md. (40)	2,364,516.42
	Quantico, Va.-Md. (47)	5,999,269.62
	Indian Head, Md.- Va. (48)	3,460,515.00
	Widewater, Va.-Md. (55)	6,234,426.54
	Nanjemoy, Md. (56)	0.00
	Mathias Point, Md.-Va. (57)	0.00
	Passapatanzy, Md.-Va. (64)	65,977.47
	King George, Va.-Md. (65)	641,543.50
	Dahlgren, Va.-Md. (66)	0.00
	Port Tobacco, Md. (161)	122,124.20
	Anacostia, D.C.-Md. (176)	0.00
		23,626,529 sq. m
	2,362.65 hectares	
	5,838.11 acres	
Middle Eastern Shore - 12	Taylor's Island, Md. (62)	0.00
	Golden Hill, Md. (63)	651,001.82
	Barren Island, Md. (72)	2,058,286.91
	Honga, Md. (73)	13,404,228.15
	Wingate, Md. (74)	5,408,853.95
	Nanticoke, Md. (75)	0.00
	Point No Point, Md. (81)	0.00
	Richland Point, Md. (82)	410,643.94
	Bloodsworth Island, Md. (83)	1,251,669.10
	Deal Island, Md. (84)	770,886.55
	Monie, Md. (85)	70,868.46
	Terrapin Sand Point, Md. (92)	200,625.97
	Marion, Md. (93)	2,939,413.35
	Great Fox Island, Md.-Va. (100)	1,406,547.54
	Crisfield, Md.-Va. (101)	1,148,377.60
	Mardela Springs, Md. (163)	0.00

(continue on next page)

Table 6 (continued)

SECTION	QUADRANGLE	AREA
Middle Eastern Shore - 12 (continued)	Wetipquin, Md. (164)	0.00
		29,721,403 sq. m
		2,972.14 hectares
		7,344.17 acres
Mid-Bay Island Complex -13	Richland Point, Md. (82)	0.00
	Bloodsworth Island, Md. (83)	7,379,138.35
	Deal Island, Md. (84)	0.00
	Kedges Straits, Md. (91)	9,034,396.14
	Terrapin Sand Point, Md. (92)	2,113,131.00
	Ewell, Md.-Va. (99)	23,766,456.38
	Great Fox Island, Md.-Va. (100)	5,474,556.40
	Tangier Island, Va. (107)	5,126,283.11
	Goose Island, Va. (179)	<u>1,773,825.53</u>
		54,667,787 sq. m
	5,466.78 hectares	
	13,508.43 acres	
Lower Eastern Shore - 14	Marion, Md. (93)	0.00
	Great Fox Island, Md.-Va. (100)	7,949,953.33
	Crisfield, Md.-Va. (101)	2,244,394.61
	Saxis, Va.-Md. (102)	20,989.38
	Tangier Island, Va. (107)	590,668.70
	Chesconessex, Va. (108)	11,039,547.00
	Parksley, Va. (109)	5,105,724.52
	Nandua Creek, Va. (113)	4,676,076.81
	Pungoteague, Va. (114)	10,083,156.89
	Jamesville, Va. (119)	6,838,222.73
	Franktown, Va. (124)	7,678,398.81
	Cape Charles, Va. (133)	4,657,806.42
	Cheriton, Va. (134)	968,198.06
	Elliotts Creek, Va. (142)	1,136,293.00
	Townsend, Va. (143)	0.00

(continue on next page)

Table 6 (continued)

SECTION	QUADRANGLE	AREA
Lower Eastern Shore - 14 (continued)	Bethel Beach, Va. (178)	0.00
	Goose Island, Va. (179)	<u>0.00</u>
		62,989,430 sq. m
		6,298.94 hectares
		15,564.71 acres
Reedville - 15	Heathsville, Va.-Md. (97)	0.00
	Burgess, Va.-Md. (98)	0.00
	Reedville, Va. (106)	3,041,640.81
	Irvington, Va. (111)	0.00
	Fleets Bay, Va. (112)	<u>5,089,223.06</u>
		8,130,864 sq. m
	813.09 hectares	
	2,009.14 acres	
Rappahannock River Complex-16	Tappahannock, Va. (95)	0.00
	Lottsburg, Va. (96)	0.00
	Dunnsville, Va. (103)	0.00
	Morattico, Va. (104)	0.00
	Lively, Va. (105)	0.00
	Urbanna, Va. (110)	0.00
	Irvington, Va. (111)	1,599,377.92
	Fleets Bay, Va. (112)	0.00
	Saluda, Va. (116)	0.00
	Wilton, Va. (117)	440,866.91
	Deltaville, Va. (118)	2,168,355.10
	Ware Neck, Va. (122)	0.00
	Mathews, Va. (123)	3,013,170.75
	Bethel Beach, Va. (178)	<u>0.00</u>
	7,221,771 sq. m	
	722.18 hectares	
	1,784.50 acres	
New Point Comfort Region-17	Mathews, Va. (123)	452,846.11
	New Point Comfort, Va. (132)	3,613,518.43

(continue on next page)

Table 6 (continued)

SECTION	QUADRANGLE	AREA
New Point Comfort Region-17 (continued)	East of New Point Comfort, Va. (177)	185,546.84
	Bethel Beach, Va. (178)	<u>57,536.15</u>
		4,309,448 sq. m
		430.94 hectares
		1,784.50 acres
Mobjack Bay Complex - 18	Ware Neck, Va. (122)	3,130,936.59
	Mathews, Va. (123)	492,775.91
	Clay Bank, Va. (130)	0.00
	Achilles, Va. (131)	7,269,886.02
	New Point Comfort, Va. (132)	7,533,040.90
	East of New Point Comfort, Va. (177)	<u>0.00</u>
		18,426,639 sq. m
		1,842.66 hectares
		4,553.23 acres
York River - 19	Toano, Va. (120)	0.00
	Gressitt, Va. (121)	0.00
	Norge, Va. (128)	0.00
	Williamsburg, Va. (129)	0.00
	Clay Bank, Va. (130)	0.00
	Achilles, Va. (131)	3,308,452.38
	New Point Comfort, Va. (132)	3,884,983.60
	Hog Island, Va. (138)	0.00
	Yorktown, Va. (139)	25,171.89
	Poquoson West, Va. (140)	1,187,626.25
	Poquoson East, Va. (141)	0.00
East of New Point Comfort, Va. (177)	<u>0.00</u>	
		8,406,234 sq. m
		840.62 hectares
		2,077.18 acres
Lower Western Shore - 20	New Point Comfort, Va. (132)	0.00
	Poquoson West, Va. (140)	4,997,149.37
	Poquoson East, Va. (141)	11,817,200.31

(continue on next page)

Table 6 (continued)

SECTION	QUADRANGLE	AREA
Lower Western Shore - 20 (continued)	Elliotts Creek, Va. (142)	0.00
	Newport News North, Va. (146)	0.00
	Hampton, Va. (147)	3,629,859.80
	Norfolk North, Va. (150)	0.00
	Little Creek, Va. (151)	0.00
	Cape Henry, Va. (152)	212,198.16
	Kempsville, Va. (156)	0.00
	Princess Anne, Va. (157)	0.00
		<hr/>
		20,656,408 sq. m
	2,065.64 hectares	
	5,104.20 acres	
James River - 21	Toano, Va. (120)	0.00
	Westover, Va. (125)	0.00
	Charles City, Va. (126)	0.00
	Brandon, Va. (127)	0.00
	Norge, Va. (128)	0.00
	Williamsburg, Va. (129)	0.00
	Savedge, Va. (135)	0.00
	Claremont, Va. (136)	0.00
	Surry, Va. (137)	0.00
	Hog Island, Va. (138)	0.00
	Yorktown, Va. (139)	0.00
	Poquoson West, Va. (140)	0.00
	Bacons Castle, Va. (144)	0.00
	Mulberry Island, Va. (145)	0.00
	Newport News North, Va. (146)	0.00
	Hampton, Va. (147)	40,070.48
	Benns Church, Va. (148)	0.00
	Newport News South, Va. (149)	0.00
	Norfolk North, Va. (150)	0.00
	Little Creek, Va. (151)	0.00
	Chuckatuck, Va. (153)	0.00
	Bowers Hill, Va. (154)	0.00
Norfolk South, Va. (155)	0.00	

(continue on next page)

Table 6 (continued)

SECTION	QUADRANGLE	AREA
James River - 21 (continued)	Kempsville, Va. (156)	0.00
	Princess Anne, Va. (157)	0.00
		<hr/>
		40,070 sq. m
		4.01 hectares
		9.90 acres
Chincoteague Bay	Selbyville, Md. (165)	0.00
	Assawoman Bay, Md. (166)	203,549.50
	Berlin, Md. (167)	153,620.34
	Ocean City, Md. (168)	330,609.61
	Public Landing, Md. (169)	0.00
	Tingles Island, Md. (170)	11,899,515.33
	Girdle Tree, Md.-Va. (171)	0.00
	Boxiron, Md.-Va. (172)	8,174,132.56
	Whittington Point, Md.-Va. (173)	4,516,484.63
	Chincoteague West, Va. (174)	139,216.00
	Chincoteague East, Va. (175)	<u>10,347,097.87</u>
		35,764,226 sq. m
		3,576.42 hectares
		8,837.35 acres

TABLE 7

Number of Square Meters of SAV in 1993 by Density Class for the 21 Sections of Chesapeake Bay and for Chincoteague Bay.

SECTION	DENSITY	AREA
Susquehanna Flats - 1	Density 1 =	15,177,640
	Density 2 =	752,937
	Density 3 =	459,625
	Density 4 =	<u>1,375,014</u>
	Total =	17,765,216
Upper Eastern Shore - 2	Density 1 =	1,174,501
	Density 2 =	320,802
	Density 3 =	310,376
	Density 4 =	<u>38,927</u>
	Total =	1,844,606
Upper Western Shore - 3	Density 1 =	35,905
	Density 2 =	498,326
	Density 3 =	255,228
	Density 4 =	<u>11,853</u>
	Total =	801,312
Chester River - 4	Density 1 =	889,735
	Density 2 =	1,360,136
	Density 3 =	2,225,875
	Density 4 =	<u>1,812,486</u>
	Total =	6,288,233
Central Western Shore - 5	Density 1 =	0
	Density 2 =	0
	Density 3 =	0
	Density 4 =	<u>0</u>
	Total =	0

(continue on next page)

TABLE 7 (continued)

SECTION	DENSITY	AREA
Eastern Bay - 6	Density 1 =	5,076,125
	Density 2 =	868,454
	Density 3 =	1,262,423
	Density 4 =	<u>165,461</u>
	Total =	7,372,464
Choptank River - 7	Density 1 =	3,472,813
	Density 2 =	11,201,823
	Density 3 =	3,638,963
	Density 4 =	<u>629,437</u>
	Total =	18,943,035
Patuxent River - 8	Density 1 =	0
	Density 2 =	87,762
	Density 3 =	9,946
	Density 4 =	<u>0</u>
	Total =	97,708
Middle Western Shore - 9	Density 1 =	0
	Density 2 =	0
	Density 3 =	0
	Density 4 =	<u>0</u>
	Total =	0
Lower Potomac River - 10	Density 1 =	92,763
	Density 2 =	1,145,772
	Density 3 =	949,431
	Density 4 =	<u>2,389,869</u>
	Total =	4,577,835

(continue on next page)

TABLE 7 (continued)

SECTION	DENSITY	AREA
Upper Potomac River - 11	Density 1 =	866,513
	Density 2 =	1,210,694
	Density 3 =	2,432,213
	Density 4 =	<u>19,117,110</u>
	Total =	23,626,529
Middle Eastern Shore - 12	Density 1 =	1,601,573
	Density 2 =	12,122,352
	Density 3 =	10,763,700
	Density 4 =	<u>5,233,779</u>
	Total =	29,721,403
Mid-Bay Island Complex - 13	Density 1 =	4,066,298
	Density 2 =	30,316,627
	Density 3 =	8,222,103
	Density 4 =	<u>12,062,759</u>
	Total =	54,667,787
Lower Eastern Shore - 14	Density 1 =	6,862,250
	Density 2 =	24,730,537
	Density 3 =	10,047,774
	Density 4 =	<u>21,348,869</u>
	Total =	62,989,430
Reedville - 15	Density 1 =	1,265,825
	Density 2 =	3,429,209
	Density 3 =	1,803,367
	Density 4 =	<u>1,632,463</u>
	Total =	8,130,864

(continue on next page)

TABLE 7 (continued)

SECTION	DENSITY	AREA
Rappahannock River Complex - 16	Density 1 =	667,773
	Density 2 =	4,365,164
	Density 3 =	1,612,282
	Density 4 =	<u>576,552</u>
	Total =	7,221,771
	New Point Comfort Region - 17	Density 1 =
Density 2 =		797,929
Density 3 =		2,472,834
Density 4 =		<u>703,293</u>
Total =		4,309,448
Mobjack Bay Complex - 18		Density 1 =
	Density 2 =	2,944,854
	Density 3 =	1,799,849
	Density 4 =	<u>12,916,931</u>
	Total =	18,426,639
	York River - 19	Density 1 =
Density 2 =		1,351,683
Density 3 =		0
Density 4 =		<u>6,916,671</u>
Total =		8,406,234
Lower Western Shore - 20		Density 1 =
	Density 2 =	2,875,213
	Density 3 =	5,685,864
	Density 4 =	<u>9,085,600</u>
	Total =	20,656,408

(continue on next page)

TABLE 7 (concluded)

SECTION	DENSITY	AREA
James River - 21	Density 1 =	0
	Density 2 =	0
	Density 3 =	40,070
	Density 4 =	<u>0</u>
	Total =	40,070
Chincoteague Bay	Density 1 =	936,199
	Density 2 =	9,415,315
	Density 3 =	6,046,968
	Density 4 =	<u>19,365,744</u>
	Total =	35,764,226
Chesapeake Bay Total	Density 1 =	45,497,719
	Density 2 =	100,374,417
	Density 3 =	53,954,745
	Density 4 =	<u>96,052,946</u>
	Total =	295,879,827

SAV

Elk River. The Upper Western Shore (Section 3) had 80 hectares of SAV compared to 186 hectares recorded in 1992. SAV was reported from the Gunpowder River area including Dundee Creek; the lower Spesutie Narrows; the Middle and Magothy rivers; and Romney and Delph creeks. SAV was noticeably reduced in Saltpeter Creek from 1992 and was absent from Seneca Creek in 1993. SAV was mapped in the Magothy River for the first time since it was last reported in 1978. *Myriophyllum spicatum*, *E. canadensis*, and *Z. palustris* were frequently cited. In the Chester River (Section 4) SAV abundance (629 hectares) was up 374 hectares from 1992. SAV was most abundant adjacent to Eastern Neck, Eastern Neck Island, and in the lower Chester River. *Ruppia maritima* was most commonly cited.

Middle Bay Zone

In 1993, 39.3% (5,467 hectares) of the SAV in the Middle Bay zone was found in the Mid-Bay Island Complex (Section 13) which includes the broad shoal area between Smith and Tangier Islands. This is a decrease of 527 hectares over 1992. In this zone, 21.4% (2972 hectares) of the SAV was present in the Middle Eastern Shore (Section 12), primarily in the Barren Island-Honga River area; the Big and Little Annemessex rivers; and the lower section of the Manokin River, with *R. maritima* reported most frequently. SAV was much less abundant in the Barren Island area. No SAV was mapped from the Central Western Shore (Section 5) and Middle Western Shore (Section 9). SAV in the Patuxent River (Section 8) increased from 0 hectares in 1992 to 10 hectares in 1993. Citizens' surveys reported *Z. palustris* and *R. maritima* at numerous locations in the South and Severn rivers. Eleven species were reported in the Patuxent River.

The Middle Bay zone also includes the entire Potomac River, where 2,820 hectares of SAV were present in 1993. SAV was concentrated in two distinct regions: 1) the Upper Potomac River (Section 11) with 2,363 hectares; and 2) the upper portion of the Lower Potomac River (Section 10) with 458 hectares, including Nanjemoy Creek and Port Tobacco River. Although the total abundance of SAV in the Upper Potomac section decreased from 1992 by 99 hectares, there was a notable increase in SAV in the Alexandria quadrangle. In particular, a large shoal area in the middle Potomac River, just above the large bed around the Woodrow Wilson Bridge, which had been previously unvegetated, supported sparse SAV for the first time since this survey was initiated. Ground truth data was reported by USGS and Citizens' surveys with 9 species reported: *M. spicatum*, *V. americana*, *H. verticillata*, *H. dubia*, *C. demersum*, *Najas minor*, *P. pectinatus*, *P. crispus*, and *N. guadalupensis*. SAV in the lower Potomac River also decreased (113 hectares) in 1993. Ground truth data was reported by Citizens' and VIMS surveys with 4 species cited: *Z. palustris*, *P. crispus*, *R. maritima*, and *M. spicatum*. SAV continued to increase in the Eastern Bay and Choptank River sections from 1992. SAV in the Eastern Bay (Section 6) increased 183 hectares from 1992 to a total of 737 hectares in 1993, while in the Choptank River (Section 7) it increased 809 hectares from 1992 to a total of 1,894 hectares in 1993. Most of the increase in the Eastern Bay occurred in Cox Creek, Crab Alley Bay, around Parson Island, and Piney Neck Point. In the Choptank River section, SAV

beds were most abundant in Harris and Broad creeks and in Trippe Bay. Two species were reported from Section 6, with *R. maritima* most commonly cited. Three species were reported from Section 7, with *R. maritima* most commonly cited.

Lower Bay Zone

Distribution and abundance of SAV in 1993, in the Lower Bay zone, were similar to 1992. In this zone, 48.4% (6,299 hectares) of the SAV was found in the Lower Eastern Shore (Section 14) around the Fox, Cedar, Webb, and Halfmoon islands; and the mouths of major creeks (i.e. Cherrystone Inlet; Hungars, Mattawoman, Nassawadox, Occohannock, Craddock, Pungoteague, Nandua, and Onancock creeks; and Beasley Bay). Along the western shore of the Chesapeake Bay, SAV was abundant in Mobjack Bay (Section 18) (14.2% of SAV in the Lower Bay zone), in the lower York River (Section 19) (6.5% of SAV in the Lower Bay zone), and in the Lower Western Shore (Section 20), specifically Back River and the Drum Island Flats area adjacent to Plum Tree Island (15.9% of SAV in the Lower Bay zone). Sparse SAV was documented for a segment of the south shore of the York River, downstream from Yorktown, for the first time in over twenty years (Orth and Gordon, 1975). There were 813 hectares of SAV mapped in the Reedville Region (Section 15) in 1993, a 4.5% increase over 1992. There were 431 hectares of SAV identified in 1993 in the New Point Comfort Region (Section 17), compared to 396 hectares in 1992. SAV abundance was up 23.1% from 1992 in both the Piankatank and Rappahannock rivers (Section 16). The James River (Section 21) had 4 hectares of SAV in 1993. *Zostera marina* and *R. maritima* were the abundant species in this zone.

Chincoteague Bay

SAV in the Chincoteague Bay section increased in distribution with 3,576 hectares mapped in 1993 compared to 3,324 hectares in 1992. Most of the SAV in Chincoteague and Sinepuxent bays, which consisted mainly of *Z. marina* and *R. maritima*, was located along the eastern side of the bay behind Assateague Island. Some small beds, consisting mainly of *R. maritima*, were located along the eastern side of Isle of Wight and Assawoman bays.

DISCUSSION OF SECTIONS ARRANGED WITHIN ZONES

Upper Bay Zone

1. Susquehanna Flats

There were 1,777 hectares of SAV in the Susquehanna Flats section in 1993 (Tables 4-7; Figure 8; Appendix C, Maps 2, 3, 8 and 9), compared to 1,792 hectares mapped in 1992. In this section 7.7% of the total coverage of SAV in this section was dense (class 4), 2.6% was moderate (class 3), 4.2% was sparse (class 2), and 85.4% was very sparse (class 1) (Table 7; Figure 3). SAV beds were located principally in two main areas: 1) sparse to dense fringing beds in the Susquehanna River consisting primarily of *M. spicatum*, with *H. dubia*, *V. americana*, *H. verticillata*, and *C. demersum* from Robert Island to the river mouth at Havre de Grace on the west side, to Stump Point at the mouth of Mill Creek on the east side, and in Mill Creek, Furnace Bay, Baker Cove, and at High Point; and 2) a large area of very sparse SAV located in the broad shoal area at the river mouth. This broad shoal area continues to consist of small patches of *M. spicatum*. Additionally, the Citizens' survey reported *H. verticillata*. In addition, SAV beds were again mapped in Spesutie Narrows where most SAV is found as small, fringing beds of *M. spicatum* and *H. verticillata*.

A total of six species (*M. spicatum*, *H. dubia*, *V. americana*, *H. verticillata*, *C. demersum*, *N. guadalupensis*) have been reported on Maps 2, 3, and 9 (Appendix C), either by Stan Kollar of Harford Community College or the Citizens' Survey.

2. Upper Eastern Shore

There were 184 hectares of SAV mapped for the Upper Eastern Shore section in 1993 (Tables 4-7; Figure 9; Appendix C, Maps 4, 9, 10, 15, 16, and 17), compared to 283 hectares mapped for 1992. In this section 2.1% of the total coverage of SAV was dense (class 4), 16.8% was moderate (class 3), 17.4% was sparse (class 2), and 63.7% was very sparse (class 1) (Table 7; Figure 3). Principal locations of beds were in the Elk River and the lower Sassafras River. Very little SAV was mapped in the Bohemia River or along the mainstem of the bay from Still Pond to Swan Point. Much of the decrease was recorded in the Elk River. Ground survey data from Stan Kollar and the Citizens' survey reported 5 species in this section (Appendix C, Maps 4, 9, 10, 16, and 17), with *M. spicatum* and *V. americana* found most frequently. *Heteranthera dubia*, *P. crispus*, *P. pectinatus*, and an unidentified species were also reported in the Elk River.

3. Upper Western Shore

There were 80 hectares of SAV mapped from the aerial photographs in 1993 for the Upper Western




Figure 8. Distribution of SAV in the Susquehanna Flats (Section 1) in 1993.




Figure 9. Distribution of SAV in the Upper Eastern Shore (Section 2) in 1993.

Shore section (Tables 4-7; Figure 10; Appendix C, Maps 8, 9, 13, 14, and 24), compared to 186 hectares in 1992. Of the total coverage of SAV in this section 1.5% was dense (class 4), 31.9% was moderate (class 3), and 62.2% was sparse (class 2), and 4.5% was very sparse (Table 7; Figure 3). SAV beds were located in Romney, Little Romney, and Delph creeks, the lower Spesutie Narrows, the Gunpowder River area, including Dundee Creek, and the Magothy and Middle rivers. SAV was noticeably reduced in Saltpeter Creek and was absent in Seneca Creek. SAV was mapped for the first time in the Magothy River since 1978. As in 1992, no SAV was reported in the Back, Patapsco, and Bush rivers.

Ground survey data from the Citizens' survey and the U.S. Fish and Wildlife Service reported 11 species in this section (Appendix C, Maps 9, 13, 14, 23, and 24). *Myriophyllum spicatum*, *E. canadensis*, and *Z. palustris* were found most frequently. *Vallisneria americana*, *P. crispus*, *P. pectinatus*, *P. perfoliatus*, *R. maritima*, *Najas flexilis*, *Najas* spp., and *C. demersum* were reported less frequently.

4. Chester River

There were 629 hectares of SAV in the Chester River section in 1993 (Tables 4-7; Figure 11; Appendix C, Maps 20, 21, 26, and 33), compared to 255 hectares in 1992. In this section, 28.8% of the total coverage of SAV was dense (class 4), 35.4% was moderate (class 3), 21.6% was sparse (class 2), and 14.1% was very sparse (class 1) (Table 7; Figure 3). This is a notable increase since 1991, when only 57 hectares were reported. Most of the SAV, and where the greatest increase occurred, was located adjacent to Eastern Neck and Eastern Neck Island, especially near Eastern Neck Narrows; in Grays Inn, Langford, and Queenstown creeks, tributaries entering the Chester River. Rock Hall Harbor; the Haven; and Swan, Langford, and Huntingfield creeks, located above Eastern Neck on the Chesapeake Bay, supported the remaining SAV beds in this section. Eight species of SAV were reported from this section by Citizens', U.S. Fish and Wildlife Service, and VIMS surveys in 1993: *Potamogeton perfoliatus*, *Z. palustris*, *R. maritima*, *M. spicatum*, *C. demersum*, *E. canadensis*, *N. spp.*, and *N. flexilis* (Appendix C, Maps 26 and 33).

Middle Bay Zone

5. Central Western Shore

There was no SAV observed from the aerial photography in the Central Western Shore section in 1993 (Tables 4-7; Figure 12). This was the same as 1992. Citizens' surveys found *Z. palustris* and *R. maritima* at numerous locations in the Severn and South rivers (Appendix C, Maps 30 and 31) as well as *Potamogeton pusillus* at Lake Ogleton.




Figure 10. Distribution of SAV in the Upper Western Shore (Section 3) in 1993.




Figure 11. Distribution of SAV in the Chester River (Section 4) in 1993.




Figure 12. Distribution of SAV in the Central Western Shore (Section 5) in 1993.

6. Eastern Bay

There were 737 hectares of SAV identified from the Eastern Bay section in 1993 (Tables 4-7; Figure 13; Appendix C, Maps 32, 33, 36, and 37), compared to 554 hectares reported in 1992, a 33.1% increase. This is the second year in a row that SAV has increased in this area. Only 68 hectares were reported in 1991. In this section, 2.2% of the total coverage of SAV in this section was dense (class 4), 17.1% was moderate (class 3), 11.8% was sparse (class 2), and 68.9% was very sparse (class 1) (Table 7; Figure 3). Most of the SAV was found in the lower Miles River; the lower Cox Creek; Wye River; the eastern shore of lower Kent Island; Parson Island; Harbor Cove on Eastern Bay; Piney Neck; Crab Alley Bay; and between Harbor Cove and Tilghman Point on Eastern Bay. Only two species of SAV were reported from this section by Citizens' and VIMS surveys: *Z. palustris* and *R. maritima*, (Appendix C, Maps 32, 33, and 36) with *R. maritima* most commonly cited.

7. Choptank River

There were 1894 hectares of SAV observed in the Choptank River section in 1993 (Tables 4-7; Figures 14 and 15; Appendix C, Maps 36, 37, 43, 44, 51, 52, 53, and 62), compared to 1085 hectares in 1992. This was an increase of 74.5%. In this section, 3.3% of the total coverage of SAV was dense (class 4), 19.2% was moderate (class 3), 59.1% was sparse (class 2), and 18.3% was very sparse (class 1) (Table 7; Figure 3). In the Choptank River, SAV was found in moderate to sparse beds in Blackwalnut Cove at the southern tip of Tilghman Island: Harris, Broad, San Domingo, Edge, Irish, and Chapel creeks; the Tred Avon River; Dickinson Bay; and Castle Haven Point. Moderate to sparse beds were also found in Trippe Bay; the lower Choptank River, including James Island, Hills Point; Oyster, Cators, and Hooper coves; and Slaughter, Brooks, Hudson, and Back creeks.

Three species of SAV were reported from this section by Citizens' and VIMS surveys: *Z. palustris*, *R. maritima*, *P. perfoliatus*, and an unidentified species (Appendix C, Maps 36, 37, 43, 44, 51, 52, and 53) with *R. maritima* most commonly cited.

8. Patuxent River

There were 10 hectares of SAV observed in 1993 in the Patuxent River section (Tables 4-7; Figures 16 and 17, Appendix C, Maps 41 and 71), compared to zero hectares in 1992. One bed was located in the upper Patuxent River where many of the Citizens' reports were located, while a second bed was noted at the mouth of the river between Solomons Island and Drum Point. In this section, 10.2% of the total coverage was moderate (class 3), while 89.8% was sparse (class 2). A total of eleven species were reported in this section. The Citizens' survey reported ten species occurring primarily in the marsh creeks in the upper portions of the Patuxent River (Appendix C, Maps 41, 49, and 159): *E. canadensis*, *C. demersum*, *V. americana*, *Z. palustris*, *N. guadalupensis*, *N. minor*, *P. crispus*, *P. pusillus*, *H. verticillata*, and *Potamogeton epihydrus*. *Zannichellia palustris* and *R. maritima* were




Figure 13. Distribution of SAV in the Eastern Bay (Section 6) in 1993.




Figure 14. Distribution of SAV in the Choptank River (Section 7) in 1993.




Figure 15. Detail of Figure 14 showing distribution of SAV in the Choptank River in 1993.




Figure 16. Distribution of SAV in the Patuxent River (Section 8) in 1993.




Figure 17. Detail of Figure 16 showing distribution of SAV in the Patuxent River in 1993.

reported from the lower portion of the Patuxent River in Saint Leonard, Island, Hominy, and Cuckold creeks; Peterson Point; and Green Holly Pond (Appendix C, Maps 60, 61, 70, and 71).

9. Middle Western Shore

There were no SAV beds identified in the Middle Western Shore section in 1993 (Tables 4-7; Figure 18) the same as in 1992.

10. Lower Potomac River

There were 458 hectares of SAV identified in the Lower Potomac River section as indicated on the 1993 aerial photography (Tables 4-7; Figures 19, 20, and 21; Appendix C, Maps 56, 57, 58, 65, 66, 67, 80, 89, and 162), compared to 571 hectares reported in 1992 with most of the changes occurring in Nanjemoy Creek. In this section, 52.2% of the total coverage of SAV was dense (class 4), 20.7% was moderate (class 3), 25.0% was sparse (class 2), and 2.0% was very sparse (class 1) (Table 7; Figure 3). Most of the SAV occurred along Nanjemoy Creek and Port Tobacco River, as well as along the shoreline adjacent to these two creeks, and fringing the eastern side of Mathias Point Neck to an area just below the Route 301 Bridge (Harry Nice Memorial Bridge). Several small beds were observed in Machodoc, Rosier, and Cuckold creeks; the St. Marys River; in Calvert Bay at the mouth of Smith Creek; Lloyd Point; above Lower Cedar Point; and in the upper Wicomico River.

Ground survey data was available from Citizens' and VIMS surveys for Maps 68, 76, 78, and 80 (Appendix C). *Zannichellia palustris* and *P. crispus* were reported from the Lower Machodoc Creek (Appendix C, Map 78). *Zannichellia palustris* was found in Neale Sound off the Wicomico River (Appendix C, Map 68). *Ruppia maritima* was reported from the St. Marys River (Appendix C, Map 80). *Myriophyllum spicatum* was reported in Popes Creek (Appendix C, Map 76).

11. Upper Potomac River

There were 2,363 hectares of SAV mapped in the Upper Potomac River section (Tables 4-7; Figures 22, 23, and 24; Appendix C, Maps 28, 29, 34, 39, 40, 47, 48, 55, 64, 65, and 161) in 1993, compared to 2,462 hectares reported in 1992, a decrease of 4%. A total of 80.9% of the SAV beds were densely vegetated (class 4), 10.3% was moderate (class 3), 5.1% was sparse (class 2), and 3.7% was very sparse (class 1) (Table 7; Figure 3). Although there was an overall reduction in SAV abundance, there was a notable change in SAV distribution in the Alexandria quadrangle (Appendix C, Map 34). An expansive shoal area in the middle of the river, just above the large dense bed surrounding the Woodrow Wilson Bridge, supported sparse SAV (bed FB1) for the first time in 1993. The western side of the mainstem Potomac River, from the Woodrow Wilson Bridge to Occoquan Bay remains very sparsely vegetated.

Ground survey data was available only from USGS and Citizens' surveys for Maps 28, 34, 39, 40,




Figure 18. Distribution of SAV in the Middle Western Shore (Section 9) in 1993.




Figure 19. Distribution of SAV in the Lower Potomac River (Section 10) in 1993.




Figure 20. Detail of Figure 19 showing distribution of SAV in the Lower Potomac River in 1993.




Figure 21. Detail of Figure 19 showing distribution of SAV in the Lower Potomac River in 1993.




Figure 22. Distribution of SAV in the Upper Potomac River (Section 11) in 1993.




Figure 23. Detail of Figure 22 showing distribution of SAV in the Upper Potomac River in 1993.




Figure 24. Detail of Figure 22 showing distribution of SAV in the Upper Potomac River in 1993.

47, 48, 55, and 161 (Appendix C). Nine species were reported: *M. spicatum*, *V. americana*, *H. verticillata*, *N. guadalupensis*, *C. demersum*, *H. dubia*, *P. pectinatus*, *P. crispus*, and *N. minor*.

12. Middle Eastern Shore

There were 2,972 hectares of SAV identified in the Middle Eastern Shore section (Tables 4-7; Figure 25; Appendix C, Maps 63, 72, 73, 74, 82, 83, 84, 85, 92, 93, 100, and 101) in 1993, compared to 3,047 hectares reported in 1992. In this section, 17.6% of the SAV was dense (class 4), 36.2% moderate (class 3), 40.8% sparse (class 2), and 5.4% very sparse (class 1) (Table 7; Figure 3). SAV beds were very abundant in: 1) the Honga River, 2) between Barren Island and Meekins Neck-Upper Hooper Island, and 3) the lower Manokin and the Big and Little Annemessex rivers. SAV abundance declined in the Barren Island area from 1992 (Barren Island quadrangle - 434 hectares in 1992 to 206 hectares in 1993), while increasing in the Honga River area (Honga quadrangle - 1,327 hectares in 1992 to 1340 hectares in 1993; Wingate quadrangle - 481 hectares in 1992 to 541 hectares in 1993; Golden Hill quadrangle - 29 hectares in 1992 to 65 hectares in 1993). No SAV beds were observed in Fishing or Monie bays, and in the Nanticoke and Wicomico rivers. Ground survey data were available for this section in 1993 from Citizens' and VIMS surveys (Appendix C, Maps 72, 73, 74, 84, 100, and 101). *Ruppia maritima* and *Z. marina* was reported most frequently.

13. Mid-Bay Island Complex

There were 5,467 hectares of SAV mapped in the Mid-Bay Island Complex in 1993 (Tables 4-7; Figure 26; Appendix C, Maps 83, 91, 92, 99, 100, 107, and 179), compared to 5,994 hectares reported in 1992, an 8.8% decrease, and with declines in abundance noted in all quadrangles. This section contains 18.5% of the SAV in the entire Chesapeake Bay, slightly less than the 21.0% in 1992. However, the density of SAV has decreased since 1992. In 1993, 37.1% of the SAV within this section was in dense and moderate beds (classes 3 and 4), compared to 70.8% in 1992. In the remaining classes, 55.4% was sparse (class 2), compared to 27.1% in 1992, and 7.4% was very sparse (class 1) compared to 2.1% in 1992 (Table 7; Figure 3).

SAV is present mainly in dense to moderate beds along the broad, expansive shoal area between Tangier Island and Smith Island; the eastern side of Tangier Island; the western side of Goose Island; Mailboat Harbor; and on the shoals and in the coves adjacent to Bloodsworth, South Marsh, Holland, Adam, and Spring islands. The Citizens' ground survey reported *R. maritima*, and *Z. marina* (Appendix C, Map 99) for this section in 1993.

Lower Bay Zone

14. Lower Eastern Shore

There were 6,299 hectares of SAV observed in the Lower Eastern Shore section in 1993 (Figure 27;




Figure 25. Distribution of SAV in the Middle Eastern Shore (Section 12) in 1993.




Figure 26. Distribution of SAV in the Mid-Bay Island Complex (Section 13) in 1993.




Figure 27. Distribution of SAV in the Lower Eastern Shore (Section 14) in 1993.

Tables 4-7; Appendix C, Maps 100, 101, 102, 107, 108, 109, 113, 114, 119, 124, 133, 134, and 142), compared to 5,920 hectares reported in 1992. In this section 33.9% of the total SAV was dense (class 4), 16.0% was moderate (class 3), 39.3% was sparse (class 2), and 10.9% was very sparse (class 1) (Table 7; Figure 3). Large, dense beds continue to persist at the mouth of Cherrystone Inlet near Cape Charles, and at the mouths of Hungars, Nassawadox, Mattawoman, Occohannock, Craddock, Pungoteague, Onancock, Nandua, and Chesconessex creeks. Large, dense beds also occur at the Big Marsh area near Chesconessex Creek, at Webb and Halfmoon islands off the mouth of Deep Creek, and on the large shoal area on the eastern side of the Fox and Cedar islands. SAV beds also were mapped along Ware Point Marsh and around Beasley Bay in the Pocomoke Sound. There was no SAV from Elliots Creek, just below Cape Charles, to Fishermans Island at the mouth of Chesapeake Bay. Ground survey data were limited in this section. Citizens' and VIMS surveys reported *R. maritima* and *Z. marina* from this section (Appendix C, Maps 100, 101, 114, 124, and 133).

15. Reedville Region

There were 813 hectares of SAV identified in the Reedville Region in 1993 (Tables 4-7; Figure 28; Appendix C, Maps 106 and 112), compared to 778 hectares reported in 1992. In this section, 20.1% of the total coverage of SAV was dense (class 4), 22.2% was moderate (class 3), 42.2% was sparse (class 2), and 15.6% was very sparse (class 1) (Table 7; Figure 3). Most beds were found in Little, Fleets, and Ingram bays; Dwymer, Indian, Dividing, Ball, and Cloverdale creeks; Dameron Marsh; and adjacent to Fleeton Point. *Zostera marina* and *R. maritima* were the two species identified by Citizens' surveys (Appendix C, Maps 106 and 112).

16. Rappahannock River Complex

There were 722 hectares of SAV observed in the Rappahannock River Complex in 1993 (Tables 4-7; Figure 29; Appendix C, Maps 111, 117, 118, and 123), compared to 587 hectares reported in 1992. In this section 8.0% of the total coverage of SAV was dense (class 4), 22.3% was moderate (class 3), 60.4% was sparse (class 2), and 9.2% was very sparse (class 1) (Table 7; Figure 3). SAV beds were present in the Corrotoman River; along the north shore of the Rappahannock River, from the Corrotoman River to Windmill Point, Milford Haven, the lower Piankatank River; and from Ginney and Horse points to Gwynns Island. SAV beds have declined in the area between Carters Creek and the mouth of the Corrotoman River. The large SAV bed adjacent to Windmill Point that has been slowly expanding naturally since 1989 (contiguous beds BB2, CB1, DB4; Appendices C and D, Map 118), now covers an area of 44 hectares, up from 28 hectares in 1992, with both *Z. marina* and *R. maritima* present. SAV beds are abundant in Milford Haven, but are principally located along the north shore with both *Z. marina* and *R. maritima* present. *Ruppia maritima* and *Z. marina* were reported from ground surveys by VIMS staff and Citizens' surveys of SAV in Maps 111, 118, and 123 (Appendix C).




Figure 28. Distribution of SAV in the Reedville Region (Section 15) in 1993.




Figure 29. Distribution of SAV in the Rappahannock River Complex (Section 16) in 1993.

17. New Point Comfort Region

There were 431 hectares of SAV identified in the New Point Comfort Region in 1993 (Tables 4-7; Figure 30; Appendix C, Map 123, 132, 177, and 178), compared to 396 hectares reported in 1992. In this section 16.3% of the total coverage of SAV was dense (class 4), 57.4% was moderate (class 3), 18.5% was sparse (class 2), and 7.8% was very sparse (class 1) (Table 7; Figure 3). SAV beds were present from New Point Comfort just north of Horn Harbor at Potato Neck. SAV beds were also present at Winter Harbor. There was no ground truth data reported for this section.

18. Mobjack Bay Complex

The Mobjack Bay Complex contained 1,843 hectares of SAV in 1993 (Tables 4-7; Figure 31; Appendix C, Maps 122, 123, 131, and 132), compared to 1,818 hectares reported in 1992. SAV beds were abundant along the entire shoreline of Mobjack Bay, as well as in the lower reaches of the four tributaries: Severn, Ware, North, and East rivers. The Mobjack Bay area continued to harbor some of the more extensive SAV beds on the western shore of the lower Chesapeake Bay. In this section 70.1% of the total coverage of SAV was dense (class 4), 9.8% was moderate (class 3), 16.0% was sparse (class 2), and 4.2% was very sparse (class 1) (Table 7; Figure 3). *Zostera marina* and *R. maritima* were reported by Citizens' and VIMS surveys from Maps 122, 131, and 132 (Appendix C).

19. York River

There were 841 hectares of SAV observed in the York River section in 1993 (Tables 4-7; Figure 32; Appendix C, Maps 131, 132, 139, and 140), compared to 830 hectares reported in 1992. In this section 82.3% of the total coverage is classified as dense (class 4), while 0.0% was moderately dense (class 3), 16.1% was sparse (class 2), and 1.6% was very sparse (class 1) (Table 7; Figure 3). Ground survey information was available for Maps 131, 132, 139, and 140 (Appendix C) from VIMS and Citizens' surveys. Dense SAV beds, consisting of both *Z. marina* and *R. maritima*, were located principally along the north shore from Gloucester Point to the mouth of the river and on the south shore adjacent to Goodwin Island. SAV beds were absent upstream of Gloucester Point along the north shore except for one small bed (Appendix C, Map 139, bed FA1) of *Z. marina* near Gloucester Point, a result of a VIMS transplanting project using seeds in 1989, 1990, and 1991. SAV was documented for the first time since 1971 along the south shore from Yorktown to the Coast Guard pier (Orth and Gordon, 1975). These two very sparse beds (Appendix C, Map 140, beds AB1 and BB1) consisted of small patches of *Z. marina*. These patches were most likely the result of seed recruitment in the spring of 1991 from SAV beds downriver on the south shore (Goodwin Island), or along the north shore directly across the river. This conclusion is based upon an understanding of the reproductive biology of *Z. marina*, which produces reproductive shoots in the second year of growth. These patches were too small in 1992 to have been documented by aerial photography. The presence of reproductive shoots in 1993 indicated that these patches had to be growing here in 1992,




Figure 30. Distribution of SAV in the New Point Comfort Region (Section 17) in 1993.




Figure 31. Distribution of SAV in the Mobjack Bay Complex (Section 18) in 1993.




Figure 32. Distribution of SAV in the York River (Section 19) in 1993.

SAV

but were from seeds produced and germinated in 1991.

Zostera marina was transplanted in the form of whole plants to the following sites: along the north shore at Claybank, and Mumford and Catlett Islands; and along the south shore at Yorktown. The transplanting effort was conducted by VIMS staff during the fall of 1992. Transplants remained present through the summer of 1993. By the late summer of 1993, transplants survived at the Mumford Island and Yorktown sites, but did not survive at the Catlett Island or Clay Bank sites.

20. Lower Western Shore

There were 2,066 hectares of SAV mapped in the Lower Western Shore section in 1992 (Tables 4-7; Figure 33; Appendix C, Maps 140, 141, 147, and 152), compared to 2,029 hectares reported in 1992.

In this section 44.0% of the total coverage was mapped as dense (class 4), 27.5% as moderate (class 3), 13.9% as sparse (class 2), and 14.6% as very sparse (class 1) (Table 7; Figure 3). SAV was mapped in Broad Bay; Back River, including the lower Northwest Branch; the lower Poquoson River; the mouth of the Poquoson River off Pasture and Hunts Neck; the lower Chisman Creek; Drum Island Flats; Poquoson Flats; adjacent to Crab Neck just south of Goodwin Island; and on the south side of Goodwin Island. No SAV was present in the Southwest Branch of Back River; in Back Creek; or from Northend Point to Old Point Comfort. Ground surveys by Citizens' and VIMS (Appendix C, Maps 140 and 152) reported both *Z. marina* and *R. maritima*.

21. James River

There were 4 hectares of SAV in the mainstem of the James River in 1993 (Tables 4-7; Figures 34 and 35; Appendix C, Map 147), compared to 3.5 hectares in 1992. This single, very dense bed, (class 4) (Table 7; Figure 3) located at the mouth of Hampton Creek adjacent to the Veteran's Hospital, consists of *Z. marina*, the species reported in previous ground surveys, and continues to remain the only SAV detected bed in the James River.

Chincoteague Bay

There were 3,576 hectares of SAV identified from the Eastern Shore of Virginia and Maryland in 1993. Chincoteague and Sinepuxent bays had 3556 hectares, and a small amount (20.35 hectares) was present in Isle of Wight and Assawoman bays (Tables 4-7; Figure 36; Appendix C, Maps 166, 167, 168, 170, 172, 173, 174, and 175), compared to 3,324 hectares reported in 1992. In this section 54.1% of the total coverage was mapped as dense (class 4), 16.9% as moderate (class 3), 26.3% as sparse (class 2), and 2.6% as very sparse (class 1) (Table 7; Figure 3). The Citizens' survey found both *Z. marina* and *R. maritima* throughout Chincoteague and Sinepuxent bays, as well as Assawoman bays (Appendix C, Maps 166, 167, 168, 170, 172, 173, and 175). All of the SAV in Chincoteague Bay continues to be present on the eastern side of the bay adjacent to Assateague Island. The vegetation remains concentrated in four relatively distinct areas identical to that reported




Figure 33. Distribution of SAV in the Lower Western Shore (Section 20) in 1993.




Figure 34. Distribution of SAV in the James River (Section 21) in 1993.




Figure 35. Detail of Figure 34 showing the distribution of SAV in the James River in 1993.




Figure 36. Distribution of SAV in Chincoteague Bay in 1993.

in the earlier surveys from 1986 through 1992. They were located west of the northern end of Chincoteague Island; west of the Tingles Island area; and in Green Run and West bays. SAV in Isle of Wight and Assawoman bays also remains present on the eastern side adjacent to Ocean City.

SAV

LITERATURE CITED

- Anderson, R. R. and R. T. Macomber. 1980. Distribution of submersed vascular plants, Chesapeake Bay, Maryland. Final Report to U.S. EPA, Chesapeake Bay Program. Grant No. R805970. 126 pp.
- Chesapeake Executive Council. 1989. Submerged Aquatic Vegetation Policy for the Chesapeake Bay and Tidal Tributaries. Annapolis, MD. July.
- Chesapeake Executive Council. 1990. Implementation Plan for the Submerged Aquatic Vegetation Policy. Annapolis, MD. July.
- Environmental Systems Research Institute, Inc. 1989. Volume I & II, User Guide. ESRI, Redlands, CA.
- Godfrey, R. K. and J. W. Wooten. 1981. Aquatic and Wetland Plants of Southeastern United States: Dicotyledons. The University of Georgia Press, Athens. 933 pp.
- Godfrey, R. K. and J. W. Wooten. 1979. Aquatic and Wetland Plants of Southeastern United States: Monocotyledons. The University of Georgia Press, Athens. 712 pp.
- Harvill, A. M. Jr., C. E. Stevens and D. M. E. Ware. 1977. Atlas of the Virginia Flora: Part I, Pteridophytes through Monocotyledons. Virginia Botanical Associates, Farmville. 59 pp.
- Harvill, A. M. Jr., T. R. Bradley and C. E. Stevens. 1981. Atlas of the Virginia Flora: Part II, Dicotyledons. Virginia Botanical Associates, Farmville. 148 pp.
- Humm, Harold J. 1979. The Marine Algae of Virginia. Special Papers in Marine Science, Number 3, Virginia Institute of Marine Science. The University Press of Virginia, Charlottesville. 263 pp.
- Kartesz, J. T. and R. Kartesz. 1980. A Synonymized Checklist of the Vascular Flora of the United States, Canada and Greenland: Volume II, The Biota of North America. The University of North Carolina Press, Chapel Hill. 498 pp.
- Orth, R. J. and Hayden Gordon. 1975. Remote sensing of submerged aquatic vegetation in the lower Chesapeake Bay. Final Report to National Aeronautical and Space Administration, Langley Research Center. Contract NAS1-10720. 62 pp.
- Orth, R. J. and K. A. Moore. 1981. Submerged aquatic vegetation in the Chesapeake Bay: past, present and future. pp. 271-283. In: Proc. 46th North American Wildlife and Natural Resources Conf., Wildlife Manage. Inst., Wash., D.C.
- Orth, R. J. and K. A. Moore. 1982. The biology and propagation of *Zostera marina*, eelgrass, in the

- Chesapeake Bay, Virginia. Final Report to U.S. EPA, Chesapeake Bay Program. Grant No. R805953. 187 pp.
- Orth, R. J. and K. A. Moore. 1983. Chesapeake Bay: an unprecedented decline in submerged aquatic vegetation. *Sci.* 222:51-53.
- Orth, R. J. and K. A. Moore. 1984. Distribution and abundance of submerged aquatic vegetation in Chesapeake Bay: an historical perspective. *Est.* 7:531-540.
- Orth, R. J. and K. A. Moore. 1988. Submerged aquatic vegetation in the Chesapeake Bay: A barometer of Bay health. pp. 619-629. In: M. Lynch (Ed.), *Understanding the Estuary: Advances in Chesapeake Bay Res.* Chesapeake Res. Consort. Pub. No. 129. CBP/TRS/24/88.
- Orth, R. J., K. A. Moore and H. H. Gordon. 1979. Distribution and abundance of submerged aquatic vegetation in the lower Chesapeake Bay, Virginia. Final Report to U.S. EPA, Chesapeake Bay Program. EPA-600/8-79-029/SAV1.
- Orth, R. J. and J. F. Nowak. 1990. Distribution of submerged aquatic vegetation in the Chesapeake Bay and tributaries and Chincoteague Bay - 1989. Final Report to U.S. EPA. 247pp.
- Orth, R. J., J. F. Nowak, A. A. Frisch, K. P. Kiley, and J. R. Whiting. 1991. Distribution of submerged aquatic vegetation in the Chesapeake Bay and tributaries and Chincoteague Bay - 1990. Final Report to U.S. EPA. 261pp.
- Orth, R. J., J. F. Nowak, G. F. Anderson, K. P. Kiley, and J. R. Whiting. 1992. Distribution of submerged aquatic vegetation in the Chesapeake Bay and tributaries and Chincoteague Bay - 1991. Final Report to U.S. EPA. 268pp.
- Orth, R. J., J. F. Nowak, G. F. Anderson, and J. R. Whiting. 1993. Distribution of submerged aquatic vegetation in the Chesapeake Bay and tributaries and Chincoteague Bay - 1992. Final Report to U.S. EPA. 268pp.
- Orth, R. J., J. Simons, R. Allaire, V. Carter, L. Hindman, K. Moore and N. Rybicki. 1985. Distribution of submerged aquatic vegetation in the Chesapeake Bay and tributaries - 1984. Final Report to U.S. EPA, Coop. Agreement X-003301-01. 155 pp.
- Orth, R. J., J. Simons, J. Capelli, V. Carter, L. Hindman, S. Hodges, K. Moore and N. Rybicki. 1986. Distribution of submerged vegetation in the Chesapeake Bay and tributaries - 1985. Final Report to U.S. EPA. 296 pp.
- Orth, R. J., J. Simons, J. Capelli, V. Carter, A. Frisch, L. Hindman, S. Hodges, K. Moore and N.

SAV

- Rybicki. 1987. Distribution of submerged aquatic vegetation in the Chesapeake Bay and tributaries and Chincoteague Bay - 1986. Final Report to U.S. EPA. 180 pp.
- Orth, R. J., A. A. Frisch, and J. F. Nowak. 1988. Quality assurance project plan for the 1987 submerged aquatic vegetation distribution and abundance survey of the Chesapeake and Chincoteague Bays. Plan submitted to U.S. EPA, Chesapeake Bay Liason Office, Annapolis, MD. 47 pp.
- Orth, R. J., A. A. Frisch, J. F. Nowak, and K. A. Moore. 1989. Distribution of submerged aquatic vegetation in the Chesapeake Bay and tributaries and Chincoteague Bay - 1987. Final Report to U.S. EPA. 247pp.
- Radford, A. E., H. E. Ahles and C. R. Bell. 1968. *Manual of the Vascular Flora of the Carolinas*. The University of North Carolina Press, Chapel Hill. 1183 pp.
- Stevenson, J. C. and N. Confer. 1978. Summary of Available Information on Chesapeake Bay Submerged Vegetation. U.S. Dept. of Interior, Fish and Wildlife Serv. FWS/OBS-78/66. 335 p.
- Wood, R. D. and K. Imahori. 1965. A Revision of the Characeae: Volume I, Monograph of the Characeae. Verlag Von J. Cramer, Weinheim. 904 pp.
- Wood, R. D. and K. Imarhori. 1964. A Revision of the Characeae: Volume II, Iconograph of the Characeae. Verlag Von J. Cramer, Weinheim. 395 icones with Index.

APPENDICES

APPENDIX A




APPENDIX A

Species of Submerged Aquatic Plants Found in Chesapeake Bay and Tributaries Exclusive of Marine Algae (Classification and Nomenclature Derived from: Godfrey and Wooten, 1979, 1981; Harvill et al., 1977, 1981; Kartesz and Kartesz, 1980; Radford et al., 1968; Wood and Imahori, 1965, 1964)

Family	Species	Common name
Characeae (muskgrass)	<i>Chara braunii</i> Gm.	Muskgrass
	<i>Chara zeylanica</i> Klein. ex Willd., em.	Muskgrass
	<i>Nitella flexilis</i> (L.) Ag., em.	Stonewort
Potamogetonaceae (pondweed)	<i>Potamogeton perfoliatus</i> L. var. <i>bupleuroides</i> (Fernald) Farwell	Redhead grass
	<i>Potamogeton epihydrus</i>	Leafy pondweed
	<i>Potamogeton pectinatus</i> L.	Sago pondweed
	<i>Potamogeton crispus</i> L.	Curly pondweed
	<i>Potamogeton pusillus</i> L.	Slender pondweed
Ruppiales	<i>Ruppia maritima</i> L.	Widgeon grass
Zannichelliaceae	<i>Zannichellia palustris</i> L.	Horned pondweed
Najadaceae	<i>Najas guadalupensis</i> (Sprengel) Magnus	Southern naiad
	<i>Najas gracillima</i> (A. Braun) Magnus	Slender naiad
	<i>Najas minor</i> Allioni	no common name
	<i>Najas flexilis</i> (Willd.) Rostk. & Schmidt	Northern naiad
Hydrocharitaceae (frogbit)	<i>Vallisneria americana</i> Michaux	Wild celery, tapegrass
	<i>Elodea canadensis</i> (Michaux)	Common elodea
	<i>Egeria densa</i> Planchon	Water-weed
	<i>Hydrilla verticillata</i> (L.f.) Boyle	Hydrilla
Pontedariaceae (pickerelweed)	<i>Heteranthera dubia</i> (Jacquin) MacMillian	Water stargrass
Ceratophyllaceae (coontail)	<i>Ceratophyllum demersum</i> L.	Coontail
Trapaceae	<i>Trapa natans</i> L.	Water chestnut
Haloragaceae (watermilfoil)	<i>Myriophyllum spicatum</i> L.	Eurasian watermilfoil
Zosteraceae	<i>Zostera marina</i> (L.)	Eelgrass

APPENDIX B



APPENDIX B

Latitude and Longitude Coordinate Points Defining the 21 Chesapeake Bay Sections and Chincoteague Bay. (For Section Locations and Descriptions See Figure 7 and Table 3.)

	Latitude Deg Min	Longitude Deg Min		Latitude Deg Min	Longitude Deg Min
Sec. 1. Susquehanna Flats			Sec. 4. Chester River		
	39 27.00	76 10.00		39 00.00	76 20.00
	39 39.15	76 10.00		39 10.00	76 20.00
	39 39.15	75 51.00		39 09.25	76 16.00
	39 27.50	76 00.00		39 12.55	76 10.40
	39 26.50	76 01.31		39 20.00	76 00.00
				39 19.50	75 45.00
Sec. 2. Upper Eastern Shore				39 05.00	75 45.00
	39 10.00	76 20.00		39 05.00	76 00.00
	39 20.00	76 12.50		38 57.10	76 11.85
	39 26.50	76 01.31		39 00.00	76 19.10
	39 27.50	76 00.00		38 50.00	76 01.65
	39 39.15	75 51.00		39 05.00	76 00.00
	39 39.15	75 45.00		39 05.00	75 45.00
	39 19.50	75 45.00		38 45.00	75 45.00
	39 20.00	76 00.00		38 45.00	75 50.00
	39 12.55	76 10.40		38 21.93	75 55.00
	39 09.25	76 16.00		38 25.00	76 06.80
			Sec. 5 Central Western Shore		
Sec. 3. Upper Western Shore				38 42.90	76 35.00
	39 12.40	76 49.00		38 55.00	76 37.50
	39 30.00	76 20.00		39 12.40	76 49.00
	39 27.00	76 10.00		39 11.15	76 40.00
	39 26.50	76 01.31		39 06.82	76 35.40
	39 20.00	76 12.50		39 03.50	76 32.30
	39 10.00	76 20.00		39 00.00	76 20.00
	39 00.00	76 20.00		38 55.00	76 25.00
	39 03.50	76 32.30		38 45.00	76 25.00
	39 06.82	76 35.40			
	39 11.15	76 40.00			

Latitude Deg Min	Longitude Deg Min	Latitude Deg Min	Longitude Deg Min
Sec. 6 Eastern Bay		38 18.00	76 22.83
38 45.00	76 25.00	Sec. 9. Middle Western Shore	
38 55.00	76 25.00		
39 00.00	76 20.00	38 02.85	76 19.40
39 00.00	76 19.10	38 05.00	76 21.54
38 57.10	76 11.85	38 15.00	76 25.45
39 05.00	76 00.00	38 18.00	76 22.83
38 50.00	76 01.65	38 21.66	76 23.50
38 44.10	76 10.50	38 30.00	76 32.30
38 50.00	76 16.50	38 42.90	76 35.00
38 45.00	76 20.00	38 45.00	76 25.00
38 42.50	76 20.50	38 23.50	76 20.00
		38 05.00	76 10.00
Sec. 7. Choptank River			
38 23.50	76 20.00	Sec. 10. Lower Potomac River	
38 45.00	76 25.00		
38 42.50	76 20.50	37 53.40	76 14.45
38 45.00	76 20.00	37 55.50	76 18.15
38 50.00	76 16.50	37 53.85	76 28.00
38 44.10	76 10.50	38 06.15	76 53.00
38 50.00	76 01.65	38 15.00	77 06.40
39 05.00	76 00.00	38 20.00	77 09.40
39 05.00	75 45.00	38 24.20	77 14.08
38 45.00	75 45.00	38 35.00	77 00.00
38 45.00	75 50.00	38 15.00	76 25.45
38 21.93	75 55.00	38 05.00	76 21.54
38 25.00	76 06.80	38 02.85	76 19.40
		38 05.00	76 10.00
		37 50.00	76 10.00
Sec. 8. Patuxent River			
38 15.00	76 25.45	Sec. 11. Upper Potomac River	
38 35.00	77 00.00		
38 58.00	76 45.00	38 15.00	77 06.40
38 55.00	76 37.50	38 20.00	77 24.80
38 42.90	76 35.00	38 27.65	77 25.00
38 30.00	76 32.30	39 01.80	77 17.10
38 21.66	76 23.50	38 58.00	76 45.00
		38 35.00	77 00.00

Latitude Deg Min	Longitude Deg Min	Latitude Deg Min	Longitude Deg Min
Sec. 11. Upper Potomac River (continued)		38 00.73	75 49.50
		38 00.00	75 38.00
38 24.20	77 14.08	38 00.00	75 30.00
38 20.00	77 09.40	37 46.45	75 39.30
		37 20.00	75 55.50
Sec. 12. Middle Eastern Shore		Sec. 15. Reedville	
38 11.10	76 13.30	37 38.75	76 10.00
38 23.50	76 20.00	37 37.40	76 21.40
38 25.00	76 06.80	37 38.05	76 23.50
38 21.93	75 55.00	37 44.35	76 23.00
38 45.00	75 50.00	37 48.00	76 28.00
38 40.00	75 37.00	37 53.85	76 28.00
38 00.00	75 38.00	37 55.50	76 18.15
38 00.73	75 49.50	37 53.40	76 14.45
37 57.10	75 50.30	37 50.00	76 10.00
37 55.00	75 55.10		
38 11.70	75 59.00	Sec. 16. Rappahannock River Complex	
38 13.60	76 05.83	37 26.50	76 10.00
Sec. 13. Mid-Bay Island Complex		37 25.00	76 18.08
37 45.00	75 58.30	37 28.00	76 20.00
37 50.00	76 10.00	37 29.00	76 25.00
38 05.00	76 10.00	37 32.00	76 35.00
38 11.10	76 13.30	37 49.15	76 48.00
38 13.60	76 05.83	37 53.73	76 49.65
38 11.70	75 59.00	37 58.00	76 45.45
37 55.00	75 55.10	37 48.00	76 28.00
Sec. 14. Lower Eastern Shore		37 44.35	76 23.00
37 00.00	75 58.95	37 38.05	76 23.50
37 20.00	76 10.00	37 37.40	76 21.40
37 38.75	76 10.00	37 38.75	76 10.00
37 50.00	76 10.00	Sec. 17. New Point Comfort Region	
37 45.00	75 58.30	37 17.45	76 16.16
37 55.00	75 55.10	37 19.45	76 16.62
37 57.10	75 50.30		

	Latitude Deg Min	Longitude Deg Min	Latitude Deg Min	Longitude Deg Min
Sec. 17. New Point Comfort Region (continued)			37 16.50	76 28.50
			37 17.00	76 25.42
			37 16.25	76 22.50
37	20.00	76 17.40	37 17.00	76 19.33
37	21.00	76 17.40	37 14.00	76 19.33
37	22.25	76 19.50		
37	25.00	76 18.00		
37	26.50	76 10.00		
37	20.00	76 10.00		
Sec. 18. Mobjack Bay Complex			Sec. 20. Lower Western Shore	
			36 49.11	75 58.05
37	17.00	76 19.33	36 45.75	76 07.00
37	16.25	76 22.50	36 55.85	76 16.00
37	17.00	76 25.42	36 57.79	76 16.00
37	16.50	76 28.50	36 58.00	76 17.70
37	20.00	76 31.88	37 01.05	76 18.52
37	25.75	76 31.00	37 03.68	76 19.80
37	29.00	76 25.00	37 00.60	76 24.00
37	28.00	76 20.00	37 07.30	76 28.20
37	25.00	76 18.00	37 12.50	76 27.50
37	22.25	76 19.50	37 13.25	76 24.00
37	21.00	76 17.40	37 14.00	76 22.50
37	20.00	76 17.40	37 14.00	76 19.33
37	19.30	76 16.62	37 17.00	76 19.33
37	17.45	76 16.16	37 17.45	76 16.16
			37 20.00	76 10.00
			37 00.00	75 58.95
Sec 19. York River			Sec. 21. James River	
			36 45.75	76 07.00
37	14.00	76 22.50	36 40.00	76 10.00
37	13.25	76 24.00	36 40.00	76 30.00
37	12.50	76 27.50	36 40.00	76 40.00
37	07.30	76 28.20	36 55.63	76 40.00
37	14.00	76 36.50	37 17.30	77 18.00
37	16.72	76 43.65	37 20.15	77 14.00
37	26.29	76 49.77	37 27.45	77 08.10
37	30.55	76 40.00	37 26.29	76 49.77
37	28.56	76 35.00	37 16.72	76 43.65
37	20.00	76 31.88	37 14.00	76 36.50

Latitude Deg Min	Longitude Deg Min	Latitude Deg Min	Longitude Deg Min
Sec. 21. James River (continued)		Chincoteague Bay	
37 07.30	76 28.20	37 52.50	75 30.00
37 00.60	76 24.00	38 00.00	75 30.00
37 03.68	76 19.80	38 07.50	75 22.50
37 01.05	76 18.52	38 15.00	75 17.50
36 58.00	76 17.70	38 15.00	75 15.00
36 57.79	76 16.00	38 22.50	75 15.00
36 55.85	76 16.00	38 30.00	75 10.00
		38 30.00	75 02.50
		38 22.50	75 02.50
		38 15.00	75 07.50
		38 07.50	75 10.00
		38 00.00	75 15.00
		37 52.50	75 20.00
		37 51.00	75 22.30
		37 51.00	75 30.00


APPENDIX C

USGS 7.5 Minute Quadrangles for Chesapeake Bay and Chincoteague Bay Showing Distribution, Abundance, and Ground Truthing of SAV in 1993. [Boundaries of Individual SAV Beds Are Delineated by Solid Lines. Each Bed Is Identified with an Unique Two Letter (AA-ZA, AB-ZB, etc.) and One Number (1-4) Designation. These Numbers Represent the Density Classification Discussed in the Text and Figure 6, i.e. 1 = <10%; 2 = 10-40%; 3 = 40-70%; 4 = 70-100%. Ground Truthing is Represented by Symbols and Species Codes which Are Explained in the Legend.]

KEY FOR 1993 SAV MAPS


SPECIES

SURVEY STATIONS

Zm	<i>Zostera marina</i> (eelgrass)	
Rm	<i>Ruppia maritima</i> (widgeon grass)	▲ VIMS Field Survey
Ms	<i>Myriophyllum spicatum</i> (Eurasian watermilfoil)	* Harford Community College
Ppf	<i>Potamogeton perfoliatus</i> (redhead-grass)	● Citizens Field Observation
Ppc	<i>Potamogeton pectinatus</i> (sago pondweed)	✱ U.S. Fish and Wildlife Service
Zp	<i>Zannichellia palustris</i> (horned pondweed)	✱ U.S. Fish and Wildlife Service
N	<i>Najas</i> spp. (naiad)	◆ U.S. Geological Survey
Ec	<i>Elodea canadensis</i> (common elodea)	
Va	<i>Vallisneria americana</i> (wild celery)	
Tn	<i>Trapa natans</i> (water chestnut)	
Pe	<i>Potamogeton epihydrus</i> (leafy pondweed)	
Hv	<i>Hydrilla verticillata</i> (hydrilla)	
Hd	<i>Heteranthera dubia</i> (water stargrass)	
Per	<i>Potamogeton crispus</i> (curly pondweed)	
Cd	<i>Ceratophyllum demersum</i> (coontail)	
Ppu	<i>Potamogeton pusillus</i> (slender pondweed)	
Ngu	<i>Najas guadalupensis</i> (southern naiad)	 Indicates 'NO SAV' polygon
Ngr	<i>Najas gracillima</i> (slender naiad)	
C	<i>Chara</i> sp. (muskgrass)	
Nm	<i>Najas minor</i>	
Nfl	<i>Najas flexilis</i> (northern naiad)	
U	Unknown species composition	

SUBMERGED AQUATIC VEGETATION 1993

Aberdeen, Md. (002)



Scale (meters): 


Sources: Virginia Institute of Marine Science
U.S. Geological Survey

Date Flown: 9-6-93

Produced by:
Virginia Institute of Marine Science
School of Marine Science
College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Havre de Grace, Md. (003)





Scale (meters): 0 1000 2000 3000
 Sources: Virginia Institute of Marine Science
 U.S. Geological Survey

Produced by:
 Virginia Institute of Marine Science
 School of Marine Science
 College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

North East, Md. (004)



Scale (meters): 


Sources: Virginia Institute of Marine Science
 U.S. Geological Survey

Date Flown: 9-6-93

Produced by:
 Virginia Institute of Marine Science
 School of Marine Science
 College of William and Mary 105

SUBMERGED AQUATIC VEGETATION 1993

Perryman, Md. (008)



0 1000 2000 3000

Scale (meters):

Sources: Virginia Institute of Marine Science
U.S. Geological Survey

Produced by:

Virginia Institute of Marine Science
School of Marine Science
College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Spesutie, Md.(009)




Scale (meters): 0 1000 2000 3000
 Sources: Virginia Institute of Marine Science
 U.S. Geological Survey
 Date Flown: 9-6-93

Produced by:
 Virginia Institute of Marine Science
 School of Marine Science
 College of William and Mary 107

SUBMERGED AQUATIC VEGETATION 1993

Earleville, Md. (010)




Scale (meters): 0 1000 2000 3000


Sources: Virginia Institute of Marine Science
U.S. Geological Survey

Produced by:
Virginia Institute of Marine Science
School of Marine Science
College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Middle River, Md. (013)



Scale (meters): 


Sources: Virginia Institute of Marine Science
U.S. Geological Survey

Date Flown: 9-6-93

Produced by:
Virginia Institute of Marine Science
School of Marine Science
College of William and Mary 109

SUBMERGED AQUATIC VEGETATION 1993

Gunpowder Neck, Md. (014)



Scale (meters): 0 1000 2000 3000

Sources: Virginia Institute of Marine Science
U.S. Geological Survey

Date Flown: 9-6-93


Produced by:
Virginia Institute of Marine Science
School of Marine Science
College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Hanesville, Md. (015)



Scale (meters):



Sources: Virginia Institute of Marine Science
U.S. Geological Survey


Date Flown: 7-28-93

Produced by:

Virginia Institute of Marine Science
School of Marine Science
College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Betterton, Md. (016)




Scale (meters): 0 1000 2000 3000

Sources: Virginia Institute of Marine Science
U.S. Geological Survey

Produced by:
Virginia Institute of Marine Science
School of Marine Science
College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Galena, Md. (017)



Scale (meters): 0 1000 2000 3000


Sources: Virginia Institute of Marine Science
 U.S. Geological Survey

Date Flown: 9-6-93

Produced by:
 Virginia Institute of Marine Science
 School of Marine Science
 College of William and Mary 113

SUBMERGED AQUATIC VEGETATION 1993

Swan Point, Md. (020)




Scale (meters): 0 1000 2000 3000
Sources: Virginia Institute of Marine Science
U.S. Geological Survey

Produced by:
Virginia Institute of Marine Science
School of Marine Science
College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Rock Hall, Md. (021)



Scale (meters): 0 1000 2000 3000


Sources: Virginia Institute of Marine Science
 U.S. Geological Survey

Date Flown: 7-28-93

Produced by:
 Virginia Institute of Marine Science
 School of Marine Science
 College of William and Mary 115

SUBMERGED AQUATIC VEGETATION 1993

Round Bay, Md. (023)




Scale (meters): 0 1000 2000 3000

Sources: Virginia Institute of Marine Science
U.S. Geological Survey

Produced by:
Virginia Institute of Marine Science
School of Marine Science
College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Gibson Island, Md. (024)



0 1000 2000 3000

Scale (meters):

Sources: Virginia Institute of Marine Science

U.S. Geological Survey

Date Flown: 7-28-93

Produced by:


Virginia Institute of Marine Science

School of Marine Science

College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Langford Creek, Md. (026)




Scale (meters): 0 1000 2000 3000
 Sources: Virginia Institute of Marine Science
 U.S. Geological Survey
 Date Flown: 7-28-93

Produced by:
 Virginia Institute of Marine Science
 School of Marine Science
 College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Washington West, Md.-D.C.-Va. (028)



Scale (meters): 0 1000 2000 3000

Sources: Virginia Institute of Marine Science


U.S. Geological Survey

Date Flown: 9-12-93

Produced by:
 Virginia Institute of Marine Science
 School of Marine Science
 College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Washington East, D.C.-Md. (029)




Scale (meters): 0 1000 2000 3000
 Sources: Virginia Institute of Marine Science
 U.S. Geological Survey

Produced by:
 Virginia Institute of Marine Science
 School of Marine Science
 College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

South River, Md. (030)




Scale (meters): 0 1000 2000 3000
 Sources: Virginia Institute of Marine Science
 U.S. Geological Survey
 Date Flown: 7-17-93

Produced by:
 Virginia Institute of Marine Science
 School of Marine Science
 College of William and Mary 121

SUBMERGED AQUATIC VEGETATION 1993

Annapolis, Md. (031)




Scale (meters): 0 1000 2000 3000
 Sources: Virginia Institute of Marine Science
 U.S. Geological Survey

Produced by:
 Virginia Institute of Marine Science
 School of Marine Science
 College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Kent Island, Md. (032)



0 1000 2000 3000

Scale (meters):

Sources: Virginia Institute of Marine Science

U.S. Geological Survey

Date Flown: 7-16-93

Produced by:

Virginia Institute of Marine Science


School of Marine Science

College of William and Mary

123

SUBMERGED AQUATIC VEGETATION 1993

Queenstown, Md. (033)




Scale (meters): 0 1000 2000 3000
 Sources: Virginia Institute of Marine Science
 U.S. Geological Survey

Produced by:
 Virginia Institute of Marine Science
 School of Marine Science
 College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Alexandria, Va.-D.C.-Md. (034)




0 1000 2000 3000

Scale (meters):
 Sources: Virginia Institute of Marine Science
 U.S. Geological Survey
 Date Flown: 9-12-93

Produced by:
 Virginia Institute of Marine Science
 School of Marine Science
 College of William and Mary 125

SUBMERGED AQUATIC VEGETATION 1993

Claiborne, Md. (036)




Scale (meters): 0 1000 2000 3000
 Sources: Virginia Institute of Marine Science
 U.S. Geological Survey

Produced by:
 Virginia Institute of Marine Science
 School of Marine Science
 College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

St. Michaels, Md. (037)



Scale (meters): 0 1000 2000 3000


Sources: Virginia Institute of Marine Science
U.S. Geological Survey

Date Flown: 7-16-93

Produced by:
Virginia Institute of Marine Science
School of Marine Science
College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Fort Belvoir, Va.-Md. (039)



Scale (meters): 0 1000 2000 3000


Sources: Virginia Institute of Marine Science
U.S. Geological Survey

Date Flown: 9-12-93

Produced by:
Virginia Institute of Marine Science
School of Marine Science
College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Mt. Vernon, Md.-Va. (040)



0 1000 2000 3000

Scale (meters):

Sources: Virginia Institute of Marine Science

U.S. Geological Survey

Date Flown: 9-12-93

Produced by:


Virginia Institute of Marine Science

School of Marine Science

College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993


Lower Marlboro, Md. (041)



Scale (meters): 0 1000 2000 3000
 Sources: Virginia Institute of Marine Science
 U.S. Geological Survey

Produced by:
 Virginia Institute of Marine Science
 School of Marine Science
 College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993 Tilghman, Md. (043)




Scale (meters): 0 1000 2000 3000
 Sources: Virginia Institute of Marine Science
 U.S. Geological Survey
 Date Flown: 7-16-93

Produced by:
 Virginia Institute of Marine Science
 School of Marine Science
 College of William and Mary 131

SUBMERGED AQUATIC VEGETATION 1993

Oxford, Md. (044)





Scale (meters): 0 1000 2000 3000
 Sources: Virginia Institute of Marine Science
 U.S. Geological Survey

Produced by:
 Virginia Institute of Marine Science
 School of Marine Science
 College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Trappe, Md. (045)




Scale (meters): 
 Sources: Virginia Institute of Marine Science
 U.S. Geological Survey
 Date Flown: 7-18-93


Produced by:
 Virginia Institute of Marine Science
 School of Marine Science
 College of William and Mary 133

SUBMERGED AQUATIC VEGETATION 1993

Quantico, Va.-Md. (047)



Scale (meters):



Sources: Virginia Institute of Marine Science
U.S. Geological Survey


Date Flown: 9-12-93

Produced by:

Virginia Institute of Marine Science
School of Marine Science
College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Indian Head, Va.-Md. (048)



0 1000 2000 3000

Scale (meters):

Sources: Virginia Institute of Marine Science

U.S. Geological Survey

Date Flown: 9-12-93

Produced by:


Virginia Institute of Marine Science

School of Marine Science

College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Benedict, Md. (049)



Scale (meters): 0 1000 2000 3000


Sources: Virginia Institute of Marine Science
U.S. Geological Survey


Date Flown: 7-16-93

Produced by:
Virginia Institute of Marine Science
School of Marine Science
College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Hudson, Md. (051)




Scale (meters): 
 Sources: Virginia Institute of Marine Science
 U.S. Geological Survey
 Date Flown: 6-24-93

Produced by:
 Virginia Institute of Marine Science
 School of Marine Science
 College of William and Mary 137

SUBMERGED AQUATIC VEGETATION 1993

Church Creek, Md. (052)




Scale (meters): 0 1000 2000 3000
 Sources: Virginia Institute of Marine Science
 U.S. Geological Survey

Produced by:
 Virginia Institute of Marine Science
 School of Marine Science
 College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Cambridge, Md. (053)




Scale (meters): 0 1000 2000 3000
 Sources: Virginia Institute of Marine Science
 U.S. Geological Survey
 Date Flown: 7-18-93

Produced by:
 Virginia Institute of Marine Science
 School of Marine Science
 College of William and Mary 139

SUBMERGED AQUATIC VEGETATION 1993

Widewater, Va.- Md. (055)




Scale (meters): 0 1000 2000 3000

Sources: Virginia Institute of Marine Science
U.S. Geological Survey

Produced by:
Virginia Institute of Marine Science
School of Marine Science
College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Nanjemoy, Md. (056)



Scale (meters): 


Sources: Virginia Institute of Marine Science
U.S. Geological Survey

Date Flown: 9-12-93

Produced by:
Virginia Institute of Marine Science
School of Marine Science
College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Mathias Point, Md.- Va. (057)




Scale (meters): 0 1000 2000 3000
 Sources: Virginia Institute of Marine Science
 U.S. Geological Survey

Produced by:
 Virginia Institute of Marine Science
 School of Marine Science
 College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Popes Creek, Md. (058)



Scale (meters): 0 1000 2000 3000


Sources: Virginia Institute of Marine Science
U.S. Geological Survey

Date Flown: 7-16-93

Produced by:
Virginia Institute of Marine Science
School of Marine Science
College of William and Mary 143

SUBMERGED AQUATIC VEGETATION 1993

Broomes Island, Md. (060)




Scale (meters): 0 1000 2000 3000
 Sources: Virginia Institute of Marine Science
 U.S. Geological Survey

Produced by:
 Virginia Institute of Marine Science
 School of Marine Science
 College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Cove Point, Md.(061)




Scale (meters): 0 1000 2000 3000
 Sources: Virginia Institute of Marine Science
 U.S. Geological Survey
 Date Flown: 7-16-93

Produced by:
 Virginia Institute of Marine Science
 School of Marine Science
 College of William and Mary 145

SUBMERGED AQUATIC VEGETATION 1993

Taylor's Island, Md. (062)




Scale (meters): 0 1000 2000 3000
 Sources: Virginia Institute of Marine Science
 U.S. Geological Survey
 Date Flown: 6-24-93

Produced by:
 Virginia Institute of Marine Science
 School of Marine Science
 College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Golden Hill, Md. (063)



Scale (meters):




Sources: Virginia Institute of Marine Science
U.S. Geological Survey

Date Flown: 6-24-93

Produced by:
Virginia Institute of Marine Science
School of Marine Science
College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Passapatanzy, Md.-Va. (064)



Scale (meters): 0 1000 2000 3000


Sources: Virginia Institute of Marine Science
U.S. Geological Survey

Date Flown: 9-12-93

Produced by:
Virginia Institute of Marine Science
School of Marine Science
College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

King George, Va.-Md. (065)




Scale (meters): 0 1000 2000 3000
 Sources: Virginia Institute of Marine Science
 U.S. Geological Survey
 Date Flown: 9-12-93

Produced by:
 Virginia Institute of Marine Science
 School of Marine Science
 College of William and Mary 149

SUBMERGED AQUATIC VEGETATION 1993

Dahlgren, Va.-Md. (066)



Scale (meters): 0 1000 2000 3000


Sources: Virginia Institute of Marine Science
 U.S. Geological Survey

Date Flown: 7-16-93

Produced by:
 Virginia Institute of Marine Science
 School of Marine Science
 College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Colonial Beach North, Md.-Va. (067)



0 1000 2000 3000

Scale (meters):

Sources: Virginia Institute of Marine Science

U.S. Geological Survey

Date Flown: 7-16-93

Produced by:


Virginia Institute of Marine Science


School of Marine Science

College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Rock Point, Md. (068)




Scale (meters):  0 1000 2000 3000


Sources: Virginia Institute of Marine Science
U.S. Geological Survey

Produced by:
Virginia Institute of Marine Science
School of Marine Science
College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Hollywood, Md. (070)



Scale (meters): 


Sources: Virginia Institute of Marine Science
 U.S. Geological Survey

Date Flown: 7-16-93

Produced by:
 Virginia Institute of Marine Science
 School of Marine Science
 College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Solomons Island, Md. (071)




Scale (meters): 0 1000 2000 3000
 Sources: Virginia Institute of Marine Science
 U.S. Geological Survey

Produced by:
 Virginia Institute of Marine Science
 School of Marine Science
 College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Barren Island, Md. (072)




Scale (meters): 0 1000 2000 3000
 Sources: Virginia Institute of Marine Science
 U.S. Geological Survey
 Date Flown: 6-24-93

Produced by:
 Virginia Institute of Marine Science
 School of Marine Science
 College of William and Mary 155

SUBMERGED AQUATIC VEGETATION 1993

Honga, Md. (073)




Scale (meters): 0 1000 2000 3000
 Sources: Virginia Institute of Marine Science
 U.S. Geological Survey

Produced by:
 Virginia Institute of Marine Science
 School of Marine Science
 College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Wingate, Md. (074)



Scale (meters): 0 1000 2000 3000


Sources: Virginia Institute of Marine Science
U.S. Geological Survey

Date Flown: 6-23-93

Produced by:
Virginia Institute of Marine Science
School of Marine Science
College of William and Mary 157

SUBMERGED AQUATIC VEGETATION 1993

Colonial Beach South, Va.-Md. (076)



Scale (meters): 0 1000 2000 3000


Sources: Virginia Institute of Marine Science
U.S. Geological Survey

Date Flown: 7-17-93

Produced by:
Virginia Institute of Marine Science
School of Marine Science
College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

St. Clements Island, Va.-Md. (078)



Scale (meters): 0 1000 2000 3000


Sources: Virginia Institute of Marine Science
U.S. Geological Survey

Date Flown: 7-17-93

Produced by:
Virginia Institute of Marine Science
School of Marine Science
College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

St. Marys City, Md. (080)




Scale (meters):
 Sources: Virginia Institute of Marine Science
 U.S. Geological Survey

Produced by:
 Virginia Institute of Marine Science
 School of Marine Science
 College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Richland Point, Md. (082)




Scale (meters): 0 1000 2000 3000
Sources: Virginia Institute of Marine Science
U.S. Geological Survey
Date Flown: 6-24-93

Produced by:
Virginia Institute of Marine Science
School of Marine Science
College of William and Mary 161

SUBMERGED AQUATIC VEGETATION 1993

Bloodsworth Island, Md. (083)




Scale (meters): 0 1000 2000 3000
 Sources: Virginia Institute of Marine Science
 U.S. Geological Survey

Produced by:
 Virginia Institute of Marine Science
 School of Marine Science
 College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Deal Island, Md. (084)




Scale (meters): 0 1000 2000 3000
 Sources: Virginia Institute of Marine Science
 U.S. Geological Survey
 Date Flown: 6-23-93

Produced by:
 Virginia Institute of Marine Science
 School of Marine Science
 College of William and Mary 163

SUBMERGED AQUATIC VEGETATION 1993

Monie, Md. (085)




Scale (meters): 0 1000 2000 3000
 Sources: Virginia Institute of Marine Science
 U.S. Geological Survey

Produced by:
 Virginia Institute of Marine Science
 School of Marine Science
 College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

St. George Island, Va.-Md. (089)



Scale (meters): 0 1000 2000 3000


Sources: Virginia Institute of Marine Science
U.S. Geological Survey

Date Flown: 7-17-93

Produced by:
Virginia Institute of Marine Science
School of Marine Science
College of William and Mary 165

SUBMERGED AQUATIC VEGETATION 1993

Kedges Straits, Md. (091)




Scale (meters): 0 1000 2000 3000
 Sources: Virginia Institute of Marine Science
 U.S. Geological Survey

Produced by:
 Virginia Institute of Marine Science
 School of Marine Science
 College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Terrapin Sand Point, Md. (092)




Scale (meters): 0 1000 2000 3000

Sources: Virginia Institute of Marine Science
U.S. Geological Survey

Date Flown: 6-23-93

Produced by:
Virginia Institute of Marine Science
School of Marine Science
College of William and Mary 167

SUBMERGED AQUATIC VEGETATION 1993 Marion, Md. (093)



Scale (meters): 0 1000 2000 3000


Sources: Virginia Institute of Marine Science
U.S. Geological Survey

Date Flown: 6-24-93

Produced by:
Virginia Institute of Marine Science
School of Marine Science
College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Ewell, Md.-Va. (099)



Scale (meters): 0 1000 2000 3000


Sources: Virginia Institute of Marine Science
U.S. Geological Survey

Date Flown: 6-24-93

Produced by:
Virginia Institute of Marine Science
School of Marine Science
College of William and Mary 169

SUBMERGED AQUATIC VEGETATION 1993

Great Fox Island, Va.-Md. (100)




Scale (meters): 0 1000 2000 3000


Sources: Virginia Institute of Marine Science
U.S. Geological Survey

Produced by:
Virginia Institute of Marine Science
School of Marine Science
College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Crisfield, Md.-Va. (101)



Scale (meters): 


Sources: Virginia Institute of Marine Science
 U.S. Geological Survey

Date Flown: 6-24-93

Produced by:
 Virginia Institute of Marine Science
 School of Marine Science
 College of William and Mary 171

SUBMERGED AQUATIC VEGETATION 1993

Saxis, Va.-Md. (102)




Scale (meters): 0 1000 2000 3000

Sources: Virginia Institute of Marine Science
U.S. Geological Survey

Produced by:
Virginia Institute of Marine Science
School of Marine Science
College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Reedville, Va. (106)



Scale (meters): 0 1000 2000 3000


Sources: Virginia Institute of Marine Science
U.S. Geological Survey

Date Flown: 6-24-93

Produced by:
Virginia Institute of Marine Science
School of Marine Science
College of William and Mary 173

SUBMERGED AQUATIC VEGETATION 1993

Tangier Island, Va.(107)




Scale (meters): 0 1000 2000 3000
 Sources: Virginia Institute of Marine Science
 U.S. Geological Survey

Produced by:
 Virginia Institute of Marine Science
 School of Marine Science
 College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Chesconessex, Va. (108)



Scale (meters): 0 1000 2000 3000


Sources: Virginia Institute of Marine Science
U.S. Geological Survey

Date Flown: 6-7-93

Produced by:
Virginia Institute of Marine Science
School of Marine Science
College of William and Mary 175

SUBMERGED AQUATIC VEGETATION 1993

Parksley, Va. (109)




Scale (meters): 0 1000 2000 3000

Sources: Virginia Institute of Marine Science
U.S. Geological Survey

Produced by:
Virginia Institute of Marine Science
School of Marine Science
College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Irvington, Va. (111)



Scale (meters): 0 1000 2000 3000


Sources: Virginia Institute of Marine Science
 U.S. Geological Survey

Date Flown: 6-7-93

Produced by:
 Virginia Institute of Marine Science
 School of Marine Science
 College of William and Mary 177

SUBMERGED AQUATIC VEGETATION 1993

Fleets Bay, Va. (112)




Chesapeake Bay

Scale (meters): 0 1000 2000 3000
 Sources: Virginia Institute of Marine Science
 U.S. Geological Survey

Produced by:
 Virginia Institute of Marine Science
 School of Marine Science
 College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Nandua Creek, Va.(113)




Scale (meters): 0 1000 2000 3000
 Sources: Virginia Institute of Marine Science
 U.S. Geological Survey
 Date Flown: 6-7-93

Produced by:
 Virginia Institute of Marine Science
 School of Marine Science
 College of William and Mary 179

SUBMERGED AQUATIC VEGETATION 1993

Wilton, Va. (117)



Scale (meters): 0 1000 2000 3000


Sources: Virginia Institute of Marine Science
 U.S. Geological Survey

Date Flown: 5-27-93

Produced by:
 Virginia Institute of Marine Science
 School of Marine Science
 College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Deltaville, Va. (118)



Scale (meters): 0 1000 2000 3000


Sources: Virginia Institute of Marine Science
U.S. Geological Survey

Date Flown: 5-27-93

Produced by:
Virginia Institute of Marine Science
School of Marine Science
College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Jamesville, Va. (119)




Chesapeake Bay

Scale (meters): 0 1000 2000 3000
 Sources: Virginia Institute of Marine Science
 U.S. Geological Survey
 Date Flown: 5-27-93

Produced by:
 Virginia Institute of Marine Science
 School of Marine Science
 College of William and Mary 183

SUBMERGED AQUATIC VEGETATION 1993

Ware Neck, Va. (122)




Scale (meters): 0 1000 2000 3000
 Sources: Virginia Institute of Marine Science
 U.S. Geological Survey
 Date Flown: 6-10-93

Produced by:
 Virginia Institute of Marine Science
 School of Marine Science
 College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Mathews, Va. (123)



Scale (meters): 0 1000 2000 3000

Sources: Virginia Institute of Marine Science
U.S. Geological Survey


Date Flown: 5-27-93

Produced by:
Virginia Institute of Marine Science
School of Marine Science
College of William and Mary

185

SUBMERGED AQUATIC VEGETATION 1993

Franktown, Va. (124)



Scale (meters): 0 1000 2000 3000


Sources: Virginia Institute of Marine Science
U.S. Geological Survey

Date Flown: 5-27-93

Produced by:
Virginia Institute of Marine Science
School of Marine Science
College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Achilles, Va. (131)




Scale (meters):
 Sources: Virginia Institute of Marine Science
 U.S. Geological Survey
 Date Flown: 6-10-93

Produced by:
 Virginia Institute of Marine Science
 School of Marine Science
 College of William and Mary 187

SUBMERGED AQUATIC VEGETATION 1993

New Point Comfort, Va. (132)




Scale (meters): 0 1000 2000 3000
 Sources: Virginia Institute of Marine Science
 U.S. Geological Survey

Produced by:
 Virginia Institute of Marine Science
 School of Marine Science
 College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Cape Charles, Va. (133)




Scale (meters): 0 1000 2000 3000
 Sources: Virginia Institute of Marine Science
 U.S. Geological Survey
 Date Flown: 5-27-93

Produced by:
 Virginia Institute of Marine Science
 School of Marine Science
 College of William and Mary 189

SUBMERGED AQUATIC VEGETATION 1993

Cheriton, Va. (134)



Scale (meters): 0 1000 2000 3000


Sources: Virginia Institute of Marine Science
U.S. Geological Survey


Date Flown: 5-27-93

Produced by:
Virginia Institute of Marine Science
School of Marine Science
College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Yorktown, Va. (139)




Scale (meters): 
 Sources: Virginia Institute of Marine Science
 U.S. Geological Survey
 Date Flown: 6-10-93

Produced by:
 Virginia Institute of Marine Science
 School of Marine Science
 College of William and Mary 191

SUBMERGED AQUATIC VEGETATION 1993

Poquoson West, Va. (140)



Scale (meters): 0 1000 2000 3000


Sources: Virginia Institute of Marine Science
U.S. Geological Survey

Date Flown: 6-10-93


Produced by:
Virginia Institute of Marine Science
School of Marine Science
College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Poquoson East, Va. (141)



Scale (meters):



Sources: Virginia Institute of Marine Science

U.S. Geological Survey

Date Flown: 6-10-93


Produced by:

Virginia Institute of Marine Science

School of Marine Science

College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993 Elliotts Creek, Va. (142)



Scale (meters): 0 1000 2000 3000


Sources: Virginia Institute of Marine Science
U.S. Geological Survey

Date Flown: 5-27-93

Produced by:
Virginia Institute of Marine Science
School of Marine Science
College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Hampton, Va. (147)



Scale (meters): 0 1000 2000 3000


Sources: Virginia Institute of Marine Science
U.S. Geological Survey

Date Flown: 6-10-93

Produced by:
Virginia Institute of Marine Science
School of Marine Science
College of William and Mary 195

SUBMERGED AQUATIC VEGETATION 1993

Cape Henry, Va. (152)




Scale (meters): 0 1000 2000 3000

Sources: Virginia Institute of Marine Science
U.S. Geological Survey

Produced by:
Virginia Institute of Marine Science
School of Marine Science
College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Bristol, Md. (159)



Scale (meters): 0 1000 2000 3000


Sources: Virginia Institute of Marine Science
 U.S. Geological Survey

Date Flown: 7-16-93

Produced by:
 Virginia Institute of Marine Science
 School of Marine Science
 College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Port Tobacco, Md. (161)



Scale (meters): 


Sources: Virginia Institute of Marine Science
U.S. Geological Survey


Date Flown: 9-14-93

Produced by:
Virginia Institute of Marine Science
School of Marine Science
College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Charlotte Hall, Md. (162)




Scale (meters): 
 Sources: Virginia Institute of Marine Science
 U.S. Geological Survey
 Date Flown: 7-16-93

Produced by:
 Virginia Institute of Marine Science
 School of Marine Science
 College of William and Mary 199

SUBMERGED AQUATIC VEGETATION 1993

Assawoman Bay, Md. (166)




Scale (meters): 0 1000 2000 3000

Sources: Virginia Institute of Marine Science
U.S. Geological Survey

Produced by:
Virginia Institute of Marine Science
School of Marine Science
College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Berlin, Md. (167)




Scale (meters): 0 1000 2000 3000
 Sources: Virginia Institute of Marine Science
 U.S. Geological Survey
 Date Flown: 5-27-93

Produced by:
 Virginia Institute of Marine Science
 School of Marine Science
 College of William and Mary 201

SUBMERGED AQUATIC VEGETATION 1993

Ocean City, Md. (168)



Scale (meters): 0 1000 2000 3000


Sources: Virginia Institute of Marine Science
U.S. Geological Survey

Date Flown: 5-27-93

Produced by:
Virginia Institute of Marine Science
School of Marine Science
College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Tingles Island, Md. (170)



0 1000 2000 3000

Scale (meters):

Sources: Virginia Institute of Marine Science

U.S. Geological Survey

Date Flown: 5-27-93

Produced by:


Virginia Institute of Marine Science

School of Marine Science

College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Boxiron, Md.-Va. (172)




Scale (meters): 0 1000 2000 3000

Sources: Virginia Institute of Marine Science
U.S. Geological Survey

Produced by:
Virginia Institute of Marine Science
School of Marine Science
College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Whittington Point, Md.-Va. (173)



Scale (meters): 0 1000 2000 3000


Sources: Virginia Institute of Marine Science
U.S. Geological Survey

Date Flown: 5-27-93

Produced by:
Virginia Institute of Marine Science
School of Marine Science
College of William and Mary 205

SUBMERGED AQUATIC VEGETATION 1993

Chincoteague West, Va. (174)




Scale (meters): 0 1000 2000 3000
 Sources: Virginia Institute of Marine Science
 U.S. Geological Survey

Produced by:
 Virginia Institute of Marine Science
 School of Marine Science
 College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Chincoteague East, Va. (175)



Scale (meters): 0 1000 2000 3000


Sources: Virginia Institute of Marine Science
U.S. Geological Survey

Date Flown: 5-27-93

Produced by:
Virginia Institute of Marine Science
School of Marine Science
College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

East of New Point Comfort, Va. (177)




Sources: Virginia Institute of Marine Science
U.S. Geological Survey

Produced by:
Virginia Institute of Marine Science
School of Marine Science
College of William and Mary

SUBMERGED AQUATIC VEGETATION 1993

Bethel Beach, Va. (178)




Scale (meters): 0 1000 2000 3000
Sources: Virginia Institute of Marine Science
U.S. Geological Survey
Date Flown: 5-27-93

Produced by:
Virginia Institute of Marine Science
School of Marine Science
College of William and Mary 209

SUBMERGED AQUATIC VEGETATION 1993

Goose Island, Va. (179)




Scale (meters): 0 1000 2000 3000
Sources: Virginia Institute of Marine Science
U.S. Geological Survey

Produced by:
Virginia Institute of Marine Science
School of Marine Science
College of William and Mary

APPENDIX D

Number of Square Meters of SAV for Individual Beds and Totals for Density Categories for Each
USGS 7.5 Minute Quadrangle in 1993.



APPENDIX D

Number of Square Meters of SAV for Individual Beds and Totals for Density Categories for Each USGS 7.5 Minute Quadrangle in 1993. [See Maps in Appendix C for Location of Each Bed. Quadrangles Are Listed Numerically by VIMS Map Number. Slight Differences (1 Square Meter) in Quadrangle Totals from Density Totals Are Due to Rounding.]

Aberdeen, Md.		LA4	2,586
VIMS Map # 002		MA4	35,171
		NA4	2,224
AA4	10,623	OA4	3,327
BA4	3,965	PA4	7,987
CA4	4,605	QA4	7,730
DA4	3,231	RA4	83,478
EA3	38,260	SA4	10,085
FA4	1,985	TA3	176,400
GA4	4,004	UA3	27,446
HA4	4,950	VA3	7,336
IA4	5,202	WA4	36,877
JA4	5,310	XA4	13,206
		YA4	16,779
Density 1 =	0	ZA3	8,578
Density 2 =	0	AB4	45,036
Density 3 =	38,260	BB4	19,711
Density 4 =	<u>43,875</u>	CB4	2,797
		DB4	355
Total =	82,135	EB4	4,225
		FB4	6,399
Havre de Grace, Md.		GB4	8,972
VIMS Map # 003		HB4	10,674
		IB4	3,894
AA2	6,912	JB4	58,009
BA2	36,408	KB3	12,680
CA2	15,480	LB1	139,702
DA2	492,198	MB4	303,096
EA1	15,035,720	NB4	102,168
FA4	291,283	OB3	41,014
GA3	14,209	PB3	40,838
HA4	3,486	QB2	11,377
IA4	2,401	RB2	6,160
JA4	5,576		
KA4	187,414	Density 1 =	15,175,422

Density 2 = 568,534
 Density 3 = 328,501
 Density 4 = 1,274,946

Total = 17,347,404

North East, Md.
 VIMS Map # 004

AA2 16,494
 BA1 4,915
 CA1 35,490
 DA1 387,755
 EA1 5,404
 FA1 11,723

Density 1 = 445,288
 Density 2 = 16,494
 Density 3 = 0
 Density 4 = 0

Total = 461,782

Perryman, Md.
 VIMS Map # 008

AA4 11,853
 BA3 1,735
 CA2 22,322
 DA2 18,468
 EA2 4,915
 FA2 2,780
 GA2 947
 HA2 5,555
 IA2 11,749

Density 1 = 0
 Density 2 = 66,735
 Density 3 = 1,735
 Density 4 = 11,853

Total = 80,323

Spesutie, Md.
 VIMS Map # 009

AA1 2,853
 BA2 16,502
 CA3 42,960
 DA3 4,046
 EA4 2,461
 FA4 2,497
 GA1 6,993
 HA2 9,069
 IA2 7,266
 JA2 9,968
 KA2 1,222
 LA2 3,700
 MA2 5,471
 NA2 14,313
 OA2 14,797
 PA2 7,030
 QA2 3,856
 RA2 34,459
 SA4 52,102
 TA3 37,248
 UA2 12,322
 VA2 6,115
 WA3 3,084
 XA2 7,713
 YA3 39,576
 ZA3 12,957
 AB2 4,619
 BB2 2,146
 CB2 9,178
 DB2 5,491
 EB2 2,292
 FB2 2,997
 GB2 2,896
 HB3 22,824
 IB2 17,177
 JB2 26,946
 KB4 4,090
 LB2 1,965

Density 1 = 9,846

Density 2 =	229,511
Density 3 =	162,695
Density 4 =	<u>61,151</u>

Total =	463,203
---------	---------

Earleville, Md.
VIMS Map # 010

AA2	27,658
BA2	20,864
CA3	22,022
DA2	4,763
EA1	5,054
FA3	28,008
GA1	4,283
HA1	9,825
IA1	7,475
JA2	22,850
KA2	83,240
LA2	67,615
MA3	102,357
NA1	13,908
OA1	92,305
PA4	22,119

Density 1 =	132,850
Density 2 =	226,988
Density 3 =	152,387
Density 4 =	<u>22,119</u>

Total =	534,344
---------	---------

Middle River, Md.
VIMS Map # 013

AA1	23,977
BA2	17,951
CA2	3,797
DA2	8,927

Density 1 =	23,977
Density 2 =	30,675

Density 3 =	0
Density 4 =	<u>0</u>

Total =	54,652
---------	--------

Gunpowder Neck, Md.
VIMS Map # 014

AA2	7,645
BA3	71,640
CA3	109,485
DA2	11,623
EA2	5,954
FA2	79,626
GA1	7,153
HA2	130,536
IA3	42,345
JA2	12,071

Density 1 =	7,153
Density 2 =	247,455
Density 3 =	223,470
Density 4 =	<u>0</u>

Total =	478,077
---------	---------

Hanesville, Md.
VIMS Map # 015

AA2	22,156
BA2	17,506
CA2	2,748

Density 1 =	0
Density 2 =	42,410
Density 3 =	0
Density 4 =	<u>0</u>

Total =	42,410
---------	--------

Betterton, Md.
VIMS Map # 016

AA3	52,089
BA3	58,893
CA1	451,654
DA2	18,408
EA1	102,758
FA1	3,036
GA1	3,101

Density 1 =	560,549
Density 2 =	18,408
Density 3 =	110,982
Density 4 =	<u>0</u>
Total =	689,939

Galena, Md.
VIMS Map # 017

AA1	25,134
BA4	11,851
CA1	7,827

Density 1 =	32,961
Density 2 =	0
Density 3 =	0
Density 4 =	<u>11,851</u>

Total =	44,812
---------	--------

Swan Point, Md.
VIMS Map # 020

AA2	15,021
BA2	36,209
CA2	25,036
DA2	22,208
EA2	22,464
FA3	16,180
GA2	16,758
HA2	5,053
IA2	17,249

Density 1 =	0
-------------	---

Density 2 =	159,999
Density 3 =	16,180
Density 4 =	<u>0</u>

Total =	176,179
---------	---------

Rock Hall, Md.
VIMS Map # 021

AA3	8,980
BA3	32,168
CA3	45,905
DA3	17,046
EA2	51,725
FA2	56,875
GA3	18,826
HA3	52,529
IA2	27,962
JA2	14,316

Density 1 =	0
Density 2 =	150,877
Density 3 =	175,454
Density 4 =	<u>0</u>

Total =	326,331
---------	---------

Gibson Island, Md.
VIMS Map # 024

AA2	8,372
BA2	10,063
CA2	2,731
DA2	11,887
EA2	36,515
FA2	41,803
GA2	9,950
HA2	3,533
IA3	7,199

Density 1 =	0
Density 2 =	124,853
Density 3 =	7,199

Density 4 = 0

Total = 132,053

Langford Creek, Md.
VIMS Map # 026

AA2	181,677
BA3	89,381
CA2	32,436
DA1	8,524
EA2	20,726
FA3	2,755
GA3	252,439
HA3	20,997
IA3	66,840
JA2	266,908
KA2	15,768
LA3	593,427
MA2	89,117
NA2	94,264
OA3	711,225
PA2	163,026
QA3	17,679
RA2	58,953
SA4	1,812,486
TA1	682,127

Density 1 =	690,651
Density 2 =	922,874
Density 3 =	1,754,742
Density 4 =	<u>1,812,486</u>

Total = 5,180,753

Washington West, Md.-D.C.-Va.
VIMS Map # 028

AA3	16,659
BA4	16,876
CA3	15,279
DA3	2,272
EA4	4,128

FA2	2,751
GA4	5,456
HA4	32,943
IA4	43,663
JA4	20,559
KA4	49,172
LA3	16,506
MA3	25,891

Density 1 =	0
Density 2 =	2,751
Density 3 =	76,608
Density 4 =	<u>172,797</u>

Total = 252,155

Washington East, D.C.-Md.
VIMS Map # 029

AA4	7,488
Density 1 =	0
Density 2 =	0
Density 3 =	0
Density 4 =	<u>7,488</u>

Total = 7,488

Kent Island, Md.
VIMS Map # 032

AA1	34,778
BA2	54,915
CA2	63,033
DA2	7,827
EA3	106,443
FA2	36,311
GA1	260,838
HA2	250,947
IA2	32,062
JA1	118,582
KA1	23,046
LA1	2,723

MA1	64,349
NA3	130,882
OA1	74,770
PA1	97,131
QA3	66,661
RA1	17,845
SA2	84,933
TA2	15,598

Density 1 =	694,062
Density 2 =	545,625
Density 3 =	303,986
Density 4 =	<u>0</u>

Total = 1,543,674

Queenstown, Md.
VIMS Map # 033

AA1	82,905
BA1	85,278
CA3	9,866
DA1	317,597
EA1	186,622
FA1	28,630
GA1	65,561
HA1	139,901
IA3	247,181
JA3	45,585
KA2	6,589
LA3	64,659
MA3	119,707
NA3	21,071
OA1	59,738
PA3	44,134
QA1	139,346
RA2	73,307
SA2	46,490
TA3	29,929

Density 1 =	1,105,577
Density 2 =	126,386
Density 3 =	582,133

Density 4 = 0

Total = 1,814,096

Alexandria, Va.-D.C.-Md.
VIMS Map # 034

AA3	23,704
BA3	817
CA3	1,587
DA3	7,243
EA3	3,463
FA2	2,035
GA2	4,477
HA4	41,631
IA3	74,270
JA3	4,799
KA4	14,762
LA2	10,171
MA3	33,668
NA2	23,199
OA1	5,469
PA4	33,798
QA1	7,661
RA2	2,621
SA4	3,438
TA3	4,177
UA4	6,669
VA4	19,418
WA4	168,982
XA2	233,107
YA4	121,705
ZA3	29,752
AB4	38,480
BB3	4,751
CB3	2,193
DB3	19,891
EB3	5,525
FB3	587,493
GB3	50,115
HB2	61,661
IB4	28,147
JB4	30,838

KB3	20,022	VA2	9,071
LB4	2,318	WA3	28,831
MB4	33,840	XA3	42,923
NB3	3,975	YA1	1,197,597
OB4	1,055,214	ZA1	272,754
PB4	202,074	AB3	120,067
QB1	174,935	BB1	356,523
RB3	1,337	CB1	177,298
SB3	12,206	DB1	462,753
TB4	60,459	EB2	18,392
UB2	86,476	FB2	237,583
Density 1 =	188,065	Density 1 =	2,502,278
Density 2 =	423,748	Density 2 =	1,395,609
Density 3 =	890,989	Density 3 =	235,859
Density 4 =	<u>1,861,772</u>	Density 4 =	<u>132,326</u>
Total =	3,364,573	Total =	4,266,072

Claiborne, Md.
VIMS Map # 036

St. Michaels, Md.
VIMS Map # 037

AA2	143,972	AA4	11,129
BA2	88,346	BA4	42,944
CA2	135,678	CA4	21,301
DA2	285,527	DA4	12,731
EA2	135,821	EA2	166,524
FA4	23,457	FA3	30,561
GA1	35,354	GA3	13,271
HA2	40,316	HA3	8,981
IA2	23,088	IA3	19,766
JA2	26,420	JA3	14,473
KA4	6,020	KA4	67,074
LA3	14,147	LA2	37,823
MA4	33,135	MA3	310,417
NA3	5,556	NA2	28,588
OA4	23,949	OA2	9,212
PA3	14,131	PA2	31,559
QA3	10,205	QA3	5,347
RA4	45,765	RA4	17,657
SA2	215,538	SA4	118,499
TA2	19,168	TA1	400,146
UA2	16,690	UA1	240,525

VA1	90,839
WA2	39,829
XA4	127,926
YA1	38,523
ZA3	80,182
AB2	27,025
BB1	15,224
CB1	51,320
DB1	172,068
EB3	240,026
FB3	215,530
Density 1 =	1,008,645
Density 2 =	340,559
Density 3 =	938,553
Density 4 =	<u>419,261</u>
Total =	2,707,020

Fort Belvoir, Va.-Md.
VIMS Map # 039

AA2	16,453
BA2	12,429
CA3	11,791
DA4	147,365
EA1	259,001
FA2	113,325
GA3	260,311
HA3	238,904
IA4	54,361
Density 1 =	259,001
Density 2 =	142,207
Density 3 =	511,006
Density 4 =	<u>201,726</u>

Total = 1,113,940

Mt. Vernon, Va.-Md.
VIMS Map # 040

AA4	60,590
-----	--------

BA4	1,528,021
CA4	147,481
DA3	87,525
EA1	7,988
FA4	57,452
GA4	55,556
HA2	39,803
IA4	39,713
JA2	32,280
KA2	6,248
LA4	7,526
MA3	284,163
NA2	2,508
OA1	7,662
Density 1 =	15,650
Density 2 =	80,838
Density 3 =	371,688
Density 4 =	<u>1,896,340</u>

Total = 2,364,516

Lower Marlboro, Md.
VIMS Map # 041

AA2	87,762
Density 1 =	0
Density 2 =	87,762
Density 3 =	0
Density 4 =	<u>0</u>

Total = 87,762

Tilghman, Md.
VIMS Map # 043

AA2	81,660
BA2	61,123
CA2	146,359
DA2	47,620
EA2	65,504
FA2	550,611

GA2	145,389
HA2	1,448,200
IA3	214,135
JA2	9,854
KA3	639,001
LA2	301,328
MA2	49,803
NA4	9,663
OA2	5,868
PA2	37,103
QA2	126,616

Density 1 =	0
Density 2 =	3,077,036
Density 3 =	853,136
Density 4 =	<u>9,663</u>
Total =	3,939,835

Oxford, Md.
VIMS Map # 044

AA2	31,931
BA3	16,257
CA3	12,318
DA3	266,368
EA3	58,281
FA4	16,961
GA2	641,424
HA2	230,766
IA2	26,201
JA2	42,658
KA2	21,775
LA3	14,626
MA2	105,290
NA2	747,753
OA2	509,250
PA2	816,097
QA3	35,259
RA2	8,784
SA3	7,206
TA3	29,194
UA3	10,427

VA3	13,864
WA2	31,442
XA4	5,137
YA4	5,093
ZA4	8,120
AB4	6,319
BB4	39,685
CB2	173,149
DB4	152,335
EB3	131,276
FB2	145,470
GB1	82,614

Density 1 =	82,614
Density 2 =	3,531,989
Density 3 =	595,076
Density 4 =	<u>233,649</u>
Total =	4,443,327

Quantico, Va.-Md.
VIMS Map # 047

AA4	622,384
BA4	1,240,126
CA4	807,689
DA4	80,918
EA2	25,480
FA4	106,001
GA3	9,604
HA4	165,627
IA4	131,418
JA1	151,976
KA4	122,806
LA2	28,421
MA4	744,292
NA4	1,113,691
OA4	290,166
PA4	115,126
QA2	4,598
RA4	4,541
SA4	234,407

Density 1 =	151,976
Density 2 =	58,498
Density 3 =	9,604
Density 4 =	<u>5,779,192</u>
Total =	5,999,270

KA1	1,946,211
LA3	748,812
MA1	930,270
NA3	112,302
OA3	735,026
PA2	556,176

Indian Head, Md. - Va.
VIMS Map # 048

AA2	133,033
BA4	1,963,088
CA2	148,112
DA3	26,142
EA3	42,922
FA2	6,106
GA4	95,480
HA4	673,932
IA4	70,812
JA4	273,590
KA2	12,416
LA3	14,882

Density 1 =	0
Density 2 =	299,667
Density 3 =	83,946
Density 4 =	<u>3,076,902</u>

Total = 3,460,515

Hudson, Md.
VIMS Map # 051

AA2	40,112
BA2	122,265
CA2	48,812
DA2	217,725
EA2	15,136
FA2	13,268
GA3	6,002
HA2	109,399
IA3	51,465
JA2	20,521

Density 1 =	2,876,481
Density 2 =	1,143,415
Density 3 =	1,653,607
Density 4 =	<u>0</u>

Total = 5,673,503

Church Creek, Md.
VIMS Map # 052

AA2	228,801
BA2	586,192
CA1	438,145
DA2	127,628
EA2	4,744
FA2	11,661
GA2	11,711
HA2	30,762
IA3	18,535
JA2	32,606

Density 1 =	438,145
Density 2 =	1,034,105
Density 3 =	18,535
Density 4 =	<u>0</u>

Total = 1,490,785

Cambridge, Md.
VIMS Map # 053

AA1	40,220
Density 1 =	40,220
Density 2 =	0
Density 3 =	0

Density 4 = 0

Total = 40,220

Widewater, Va.-Md.
VIMS Map # 055

AA3 1,379
BA3 4,039
CA4 20,070
DA4 3,220
EA3 155,903
FA4 236,839
GA2 141,296
HA4 49,146
IA1 224,465
JA4 3,804,912
KA4 66,323
LA4 148,842
MA2 37,971
NA4 432,110
OA1 27,357
PA4 793,190
QA3 4,159
RA4 76,065
SA3 7,142

Density 1 = 251,821
Density 2 = 179,267
Density 3 = 172,622
Density 4 = 5,630,716

Total = 6,234,427

Nanjemoy, Md.
VIMS Map # 056

AA4 395,871
BA4 9,985
CA4 10,290
DA3 3,360
EA2 55,501
FA2 23,469

GA4 5,122
HA4 35,538
IA4 20,877
JA4 33,261
KA4 40,129
LA4 16,166
MA4 26,413
NA2 34,780
OA2 33,306
PA3 2,148
QA2 7,537
RA3 2,697
SA2 130,386
TA2 3,225

Density 1 = 0
Density 2 = 288,203
Density 3 = 8,205
Density 4 = 593,652

Total = 890,060

Mathias Point, Md.-Va.
VIMS Map # 057

AA1 92,763
BA2 172,896
CA2 13,208
DA3 118,509
EA4 165,580
FA3 43,772
GA4 134,996
HA2 27,214
IA4 113,987
JA2 5,638
KA4 53,163
LA4 207,900
MA3 68,539
NA4 519,039
OA3 229,039
PA2 28,926
QA2 40,249
RA3 7,552

SA4	1,286
TA4	11,037
UA2	5,850
VA3	71,654
WA3	50,314
XA4	86,643
YA2	52,804
ZA4	121,215
AB4	32,165
BB3	1,152
CB3	1,230
DB2	5,673
EB4	5,464
FB3	38,939

Density 1 =	92,763
Density 2 =	352,459
Density 3 =	630,701
Density 4 =	<u>1,452,474</u>

Total = 2,528,397

Popes Creek, Md.
VIMS Map # 058

AA2	11,198
BA2	4,042

Density 1 =	0
Density 2 =	15,240
Density 3 =	0
Density 4 =	<u>0</u>

Total = 15,240

Taylors Island, Md.
VIMS Map # 062

AA2	2,643
BA2	23,766
CA2	41,110
DA2	208,877
EA2	62,213

FA2	497,848
GA2	165,481

Density 1 =	0
Density 2 =	1,001,938
Density 3 =	0
Density 4 =	<u>0</u>

Total = 1,001,938

Golden Hill, Md.
VIMS Map # 063

AA1	221,952
BA1	92,852
CA2	285,266
DA2	8,554
EA2	26,300
FA1	16,078

Density 1 =	330,882
Density 2 =	320,120
Density 3 =	0
Density 4 =	<u>0</u>

Total = 651,002

Passapatanzy, Md.-Va.
VIMS Map # 064

AA3	14,729
BA2	23,718
CA4	8,316
DA4	19,214

Density 1 =	0
Density 2 =	23,718
Density 3 =	14,729
Density 4 =	<u>27,530</u>

Total = 65,977

King George, Va.-Md.

VIMS Map # 065

AA3	301,022
BA4	281,725
CA4	207,932
Density 1 =	0
Density 2 =	0
Density 3 =	301,022
Density 4 =	<u>489,657</u>
Total =	790,678

Dahlgren, Va.-Md.
VIMS Map # 066

AA4	7,549
BA2	1,298
CA3	2,399
DA3	2,149
EA4	4,732
FA2	200,771
GA4	66,813
Density 1 =	0
Density 2 =	202,069
Density 3 =	4,548
Density 4 =	<u>79,095</u>
Total =	285,711

Colonial Beach North, Va.-Md.
VIMS Map # 067

AA3	30,754
BA3	91,212
CA4	1,897
DA4	9,310
EA2	7,376
FA3	45,228
GA3	71,195
HA3	67,588
IA2	112,948

JA4	58,263
KA2	2,265

Density 1 =	0
Density 2 =	122,589
Density 3 =	305,976
Density 4 =	<u>69,470</u>
Total =	498,036

Solomons Island, Md.
VIMS Map # 071

AA3	9,946
Density 1 =	0
Density 2 =	0
Density 3 =	9,946
Density 4 =	<u>0</u>
Total =	9,946

Barren Island, Md.
VIMS Map # 072

AA2	269,424
BA2	51,814
CA2	1,737,049
Density 1 =	0
Density 2 =	2,058,287
Density 3 =	0
Density 4 =	<u>0</u>

Total =	2,058,287
---------	-----------

Honga, Md.
VIMS Map # 073

AA2	54,418
BA3	111,650
CA2	150,599
DA3	101,676

EA4	129,423
FA3	81,309
GA3	12,278
HA3	136,613
IA2	83,842
JA2	30,691
KA2	115,130
LA2	35,841
MA2	556,600
NA2	700,560
OA2	444,626
PA1	79,705
QA3	196,656
RA2	708,805
SA2	148,834
TA2	1,096,728
UA2	519,629
VA2	73,193
WA4	649,376
XA2	453,334
YA3	2,218,390
ZA1	117,741
AB2	69,271
BB2	125,357
CB3	99,749
DB2	302,853
EB4	43,244
FB2	115,059
GB4	1,832,192
HB1	142,651
IB3	506,015
JB2	581,679
KB3	83,414
LB4	397,950
MB3	19,169
NB1	77,979
Density 1 =	418,076
Density 2 =	6,367,050
Density 3 =	3,566,918
Density 4 =	<u>3,052,184</u>
Total =	13,404,228

Wingate, Md.
VIMS Map # 074

AA2	14,888
BA3	163,024
CA2	177,955
DA1	131,752
EA2	121,494
FA4	1,253,053
GA3	326,920
HA3	3,019,290
IA3	200,478
Density 1 =	131,752
Density 2 =	314,337
Density 3 =	3,709,712
Density 4 =	<u>1,253,053</u>
Total =	5,408,854

St. Marys City, Md.
VIMS Map # 080

AA2	88,860
BA2	33,727
Density 1 =	0
Density 2 =	122,587
Density 3 =	0
Density 4 =	<u>0</u>
Total =	122,587

Richland Point, Md.
VIMS Map # 082

AA2	272,674
BA1	94,232
CA2	43,738
Density 1 =	94,232
Density 2 =	316,412
Density 3 =	0

Density 4 = 0

Total = 410,644

Bloodsworth Island, Md.
VIMS Map # 083

AA2	3,848,825
BA3	193,864
CA1	46,994
DA1	27,625
EA2	1,352,151
FA2	237,870
GA2	297,033
HA1	125,226
IA2	1,116,782
JA1	48,791
KA1	17,532
LA1	4,827
MA3	665,084
NA1	160,198
OA1	106,920
PA2	158,280
QA3	19,958
RA2	65,406
SA2	75,822
TA2	24,832
UA2	21,467
VA2	12,104
WA2	3,213

Density 1 = 538,114

Density 2 = 7,213,787

Density 3 = 878,907

Density 4 = 0

Total = 8,630,807

Deal Island, Md.
VIMS Map # 084

AA1	13,881
BA3	317,222

CA2 38,905

DA2 61,792

EA2 9,656

FA2 3,263

GA2 326,168

Density 1 = 13,881

Density 2 = 439,784

Density 3 = 317,222

Density 4 = 0

Total = 770,887

Monie, Md.
VIMS Map # 085

AA2 40,900

BA2 21,179

CA1 8,790

Density 1 = 8,790

Density 2 = 62,079

Density 3 = 0

Density 4 = 0

Total = 70,868

St. George Island, Md.-Va.
VIMS Map # 089

AA2 29,998

BA2 12,628

Density 1 = 0

Density 2 = 42,625

Density 3 = 0

Density 4 = 0

Total = 42,625

Kedges Straits, Md.
VIMS Map # 091

AA3	72,589
BA2	80,327
CA4	153,233
DA4	2,718,547
EA2	403,684
FA2	107,149
GA3	281,792
HA3	1,464,037
IA2	404,811
JA2	52,418
KA2	458,640
LA2	23,475
MA4	471,069
NA2	4,130
OA3	804,109
PA2	546,099
QA2	126,753
RA2	60,367
SA3	256,227
TA2	389,167
UA3	150,207
VA3	5,565

Density 1 =	0
Density 2 =	2,657,022
Density 3 =	3,034,526
Density 4 =	<u>3,342,848</u>

Total = 9,034,396

Terrapin Sand Point, Md.
VIMS Map # 092

AA2	28,535
BA3	1,642,170
CA1	151,415
DA3	319,546
EA4	73,255
GA3	16,748
HA2	10,795
IA3	36,889
JA1	34,405

Density 1 =	185,820
Density 2 =	39,330
Density 3 =	2,015,353
Density 4 =	<u>73,255</u>

Total = 2,313,758

Marion, Md.
VIMS Map # 093

AA2	6,318
BA2	3,808
CA2	5,916
DA2	19,783
EA2	196,379
FA3	118,760
GA2	123,843
HA4	83,722
IA2	174,969
JA2	30,656
KA3	600,519
LA2	15,759
MA2	24,666
NA3	3,044
OA2	8,390
PA2	15,019
QA3	105,620
RA2	132,729
SA3	290,373
TA1	108,946
UA2	41,187
VA2	74,382
WA4	14,717
XA3	69,264
YA3	181,703
ZA1	78,725
AB2	58,028
BB3	6,402
CB3	71,431
DB2	27,887
EB2	227,259
FB2	10,494
GB2	8,716

Density 1 =	187,671
Density 2 =	1,206,188
Density 3 =	1,447,116
Density 4 =	<u>98,439</u>

Total =	2,939,413
---------	-----------

Ewell, Md.-Va.
VIMS Map # 099

AA3	216,266
BA2	115,887
CA4	101,709
DA4	29,564
EA4	872,698
FA4	32,024
GA3	202,924
HA1	71,250
IA2	56,683
JA2	110,220
KA2	430,638
LA3	130,185
MA2	123,674
NA4	136,823
OA3	67,488
PA3	117,019
QA2	63,052
RA2	2,748,561
SA2	5,707,851
TA2	33,945
UA3	559,630
VA2	16,022
WA2	1,656,751
XA1	3,366,268
YA2	639,625
ZA4	6,159,700

Density 1 =	3,437,518
Density 2 =	11,702,910
Density 3 =	1,293,511
Density 4 =	<u>7,332,517</u>

Total =	23,766,456
---------	------------

Great Fox Island, Md.-Va.
VIMS Map # 100

AA2	2,007,255
BA2	555,855
CA2	127,135
DA2	2,168,475
EA3	615,837
FA1	15,995
GA1	14,836
HA1	14,926
IA3	145,223
JA1	69,010
KA4	79,210
LA3	691,348
MA2	140,172
NA2	250,552
OA2	640,730
PA3	177,253
QA4	3,562,095
RA2	3,555,152

Density 1 =	114,766
Density 2 =	9,445,325
Density 3 =	1,629,661
Density 4 =	<u>3,641,305</u>

Total =	14,831,057
---------	------------

Crisfield, Md.-Va.
VIMS Map # 101

AA2	58,051
BA2	652,694
CA3	792,589
DA2	7,740
EA2	276,746
FA3	179,214
GA2	63,931
HA3	87,440
IA2	96,408
JA2	14,482
KA2	13,964

LA2	17,092
MA2	98,419
NA3	8,979
OA2	114,745
PA2	6,785
QA3	116,207
RA2	3,101
SA4	92,696
TA3	22,296
UA2	3,972
VA2	3,048
WA2	31,392
XA2	45,839
YA4	367,981
ZA4	216,961

Density 1 =	0
Density 2 =	1,508,411
Density 3 =	1,206,724
Density 4 =	<u>677,637</u>

Total = 3,392,772

Saxis, Va.-Md.
VIMS Map # 102

AA3	2,199
BA3	11,247
CA3	4,770
DA2	2,774

Density 1 =	0
Density 2 =	2,774
Density 3 =	18,216
Density 4 =	<u>0</u>

Total = 20,989

Reedville, Va.
VIMS Map # 106

AA1	26,618
BA2	153,070

CA4	1,332,697
DA1	612,218
EA2	465,916
FA4	271,231
GA2	161,507
HA2	18,384

Density 1 =	638,835
Density 2 =	798,877
Density 3 =	0
Density 4 =	<u>1,603,928</u>

Total = 3,041,641

Tangier Island, Va.
VIMS Map # 107

AA3	479,651
BA2	111,017
CA4	1,387,394
DA2	33,520
EA2	116,127
FA2	55,185
GA2	112,291
HA2	858,845
IA2	2,040,244
JA3	522,678

Density 1 =	0
Density 2 =	3,327,229
Density 3 =	1,002,329
Density 4 =	<u>1,387,394</u>

Total = 5,716,952

Chesconessex, Va.
VIMS Map # 108

AA2	162,709
BA1	112,141
CA3	149,106
DA2	19,727
EA2	146,537

FA3	1,483	Parksley, Va.	
GA2	723	VIMS Map # 109	
HA3	447		
IA3	351	AA2	8,433
JA3	173,465	BA2	165,552
KA2	142,068	CA2	24,829
LA3	688,124	DA2	19,424
MA1	225,876	EA3	1,444
NA2	125,414	FA2	513,349
OA3	290,171	GA4	2,582,470
PA2	12,396	HA2	112,918
QA4	777,968	IA2	3,003
RA2	85,643	JA2	168,021
SA4	393,952	KA3	172,574
TA3	473,711	LA2	53,554
UA1	409,642	MA3	19,941
VA1	568,709	NA2	81,089
WA3	444,519	OA3	273,809
XA4	532,651	PA2	190,562
YA2	374,591	QA4	154,546
ZA4	326,647	RA3	246,851
AB2	443,885	SA2	17,035
BB4	814,470	TA2	1,615
CB3	65,346	UA2	2,854
DB3	4,215	VA2	14,304
EB3	4,311	WA2	8,237
FB4	70,933	XA3	38,561
GB4	716,478	YA2	2,505
HB2	548,499	ZA3	3,969
IB3	1,024,603	AB2	833
JB1	275,802	BB3	98,369
KB2	51,545	CB2	56,746
LB2	4,265	DB3	17,865
MB2	125,657	EB3	15,896
NB4	125,546	FB1	34,567
OB2	125,221		
Density 1 =	1,592,169	Density 1 =	34,567
Density 2 =	2,368,882	Density 2 =	1,444,863
Density 3 =	3,319,851	Density 3 =	889,278
Density 4 =	<u>3,758,645</u>	Density 4 =	<u>2,737,016</u>
Total =	11,039,547	Total =	5,105,725

Irvington, Va.
VIMS Map # 111

AA2	981
BA2	5,187
CA2	87,812
DA2	4,483
EA2	2,496
FA2	17,581
GA3	14,755
HA2	472,287
IA2	271,416
JA2	219,124
KA2	35,080
LA2	57,775
MA2	81,547
NA2	36,389
OA2	3,199
PA2	28,186
QA2	4,677
RA2	1,208
SA2	1,190
TA2	764
UA3	42,769
VA1	7,933
WA2	6,911
XA2	35,968
YA2	108,791
ZA2	50,866
Density 1 =	7,933
Density 2 =	1,533,921
Density 3 =	57,524
Density 4 =	<u>0</u>

Total = 1,599,378

Fleets Bay, Va.
VIMS Map # 112

AA2	1,092,599
BA3	95,883
CA2	31,648

DA2	2,419
EA3	325,705
FA1	429,423
GA2	154,314
HA4	28,534
IA2	337,280
JA2	8,777
KA2	26,167
LA2	22,850
MA2	22,878
NA2	47,720
OA2	64,204
PA3	990,149
QA1	186,208
RA3	380,626
SA2	113,021
TA3	11,004
UA2	27,819
VA2	17,176
WA2	12,294
XA2	298,697
YA2	119,261
ZA2	6,050
AB2	80,585
BB2	110,806
CB2	33,766
DB1	11,359
Density 1 =	626,990
Density 2 =	2,630,332
Density 3 =	1,803,367
Density 4 =	<u>28,534</u>

Total = 5,089,223

Nandua Creek, Va.
VIMS Map # 113

AA1	38,867
BA2	54,131
CA1	89,616
DA3	111,926
EA2	2,079

FA1	213,618
GA4	551,257
HA2	1,229,227
IA4	215,427
JA3	101,253
KA1	855,446
LA4	436,854
MA2	776,376

Density 1 =	1,197,547
Density 2 =	2,061,813
Density 3 =	213,179
Density 4 =	<u>1,203,538</u>

Total =	4,676,077
---------	-----------

Pungoteague, Va.
VIMS Map # 114

AA3	106,434
BA4	66,930
CA2	560,107
DA2	9,534
EA2	8,819
FA2	70,767
GA4	299,851
HA2	66,845
IA2	43,294
JA2	3,634
KA1	23,412
LA4	3,634,723
MA2	76,428
NA1	29,303
OA2	2,229,186
PA2	65,632
QA3	54,988
RA1	165,217
SA2	111,687
TA3	286,918
UA2	356,509
VA3	312,288
WA2	1,837
XA2	820

YA3	3,503
ZA2	19,047
AB3	14,553
BB2	8,760
CB3	9,791
DB2	18,817
EB2	118,470
FB3	813
GB3	1,227
HB3	47,038
IB2	8,141
JB3	77,540
KB2	243,124
LB1	320,040
MB2	365,800
NB1	18,127
OB2	28,441
PB3	34,042
QB2	120,368
RB2	9,421
SB2	30,932

Density 1 =	556,098
Density 2 =	4,576,419
Density 3 =	949,135
Density 4 =	<u>4,001,505</u>

Total =	10,083,157
---------	------------

Wilton, Va.
VIMS Map # 117

AA2	45,580
BA2	10,833
CA2	6,145
DA2	35,768
EA2	91,613
FA2	3,020
GA2	136,066
HA2	4,291
IA1	32,147
JA1	3,175
KA1	50,249

LA2	8,174	Density 1 =	448,687
MA2	13,807	Density 2 =	1,322,811
Density 1 =	85,571	Density 3 =	105,289
Density 2 =	355,296	Density 4 =	<u>291,568</u>
Density 3 =	0	Total =	2,168,355
Density 4 =	<u>0</u>		
Total =	440,867		

Jamesville, Va.
VIMS Map # 119

Deltaville, Va.
VIMS Map # 118

AA2	10,146	AA2	42,545
BA1	41,927	BA1	141,125
CA2	373,342	CA2	604,008
DA3	105,289	DA4	846,200
EA1	164,164	EA1	31,080
FA2	55,097	FA2	85,918
GA4	160,976	GA1	46,281
HA2	68,287	HA2	89,517
IA2	247,642	IA2	9,525
JA2	6,435	JA2	66,392
KA2	39,553	KA3	3,193
LA2	30,121	LA4	19,584
MA2	108,574	MA3	79,992
NA2	8,725	NA2	1,142,351
OA2	891	OA4	186,740
PA2	21,855	PA1	420,512
QA2	75,863	QA2	271,411
RA2	14,165	RA2	16,820
SA2	15,327	SA2	72,575
TA2	18,093	TA4	520,717
UA2	70,337	UA1	25,096
VA2	29,921	VA3	214,992
WA2	19,234	WA2	3,002
XA2	1,713	XA2	705,797
YA2	30,270	YA4	194,730
ZA2	2,377	ZA2	189,598
AB2	6,294	AB4	294,521
BB2	68,552	BB1	11,115
CB1	242,596	CB1	80,152
DB4	130,592	DB4	148,988
		EB2	198,661
		FB2	2,099
		GB2	72,988

Density 1 =	755,362
Density 2 =	3,573,205
Density 3 =	298,177
Density 4 =	<u>2,211,479</u>

Total = 6,838,223

Ware Neck, Va.
VIMS Map # 122

AA2	4,757
BA2	518,979
CA2	40,723
DA2	200,797
EA3	248,308
FA1	53,550
GA3	81,950
HA2	105,225
IA2	106,296
JA2	165,508
KA3	154,296
LA4	478,619
MA2	301,667
NA4	196,776
OA4	106,426
PA3	367,057

Density 1 =	53,550
Density 2 =	1,443,954
Density 3 =	851,612
Density 4 =	<u>781,821</u>

Total = 3,130,937

Mathews, Va.
VIMS Map # 123

AA1	3,070
BA1	9,290
CA3	440,487
DA3	20,681
EA2	123,206
FA2	61,728

GA2	25,549
HA2	26,579
IA4	180,713
JA4	54,320
KA2	214,113
LA2	75,055
MA2	51,958
NA4	36,990
OA4	30,952
PA2	5,946
QA1	38,463
RA2	209,846
SA4	612
TA4	19,129
UA4	8,417
VA4	57,530
WA2	151,660
XA2	61,709
YA2	50,174
ZA3	88,335
AB1	53,335
BB3	662,203
CB2	121,522
DB4	131,354
EB2	20,498
FB2	190,656
GB3	698,931
HB1	33,783

Density 1 =	137,941
Density 2 =	1,390,198
Density 3 =	1,910,636
Density 4 =	<u>520,017</u>

Total = 3,958,793

Franktown, Va.
VIMS Map # 124

AA4	108,305
BA1	912,909
CA3	56,157
DA4	300,471

EA2	61,588	Total =	7,678,399
FA2	58,182		
GA3	748,442	Achilles, Va.	
HA3	23,366	VIMS Map # 131	
IA3	145,378		
JA2	32,340	AA4	65,377
KA2	33,058	BA4	59,867
LA2	4,282	CA4	80,012
MA4	214,354	DA4	1,181,695
NA2	6,662	EA2	229,512
OA2	200,981	FA4	61,069
PA4	1,535,808	GA4	1,325,538
QA2	483,758	HA2	142,564
RA1	52,151	IA2	178,298
SA1	56,563	JA4	67,445
TA2	85,401	KA2	1,023
UA1	428,174	LA2	5,967
VA3	64,722	MA2	598
WA3	116,720	NA2	14,648
XA1	303,037	OA4	1,270,356
YA4	328,033	PA2	26,231
ZA2	147,831	QA4	1,378,495
AB4	140,078	RA3	445,523
BB2	67,563	SA4	7,047
CB2	27,962	TA2	1,042
DB2	16,221	UA4	234,301
EB2	13,104	VA2	26,253
FB3	152,500	WA4	19,282
GB2	24,113	XA2	32,445
HB2	99,752	YA4	277,911
IB2	13,129	ZA2	64,831
JB2	21,419	AB4	1,986,907
KB2	4,355	BB2	127,967
LB3	183,169	CB2	3,587
MB2	51,834	DB2	103,580
NB2	127,734	EB2	10,606
OB1	112,966	FB4	48,705
PB2	113,826	GB2	21,616
		HB2	67,485
		IB2	217,672
Density 1 =	1,865,799	JB4	782,882
Density 2 =	1,695,097	KB2	10,003
Density 3 =	1,490,454		
Density 4 =	<u>2,627,049</u>		

Density 1 =	0
Density 2 =	1,285,928
Density 3 =	445,523
Density 4 =	<u>8,846,888</u>

Total = 10,578,338

New Point Comfort, Va.
VIMS Map # 132

AA2	367,286
BA2	291,407
CA4	5,247,032
DA4	81,805
EA4	651,126
FA2	330,292
GA3	174,286
HA3	307,747
IA4	1,172,207
JA2	50,341
KA4	855,557
LA1	114,714
MA1	69,400
NA4	990,454
OA2	132,173
PA1	476,262
QA4	54,856
RA1	132,645
SA2	739,793
TA2	24,714
UA1	6,431
VA1	1,525
WA4	368,199
XA3	360,329
YA4	223,432
ZA3	112,950
AB2	33,422
BB3	1,549,208
CB1	12,946
DB4	99,006

Density 1 = 813,922
Density 2 = 1,969,427

Density 3 =	2,504,520
Density 4 =	<u>9,743,674</u>

Total = 15,031,543

Cape Charles, Va.
VIMS Map # 133

AA2	265,108
BA4	1,435
CA4	19,220
DA2	141,625
EA4	249,818
FA2	41,638
GA2	28,704
HA2	1,408
IA3	13,867
JA2	70,611
KA2	384,274
LA4	609,470
MA2	34,877
NA2	47,262
OA2	149,099
PA1	533,424
QA3	213,529
RA2	865,281
SA3	919,985
TA2	67,170

Density 1 = 533,424
Density 2 = 2,097,057
Density 3 = 1,147,382
Density 4 = 879,943

Total = 4,657,806

Cheriton, Va.
VIMS Map # 134

AA3	5,954
BA2	240,809
CA4	367,599
DA1	72,836

EA1	254,447
FA2	26,553
Density 1 =	327,284
Density 2 =	267,361
Density 3 =	5,954
Density 4 =	<u>367,599</u>
Total =	968,198

Yorktown, Va.
VIMS Map # 139

AA1	926
BA2	8,840
CA1	4,160
DA2	3,138
EA2	3,193
FA1	4,914
Density 1 =	10,000
Density 2 =	15,172
Density 3 =	0
Density 4 =	<u>0</u>
Total =	25,172

Poquoson West, Va.
VIMS Map # 140

AA4	32,214
BA4	31,315
CA4	84,533
DA3	436,065
EA2	175,275
FA4	443,452
GA1	92,844
HA2	28,001
IA3	163,205
JA2	21,829
KA2	2,211
LA3	80,416
MA2	1,584

NA2	40,241
OA3	187,771
PA2	355,538
QA1	610,566
RA3	564,252
SA3	58,524
TA2	82,888
UA4	636,101
VA2	824,099
WA1	96,023
XA4	808,815
YA2	94,806
ZA2	46,731
AB1	28,499
BB1	3,586
CB4	39,802
DB4	49,618
EB4	63,971
Density 1 =	831,519
Density 2 =	1,673,203
Density 3 =	1,490,233
Density 4 =	<u>2,189,821</u>
Total =	6,184,776

Poquoson East, Va.
VIMS Map # 141

AA1	56,607
BA2	623,254
CA4	5,572,292
DA3	787,979
EA4	573,908
FA4	62,226
GA4	19,119
HA4	9,655
IA3	1,894
JA4	8,481
KA1	946
LA1	1,723,007
MA3	2,377,833

Density 1 = 1,780,559
 Density 2 = 623,254
 Density 3 = 3,167,707
 Density 4 = 6,245,680

Total = 11,817,200

Elliotts Creek, Va.
 VIMS Map # 142

AA2 1,136,293

Density 1 = 0
 Density 2 = 1,136,293
 Density 3 = 0
 Density 4 = 0

Total = 1,136,293

Hampton, Va.
 VIMS Map # 147

AA4 40,070
 BA1 45,348
 CA3 174,783
 DA4 187,160
 EA3 296,875
 FA3 147,352
 GA4 108,539
 HA1 330,342
 IA3 102,294
 JA3 108,097
 KA4 412,110
 LA3 48,162
 MA3 59,288
 NA3 21,664
 OA1 15,110
 PA2 647,765
 QA4 531,122
 RA4 327,990
 SA1 65,858

Density 1 = 456,658

Density 2 = 647,765
 Density 3 = 958,515
 Density 4 = 1,606,992

Total = 3,669,930

Cape Henry, Va.
 VIMS Map # 152

AA2 51,606
 BA3 60,157
 CA3 9,253
 DA2 22,308
 EA1 11,766
 FA1 4,846
 GA1 52,262

Density 1 = 68,874
 Density 2 = 73,914
 Density 3 = 69,410
 Density 4 = 0

Total = 212,198

Port Tobacco, Md.
 VIMS Map # 161

AA4 122,124
 Density 1 = 0
 Density 2 = 0
 Density 3 = 0
 Density 4 = 122,124

Total = 122,124

Charlotte Hall, Md.
 VIMS Map # 162

AA4 46,044
 Density 1 = 0
 Density 2 = 0

Density 3 =	0
Density 4 =	<u>46,044</u>
Total =	46,044

Assawoman Bay, Md.
VIMS Map # 166

AA1	14,022
BA1	22,679
CA1	139,157
DA1	27,691

Density 1 =	203,549
Density 2 =	0
Density 3 =	0
Density 4 =	<u>0</u>

Total =	203,549
---------	---------

Berlin, Md.
VIMS Map # 167

AA2	7,210
BA4	28,384
CA3	16,518
DA3	42,828
EA2	6,453
FA3	14,353
GA3	10,382
HA1	27,492

Density 1 =	27,492
Density 2 =	13,663
Density 3 =	84,082
Density 4 =	<u>28,384</u>

Total =	153,620
---------	---------

Ocean City, Md.
VIMS Map # 168

AA4	9,029
-----	-------

BA3	3,687
CA3	250,277
DA2	5,509
EA2	26,463
FA2	7,752
GA2	2,600
HA2	25,293

Density 1 =	0
Density 2 =	67,617
Density 3 =	253,964
Density 4 =	<u>9,029</u>

Total =	330,610
---------	---------

Tingles Island, Md.
VIMS Map # 170

AA2	26,122
CA3	1,156,132
DA4	5,702,765
EA4	15,075
FA2	21,582
GA1	207,346
IA2	16,872
JA2	12,735
KA2	8,458
LA3	2,380,898
MA2	495,212
NA2	4,320
OA2	180,920
PA3	720,587
QA2	424,667
RA4	185,708
SA2	81,865
TA2	225,082
UA2	33,169

Density 1 =	207,346
Density 2 =	1,531,005
Density 3 =	4,257,617
Density 4 =	<u>5,903,548</u>

Total = 11,899,515

Boxiron, Md.-Va.
VIMS Map # 172

AA2	315,649
BA4	2,660,002
CA2	342,305
DA2	53,203
EA2	2,001,766
FA3	104,010
GA2	1,293,423
HA4	28,978
IA2	126,642
JA2	11,897
KA1	93,073
LA4	409,015
MA2	170,053
NA2	564,117

Density 1 =	93,073
Density 2 =	4,879,055
Density 3 =	104,010
Density 4 =	<u>3,097,994</u>

Total = 8,174,133

Whittington Point, Md.-Va.
VIMS Map # 173

AA1	16,914
BA1	297,108
CA4	474,162
DA2	906,468
EA3	497,649
FA2	1,452,866
GA3	810,230
HA3	39,417
IA2	3,261
JA1	18,410

Density 1 =	332,432
Density 2 =	2,362,595

Density 3 =	1,347,296
Density 4 =	<u>474,162</u>

Total = 4,516,485

Chincoteague West, Va.
VIMS Map # 174

AA4	139,216
Density 1 =	0
Density 2 =	0
Density 3 =	0
Density 4 =	<u>139,216</u>

Total = 139,216

Chincoteague East, Va.
VIMS Map # 175

AA4	9,713,411
BA2	498,189
CA1	72,306
DA2	21,987
EA2	2,184
FA2	39,021

Density 1 =	72,306
Density 2 =	561,381
Density 3 =	0
Density 4 =	<u>9,713,411</u>

Total = 10,347,098

East of New Point Comfort, Va.
VIMS Map # 177

AA1	172,890
BA4	12,657

Density 1 =	172,890
Density 2 =	0
Density 3 =	0

Density 4 = 12,657

Total = 185,547

Bethel Beach, Va.
VIMS Map # 178

AA3 3,580

BA3 6,281

CA1 47,676

Density 1 = 47,676

Density 2 = 0

Density 3 = 9,860

Density 4 = 0

Total = 57,536

Goose Island, Va.
VIMS Map # 179

AA2 22,041

BA2 56,920

CA3 303,577

DA2 888,525

EA1 206,368

FA3 296,395

Density 1 = 206,368

Density 2 = 967,485

Density 3 = 599,972

Density 4 = 0

Total = 1,773,826

APPENDIX E

1993 Submerged Aquatic Vegetation Ground Survey Data Listed by USGS 7.5 Minute Quadrangle
and by 1993 Bed.

KEY

* Abbreviations under column "Species" are as follows:

Zm	-	<i>Zostera marina</i> (eelgrass)
Rm	-	<i>Ruppia maritima</i> (widgeon grass)
C	-	<i>Chara</i> sp. (muskgrass)
Cd	-	<i>Ceratophyllum demersum</i> (coontail)
Ec	-	<i>Elodea canadensis</i> (common elodea)
Hd	-	<i>Heteranthera dubia</i> (water stargrass)
Hv	-	<i>Hydrilla verticillata</i> (hydrilla)
Ms	-	<i>Myriophyllum spicatum</i> (Eurasian watermilfoil)
N	-	<i>Najas</i> spp. (naiad)
Nfl	-	<i>Najas flexilis</i> (northern naiad)
Ngr	-	<i>Najas gracillima</i> (slender naiad)
Ngu	-	<i>Najas guadalupensis</i> (southern naiad)
Nm	-	<i>Najas minor</i> (no common name)
Pcr	-	<i>Potamogeton crispus</i> (curly pondweed)
Pe	-	<i>Potamogeton epihydrus</i> (leafy pondweed)
Ppc	-	<i>Potamogeton pectinatus</i> (sago pondweed)
Ppf	-	<i>Potamogeton perfoliatus</i> (redhead-grass)
Ppu	-	<i>Potamogeton pusillus</i> (slender pondweed)
Tn	-	<i>Trapa natans</i> (water chestnut)
Va	-	<i>Vallisneria americana</i> (wild celery)
Zp	-	<i>Zannichellia palustris</i> (horned pondweed)
U	-	Unknown species composition

** Abbreviations under column "Surveyor" are as follows:

Cit.	-	Citizens' Survey
Harford	-	Harford Community College
PARK	-	Patuxent River Park Staff
Pines	-	Ocean Pines Boat Club
USFWS	-	United States Fish and Wildlife Service
USGS	-	United States Geological Survey
VIMS	-	Virginia Institute of Marine Science

\ - Slash mark separates species data of independent survey sources and independent survey dates.

- No SAV bed mapped from 1992 or 1993 aerial photography but SAV bed presence was verified by 1993 groundtruth survey at this location.

* - No SAV bed mapped from 1993 aerial photography but SAV bed presence was verified by 1993 groundtruth survey at this location.

APPENDIX E

1993 Submerged Aquatic Vegetation Ground Survey Data Listed by USGS 7.5 Minute Quadrangle

Quad	1992 Bed	1993 Bed	Species**	Surveyor***	1993 Survey Date
002	FA3	*	Ms,U	Cit.	09-24
	EA3	GA4	Hd,Hv,Ms,Va,U\ Ms,Hv,Va,Cd,Hd	Cit.\Harford	09-24\09-11
	DA3	HA4	Ms	Cit.	09-24
	AA4	AA4	Hd,Hv,Ms,Va	Cit.	09-24
	GA3	IA4	Ms	Harford	09-11
	HA3	EA3	Ms,Hv,Va,Cd	Harford	09-11
	HA1	EA1	Hv,Ms\ Ms	Cit.\Harford	06-15\09-11
	AA2	AA2	Ms\ Ms,Va	Cit.\Harford	06-15\09-11
	QA4	OA4	Ms,U\ Ms,Hv,Cd	Cit.\Harford	09-24\09-11
	SA3	PA4	Ms	Cit.	09-24
003	TA4	QA4	Ms	Cit.	09-24
	VA4	RA4	Ms\ Ms,Hv,Hd,Cd	Cit.\Harford	09-24\09-11
	UA4	SA4	Ms	Cit.\Harford	09-24\09-11
	XA4	TA3	Ms\ Ms,Hv,Hd,Cd	Cit.\Harford	09-24\09-11
	YA4	UA3	Hd,Hv,Ms\ Ms,Hv	Cit.\Harford	09-24\09-11
	AB4	YA4	Hd,Ms	Cit.	09-24
	BB4	ZA3	Hd,U\ Ms,Hd,Hv,Cd	Cit.\Harford	09-24\09-11
	DB4	DB4	Hv,Ms,U\ Ms,Hv,Va,Cd	Cit.\Harford	09-24\09-11
	EB4	EB4	Hv,Ms,U\ Ms,Hv,Hd,Cd,Va	Cit.\Harford	09-24\09-11
	FB4	FB4	Hv,Ms,U\ Ms,Hv,Va	Cit.\Harford	09-24\09-11
	XB3	AB4	Ms,Hd\ Ms,Hv	Cit.\Harford	09-24\09-11
	WB2	BB4	Ms,U\ Ms,Hv,Hd,Cd	Cit.\Harford	09-24\09-11
	VB4	WA4 (east)	Ms,U\ Ms,Hd,Cd	Cit.\Harford	09-24\09-11
	VB4	WA4 (west)	Ms,U\ Ms,Hd,Cd	Cit.\Harford	09-24\09-11
	LA2	JA4	Ms\ Ms,Cd	Cit.\Harford	09-24\09-11
	KA2	IA4	Ms\ Ms,Hv	Cit.\Harford	09-24\09-11
	OA4	MA4 (Bainbridge)	Ms,Hv,U\ Ms,Hv,Cd	Cit.\Harford	09-24\09-11
	OA4	MA4 (Bainbridge)	Ms,Hv,Cd	Harford	09-11
	OA4	MA4 (Happy Valley)	Ms,Cd	Harford	09-11
	CA4	BA2	Ms	Harford	09-11

Quad	1992 Bed	1993 Bed	Species**	Surveyor***	1993 Survey Date
003	EA1	CA2	Ms,Hv,Cd	Harford	09-11
	GA4, FA1	DA2 (Stump Pt.)	Va,Ms,Hd,Hv	Harford	09-11
	GA4, FA1	DA2 (Perry)	Ms,Hd	Harford	09-11
	GA4	FA4 (Perry Pt.)	Va,Ms,Hv,Cd	Harford	09-11
	GA4	FA4 (Perry Pt. Med. Ctr.)	Hv,Ms	Harford	09-11
	GA4	FA4 (Perry Pt. Park)	Hv, Va,Ms	Harford	09-11
	GA4	FA4 (Perry)	Va,Hv,Ms	Harford	09-11
	JA2	HA4	Ms\ Ms,Hv	Harford	09-11
	MA4	KA4 (Frenchtown)	Hv,Ms,U Ms,Hv,Cd	Cit.\Harford	09-24\09-11
	MA4	KA4 (Toll Rd.)	Ms,Cd	Cit.\Harford	09-24\09-11
	NA4	LA4	Ms,Cd	Harford	09-11
	ZA3	VA3	Ms,Hv	Harford	09-11
	CB4	CB4	Ms,Hv, Va,Hd,Cd	Harford	09-11
	GB4	GB4	Hv,Ms,Hd	Harford	09-11
	HB4	HB4	Ms, Va,Hv,Hd	Harford	09-11
	IB4	IB4	Ms, Va,Hv,Hd,Cd	Harford	09-11
	JB4	JB4	Ms, Va,Hv,Hd	Harford	09-11
	LB1	LB1	Ms, Va	Harford	09-11
	NB4	MB4	Ms,Hv, Va,Hd,Cd,Ngu	Harford	09-11
	OB4	NB4 (Todd Pk.)	Ms,Hv, Va,Hd,Cd	Harford	09-11
	OB4	NB4 (Nat. Guard Reserv.)	Ms,Hv,Hd, Va	Harford	09-11
	OB4	NB4 (south)	Ms,Hv,Hd	Harford	09-11
	QB4	OB3 (south)	Ms,Hv,Hd	Harford	09-11
	QB4	OB3 (north)	Ms,Hv,Hd	Harford	09-11
	RB4	PB3 (south)	Ms, Va,Cd,Hv	Harford	09-11
	RB4	PB3 (south-middle)	Ms, Va,Cd,Hv	Harford	09-11
	RB4	PB3 (middle)	Ms,Hv, Va	Harford	09-11
RB4	PB3 (north-middle)	Ms,Hv, Va	Harford	09-11	
RB4	PB3 (north)	Ms,Hv	Harford	09-11	
SB2	QB2	Ms,Cd,Hd	Harford	09-11	
TB2	RB2	Ms,Hd	Harford	09-11	
004	BA1	DA1	Ms,U Ms	Cit.\Harford	08-07\09-11
	AA1	CA1	Ms	Cit.\Harford	08-07\09-11
	CA1	AA2	Ms	Cit.\Harford	08-07\09-11
	DA1	*	Ms	Cit.	08-07

1993 Survey Date

Surveyor***

Species**

1993 Bed

1992 Bed

Quad

Quad	1992 Bed	1993 Bed	Species**	Surveyor***	1993 Survey Date	
004	FA1	Piney Cr.#	Por, Va	Cit.	08-97	
	EA1	FA1	Ms	Harford	09-11	
		EA1	Ms	Harford	09-11	
		BA1	Ms	Harford	09-11	
009	BB1	KB4	Ms,Hv	Harford	09-29	
	BB1	JB2	Va,Hd,Ms,Ngu	Harford	09-29	
	AB2	LB2	Ms	Harford	09-29	
		VA2	Ms,Hv	Harford	09-29	
	PA2	UA2	Ms,Hv	Harford	09-29	
	OA2	TA3	Ms	Harford	09-29	
	KA2	RA2	Ms	Harford	09-29	
	IA2	RA2	Ms,Hv	Harford	09-29	
	RA3	YA3	Ms	Harford	09-29	
	UA3	AB2	Ms	Harford	09-29	
		GAI	Ms	Harford	09-29	
		BA2	Ms	Harford	09-29	
		AA1	Ms	Harford	10-03	
				Harford	10-03	
	010	IA1	HA1	Ms	Harford	09-19
		IA1	IA1	Ms	Harford	09-19
HA2		JA2	Va,Ms	Harford	09-19	
GA2, FA1		KA2	Ms, Va,Ppc,Hd	Harford	09-19	
JA2		GAI	Ms	Harford	09-19	
BA1		DA2	Ms, Va	Harford	09-19	
FA1		LA2	Va,Ms	Harford	09-19	
FA1		MA3	Ms, Va,Hd	Harford	09-19	
FA1		NA1	Va	Harford	09-19	
FA1		NA1	Ms	Harford	09-19	
FA1		OA1	Ms	Harford	09-19	
EA3		PA4	Va,Ms	Harford	09-19	
DA2		FA3	Ms, Va	Harford	09-19	
CA2		FA3	Ms, Va,Hd	Harford	09-19	
		EA1	Ms	Harford	09-19	
AA2		CA3	Ms, Va	Harford	09-19	

Quad	1992 Bed	1993 Bed	Species**	Surveyor***	1993 Survey Date	
013	CA2	CA2	Cd,Ec,Ms,Pcr,Va,Zp	Cit.	06-01	
		Galloway Cr.#	Cd,Ec,Ms,Pcr	Cit.	04-08	
		Baltimore Yacht Club#	Cd,Ec,Ms,Pcr	Cit.	04-08	
014	IA2 HA2 GA3 EA4 YA3 TA2	HA2	Va,Ms,Ec,Cd	Harford	10-06	
		GA1	Ms,Ec	Harford	10-06	
		FA2	Va,Ms,Ec	Harford	10-06	
		DA2	Va,Ms,Ec	Harford	10-06	
		IA3	Ms,Ec,Va	Harford	10-06	
		JA2	Ms	Harford	10-06	
016	CA2 BA3	GA1	Ms	Harford	10-03	
		FA1	Ms	Harford	10-03	
		EA1	Ms	Harford	10-03	
		DA2	Ms	Harford	10-03	
		CA1	Ms,Va	Harford	10-03	
		BA3	Ms,Va	Harford	10-03	
		AA3	Ms,Va	Harford	10-03	
017	BA4 CA1	BA4	Ms	Harford	10-03	
		CA1	Ms	Harford	10-03	
023	Cockley Cr.# Cattail Cr.# Mill Cr.#	Cockley Cr.#	Zp	Cit.	06-15	
		Cattail Cr.#	Zp	Cit.	06-24	
		Mill Cr.#	Zp	Cit.	06-15	
024	Little Magothy R.# Little Magothy R.# Little Magothy R.# Deep Cr.# Deep Cr.# AA2 Shore Acres# BA2 Ulmsteads Pt.#	Little Magothy R.#	Zp	Cit.	06-24	
		Little Magothy R.#	Zp	Cit.	06-24	
		Little Magothy R.#	Zp	Cit.	06-24	
		Deep Cr.#	Zp	Cit.	06-24	
		Deep Cr.#	Zp	Cit.	06-24	
		AA2	Ppc,Zp\Ppc	USFWS	06-24\08-20	
		Shore Acres#	Cd,Ms,Zp	Cit.	06-24	
		BA2	Cd,Ms,Zp	Cit.	06-24	
		Ulmsteads Pt.#	Ms,Ppf,Rm	Cit.	06-24	

Quad	1992 Bed	1993 Bed	Species**	Surveyor***	1993 Survey Date	
024		Forked Cr. #	Zp	Cit.	06-24	
		Cool Spring Cove#	Zp	Cit.	06-24	
		Springs Pond#	Zp	Cit.	06-24	
		Blackhole Cr. #	Zp	Cit.	06-24	
		Blackhole Cr. #	Zp	Cit.	06-24	
		Blackhole Cr. #	Zp	Cit.	06-24	
		Chest Neck#	Zp	Cit.	06-24	
		Broad Cr. #	Zp	Cit.	06-24	
		Park Cr. #	Zp	Cit.	06-24	
		CA2	Ppc	USFWS	06-24\08-20	
		Grays Cr. #	Zp	Cit.	06-24	
		Grays Cr. #	Zp	Cit.	06-24	
		Grays Cr. #	Zp	USFWS	06-24\08-20	
		DA2	Ppc	USFWS	06-24\08-20	
		EA2	Rm	USFWS	08-24\08-20	
		Tar Cove#	Zp	Cit.	06-24	
		Cornfield Cr. #	Zp	Cit.	06-24	
		Cornfield Cr. #	Zp	Cit.	06-24	
		IA3	Ec,Rm	USFWS	06-24\08-20	
		Redhouse Cove#	Zp	Cit.	06-24	
		HA2	Ec,Rm	USFWS	06-24\08-20	
		GA2	Zp\Nfl	Cit.\USFWS	06-24\08-20	
		Magothy Narrows#	N	USFWS	06-24\08-20	
		FA2	Ppf,Rm	USFWS	06-24\08-20	
		Bayberry#	Ec	Cit.	06-24	
	026		Durdin Cr. #	Rm	USFWS	08-18
			PA2	Ppf,Rm	USFWS	08-18
			OA3	Ppf	USFWS	08-20
			NA2	Ms	USFWS	08-20
			MA2 (north)	Ms,Ppf,Rm	USFWS	08-20
			MA2 (south)	Cd,Ec,N,Ppf,Rm	USFWS	08-20
			LA3 (south)	Rm	USFWS	08-20
			LA3 (Little Gum Pt.)	Rm	Cit.\USFWS	No Date\08-20
		LA3 (Spring Pt.)	Rm	USFWS	08-20	
		KA2	Rm,Nfl	USFWS	08-20	

Quad	1992 Bed	1993 Bed	Species**	Surveyor***	1993 Survey Date	
026	NA2	HA3	Cd,N,I,Rm\ Cd,Nfl,Rm	Cit.\USFWS	No Date\08-20	
	PA2	GA3	Rm	Cit.\USFWS	No Date\08-20	
028		DA3	Hv	USGS	07-20-22	
		CA3	Ms,Hv,U	USGS	07-20-22	
030		Muddy Cr. #	Rm	Cit.	08-09	
		Selby Bay#	Rm	Cit.	08-15	
		Selby Bay#	Zp	Cit.	05-17	
		Selby Bay#	Rm	Cit.	08-15	
		Limehouse Cove#	Rm	Cit.	08-15	
		Cedar Pt. #	Rm	Cit.	08-15	
		Glebe Bay#	Rm	Cit.	08-15	
		Glebe Bay#	Rm	Cit.	08-15	
		Beards Cr. #	Rm	Cit.	08-09	
		Beards Cr. #	Rm	Cit.	08-09	
		Crab Cr. #	Zp	Cit.	05-29	
		Spa Cr. #	Zp	Cit.	06-10	
		Hillsmere Shores #	Rm	Cit.	08-15	
	031		Flagg Pond#	Rm	Cit.	08-15
			Duvall Cr. #	Rm	Cit.	08-15
		Cherrytree Cove#	Rm	Cit.	08-15	
		Oyster Cr. #	Rm	Cit.	08-15	
		Lk. Ogleton#	Ppu,Zp	Cit.	06-01	
		Heron Lk. #	Rm	Cit.	08-15	
		Back Cr. #	Zp	Cit.	04-15	
		Spa Cr. #	Zp	Cit.	06-10	
		Meadow Pt. #	Rm	Cit.	08-15	
		Severn Beach#	Rm	Cit.	08-15	
		Hackett Pt. #	Rm	Cit.	08-15	
032			QA3 (north)	Rm,Zp	VIMS	07-20
		AA3	QA3 (south)	Rm,Zp	VIMS	07-20
		HA1	HA2	Rm	VIMS	07-20
			GAI	Rm	VIMS	07-20

Quad	1992 Bed	1993 Bed	Species**	Surveyor***	1993 Survey Date	
033		Kent Is. #	Zp	Cit.	07-15	
		Kent Narrows#	Ms, Ppf, Zp	Cit.	07-15	
		NA3	Ppf, Zp	Cit.	07-15	
		PA3	Ppf, Zp	Cit.	07-15	
		Wye R. #	Rm	Cit.	08-01	
		Wye R. #	U	Cit.	08-01	
		Wye R. #	U	Cit.	08-01	
		Hood Pt. #	Rm	Cit.	06-27	
		MA3	Ppf, Rm, Zp	Cit.	07-15	
	HA3	LA3	Ms, Ppf, Zp	Cit.	07-15	
	GA2	IA3	Rm	VIMS	07-20	
	034		YA4	Va, Hv, Ms, Hd	USGS	07-20-22
			XA2	Va, Hv, Ms, Hd	USGS	07-20-22
			WA4	Va, Hv, Ms, Hd	USGS	07-20-22
		RA2, QA1	Hd, Hv, Va	USGS	07-20-22	
		PA4	Va, Ms, Hd	USGS	07-20-22	
		MA3	Cd, Ms, Ppc, Va, Hv, Pcr, Nm, Hd	USGS	07-20-22	
		LA2	Va, Hv, Ms	USGS	07-20-22	
		OB4 (Goose Is.)	Ms, Hv, Cd	USGS	07-20-22	
		OB4 (Fox Ferry Pt.)	Ms, Hv, Cd	USGS	07-20-22	
		OB4 (south)	Hv, Ms	USGS	07-20-22	
		PB4	Hv, Ms	USGS	07-20-22	
		UB2 (north)	Va, Ms, Hv	USGS	07-20-22	
		UB2 (south)	Hv, Va, Cd, Hd, Ms	USGS	07-20-22	
		Indian Queen Bluff#	Hv, Va, Cd, Hd, Ms	USGS	07-20-22	
036			WA3	Zp	Cit.	06-13
			Northeast Branch#	Zp	Cit.	07-04
			SA2	Rm, Zp	Cit.	08-15
			Broad Cr. #	Ppf	Cit.	06-27
			Broad Cr. #	Rm	Cit.	06-27
			*	Rm	Cit.	06-27
		DB1 (south)	Rm, Zp	VIMS	07-20	
		FB2 (north)	Rm, Zp	VIMS	07-20	
	DA2	FB2 (south)	Rm, Zp	VIMS	07-20	

Quad	1992 Bed	1993 Bed	Species**	Surveyor***	1993 Survey Date	
037		RA4	Ppf	Cit.	06-27	
	BA2	MA3 (north)	Rm	VIMS	07-20	
	BA2	MA3 (south)	Rm	VIMS	07-20	
039		IA4 (north)	Va,Hv,Ms	USGS	07-20-22	
		IA4 (south)	Cd,Ms,Hv,Va,Nm	USGS	07-20-22	
		Whitestone Pt.#	Cd, Ms,Hv, Va.,Nm	USGS	07-20-22	
		Gunston Hall#	Va,Ms	USGS	07-20-22	
		FA2 (north)	Hv, Va,Ms	USGS	07-20-22	
		FA2 (south)	Va,Ms	USGS	07-20-22	
		EA1	Va,Ms,Hv	USGS	07-20-22	
		DA4	Ms,Hv, Va,Nm	USGS	07-20-22	
	040		Broad Cr.#	Hv	USGS	07-20-22
			IA4	Va,Hv	USGS	07-20-22
		GA4	Hv,Hd	USGS	07-20-22	
		Mockley Pt.#	Hv,Hd	USGS	07-20-22	
		CA4	Hv,Hd	USGS	07-20-22	
		BA4 (Nat. Colonial Farm)	Cd,Hv,Ms	USGS	07-20-22	
		BA4	Ms,Hv, Va,Cd	USGS	07-20-22	
		BA4 (River Rd.)	Ms,Hv, Va,Cd	USGS	07-20-22	
BA4		BA4 (Fenwick)	Hv,Ms, Va\ Ms,Hv	Cit.USGS	06-26\07-20-22	
		Ferry Pt.#	Hv	USGS	07-20-22	
		Sheridan Pt.#	Hv	USGS	07-20-22	
		Potomac R.#	Ppc,Hv	USGS	07-20-22	
		NA2	Hv,Nm, Va,Ms	USGS	07-20-22	
		OAI	Hv,Nm, Va,Ms	USGS	07-20-22	
041			Mataponi Cr.#	Nm,Per	PARK	07-14
			Mataponi Cr.#	Nm,Per,Ec	PARK	08-05
			Lyons Cr.#	Ec	PARK	08-05
		Cocktown Cr.#	Cd,Ec,Per,Ngu, Va	PARK	07-30	
		Swamp Cr.#	Cd,Ec,Per	PARK	07-30	
		Balls Cr.#	Rm	Cit.	07-01	
043	AA3	CA2	Rm,Zp	VIMS	07-21	

Quad	1992 Bed	1993 Bed	Species**	Surveyor***	1993 Survey Date	
043	AA3	BA2 (west)	Rm	VIMS	07-21	
	AA3	BA2 (east)	Rm	VIMS	07-21	
	AA3	AA2	Rm	VIMS	07-21	
		HA2	Rm	VIMS	07-21	
		IA3	Rm	VIMS	07-21	
	JA2	KA3 (south)	Rm	VIMS	07-21	
	JA2	KA3 (north)	Rm	VIMS	07-21	
	NA2	QA2	Rm	VIMS	07-21	
	044		Tar Cr.#	Zp	Cit.	06-16
			Plaindealing Cr.#	Zp	Cit.	06-16
		Tred Avon R.#	Zp	Cit.	06-16	
		Trippe Cr.#	Zp	Cit.	06-16	
		Bailey's Neck#	Zp	Cit.	07-16	
		Goldsborough Cr.#	Rm, U	Cit.	07-16	
		Boone Cr.#	U	Cit.	07-15	
RA2		BA3	Rm	Cit.	07-16	
QA2		CA3	Rm	Cit.	07-16	
		PA2	Rm	Cit.	07-16	
		Westpoint Rd. Cove#	U	Cit.	06-15	
045			CA4	Hv, Ms, Va, Hd, Cd, Nm, Ngu	USGS	07-20-22
			AA4	Hv, Va, Ms, Nm, Cd, Hd	USGS	07-20-22
		EA2	Hv, Va, Ms, Cd, Nm	USGS	07-20-22	
		FA4	Hv, Nm, Va	USGS	07-20-22	
		IA4	Hv, Ms, Va, Nm	USGS	07-20-22	
		KA4	Hv, Va, Ms, Cd, Hd, Nm	USGS	07-20-22	
		MA4	Hv, Va, Ms, Cd	USGS	07-20-22	
		MA4	Hv, Ms	USGS	07-20-22	
		NA4	Hv, Ms	USGS	07-20-22	
		PA4	Hv, Va, Ms, Hd	USGS	07-20-22	
		QA2	Hv, Va, Ms	USGS	07-20-22	
	048	Chapman Pt.	JA4	Hd, Ms, Cd	USGS	07-20-22
		Glymont	JA4	Ms, Hv, Cd, Va	USGS	07-20-22

Quad	1992 Bed	1993 Bed	Species**	Surveyor***	1993 Survey Date	
048		IA4	Hv,Ms,Va,Hd,Cd	USGS	07-20-22	
		HA4	Hv,Ms,Va,Hd,Cd	USGS	07-20-22	
	BA4	BA4	Nm,Hv,Va,Cd,Ms,Hd	USGS	07-20-22	
	BA4	CA2	Hv,Ms,Cd	USGS	07-20-22	
	BA4	CA2	Hv,Ms,Va	USGS	07-20-22	
049		Patuxent R. #	Ec,Ngu,Pcr,Ppu,Zp	Cit.	06-13	
		Patuxent R. #	Pcr,Ppu,Zp	Cit.	06-13	
051	TA3	LA3, MA1	Rm	Cit.	06-01	
	WA3	OA3	Rm	VIMS	07-21	
	XA2	OA3	Rm	VIMS	07-21	
	XA2, WA3	OA3	Rm	VIMS	07-21	
	TA3	MA1	Rm	VIMS	07-21	
	QA2	KA1	Rm	Cit.\VIMS	06-01\07-21	
	SA1	KA1	Rm	Cit.\VIMS	06-01\07-21	
	PA2	JA2	Rm	Cit.	06-01	
	OA2	*	Rm	Cit.	06-01	
	UA1	MA1	Rm	VIMS	07-21	
		DA2	Zp	Cit.	06-15	
	052		IA3	Zp	Cit.	06-15
		HA3	*	Zp	Cit.	06-15
FA2		HA2	Zp	Cit.	06-15	
GA3		*	Zp	Cit.	06-15	
053		Chancellor Pt. (west)#	Zp	Cit.	Spring-Summer	
		Chancellor Pt. (middle)#	Zp	Cit.	Spring-Summer	
		Chancellor Pt. (east)#	Zp	Cit.	Spring-Summer	
055		RA4	Va,Ms	USGS	07-20-22	
	Clifton Beach	PA4	Va,Ms,Hd,Cd	USGS	07-20-22	
	Wades Bay	PA4	Va,Ms,Hd,Nm,Cd,Hv	USGS	07-20-22	
	Douglas Pt.	PA4	Va,Ms,Hd	USGS	07-20-22	
		PA4	Ms,Va,Hv,Hd	USGS	07-20-22	
		NA4	Hv,Va,Ms,Hd,Cd	USGS	07-20-22	

Quad	1992 Bed	1993 Bed	Species**	Surveyor***	1993 Survey Date
055		LA4	Va,Ms,Hv,I,Id	USGS	07-20-22
		JA4	Ms,I,Hd,Va,Hv,Nm,Ng	USGS	07-20-22
	Clifton Pt.	JA4	Hv,Hd,Ms,Va,Cd,Nm,Ng	USGS	07-20-22
060		Peterson's Pt. #	Zp	Cit.	05-15
		Wells Cove #	Rm,Zp	Cit.	05-15
		Island Creek#	Zp	Cit.	06-15
061		St. Leonard Cr. #	Zp	Cit.	06-15
		Cape Leonard (west)#	Zp	Cit.	06-15
		Cape Leonard (east)#	Zp	Cit.	06-15
		Osbourne Cove (west)#	Zp	Cit.	06-15
		Osbourne Cove (east)#	Zp	Cit.	06-15
		Breeders Rd. #	Zp	Cit.	06-15
		Osbourne Cove#	Zp	Cit.	06-15
		Neale Sound. #	Zp	Cit.	05-18
070		Cuckold Cr. #	Zp	Cit.	06-06
		Green Holly Pond#	Zp	Cit.	06-12
071		Hominy Cr. #	Zp	Cit.	04-15
		CA2	Rm	Cit.	06-15
072		BA2	Rm	VIMS	07-22
		Great Cove#	Rm	VIMS	07-22
		Cove Pt. #	Rm	VIMS	07-22
		IB3	Rm	VIMS	07-22
073		NB3	Rm	VIMS	07-22
		NB3	Rm	VIMS	07-22
		HA4	Rm	VIMS	07-22
		FB3	Rm	VIMS	07-22
		FB3	Rm	VIMS	07-22
		FB3	Rm	VIMS	07-22
		QA3	Rm	VIMS	07-22
		TA2	Rm	VIMS	07-22
	NA2 (north)	Rm	VIMS	07-22	

Quad	1992 Bed	1993 Bed	Species**	Surveyor***	1993 Survey Date
073	TA2	NA2 (south)	Rm	VIMS	07-22
		UA2	Rm	VIMS	07-22
	AB3	UA2 (north)	Rm	VIMS	07-22
	AB3	UA2 (south)	Rm	VIMS	07-22
074	EA4	FA4	Rm	Cit.	05-15
076		Popes Cr. #	Ms	Cit.	10-09
078		Weatherall Cr. (east)#	Zp	Cit.	06-06
		Weatherall Cr. (west)#	Zp	Cit.	06-06
		Lvr. Machodoc Cr.#	Per	Cit.	06-12
		E. of Herring Pond#	Zp	Cit.	06-12
080	AA2	*	Rm	Cit.	08-31
	BA2	BA2	Rm	Cit.	08-31
084	AA2	BA3	Rm	Cit.	10-13
	BA2	CA2	Rm	Cit.	10-13
	CA2	DA2	Rm	Cit.	10-13
	DA2	GA2	Rm	Cit.	10-13
	EA2	GA2	Rm	Cit.	10-13
		Laws Thorofare#	Rm	Cit.	10-13
099	AA2	SA2	Rm	Cit.	05-22
	BA2	XAI, RA2	Rm,Zm	Cit.	05-22
	CA3	ZA4, SA2	Rm	Cit.	05-22
	RA3	WA2, UA3	Rm	Cit.	05-22
		Cheeseman Is.#	Rm	Cit.	05-22
		Hog Neck#	Rm	Cit.	05-22
100	NA2	JA1	Rm,Zm	VIMS	07-23
	LA2	HA1	Rm,Zm	VIMS	07-23
	WA2	QA4	Rm,Zm	VIMS	07-23
	WA2	RA2	Rm,Zm	VIMS	07-23
	WA2	RA2	Zm	VIMS	07-23

Quad	1992 Bed	1993 Bed	Species**	Surveyor***	1993 Survey Date
100	XA4	RA2	Rm,Zm	VIMS	07-23
	XA4	RA2	Zm	VIMS	07-23
101	NA2	*	Rm,Zm	VIMS	07-23
	OA2	MA2	Rm,Zm	VIMS	07-23
	CA3	CA3 (north)	Rm,Zm	VIMS	07-23
	CA3	CA3 (middle)	Rm,Zm	VIMS	07-23
	CA3	CA3 (south)	Rm,Zm	VIMS	07-23
	FA2	EA2	Rm,Zm	VIMS	07-23
	GA3	FA3 (north)	Rm,Zm	VIMS	07-23
	GA3	FA3 (south)	Rm,Zm	VIMS	07-23
106	GA4	FA4	Rm,Zm	Cit.	07-06
	HA3	*	Zm	Cit.	07-06
	EA1	DA1	Zm	Cit.	07-06
	DA4	CA4	Rm,Zm	Cit.	07-16
	AA1	AA1	Zm	Cit.	08-07
	BA2	*	Zm	Cit.	08-07
	CA2	BA2	Zm	Cit.	08-07
		Sandy Pt.#	Rm	Cit.	06-15
		GA2	Rm	Cit.	06-15
		Bull Neck#	Rm	Cit.	06-15
	111	DA3	CA2	Rm,Zm	VIMS
GA4		EA2	Rm	VIMS	06-17
HA4		GA3	Rm	VIMS	06-17
IA1		HA2	Rm,Zm	VIMS	06-17
AB2		IA2	Rm	VIMS	06-17
YA3		ZA2	Rm	VIMS	06-17
LA4		JA2	Rm	VIMS	06-17
112	IB2	BB2	Rm,Zm	Cit.	07-01
	JB2	CB2	Rm,Zm	Cit.	07-01
	HB2	AB2	Zm	Cit.	07-01
	FB2	YA2	Zm	Cit.	07-01
	EB3	XA2	Zm	Cit.	07-01

Quad	1992 Bed	1993 Bed	Species**	Surveyor***	1993 Survey Date
112	PA3	*	Rm	Cit.	09-15
	MA4	LA2	Rm	Cit.	09-15
	LA2	KA2	Rm	Cit.	09-15
	QA3	MA2	Zm	Cit.	09-15
	TA3	PA3	Zm	Cit.	09-15
114	XA2	DB2	U	Cit.	09-09
	YA2	EB2	U	Cit.	09-09
118	GA2	IA2	Zm\Zm, Rm	Cit.\VIMS	08-07\06-16
	JA3	MA2	Zm	VIMS	06-16
	LA2	QA2	Zm	Cit.	08-07
	MA2	*	Zm	Cit.	08-07
	EA4	GA4	Zm\Zm, Rm	Cit.\VIMS	08-08\06-16
	FA2	HA2	Zm	Cit.	08-08
	CA4	FA2	Zm	Cit.	06-16
		PA2	Rm	VIMS	06-16
		OA2	Rm	VIMS	06-16
		CB1	Rm,Zm	VIMS	06-16
		DB4	Rm,Zm	VIMS	06-16
122	BA2	BA2	Rm,Zm	VIMS	06-05
123	NA2	LA2	Zm	VIMS	06-19
		MA2	Zm	VIMS	06-19
		BB3	Zm,Rm	VIMS	06-19
		AB1	Zm	VIMS	06-19
		ZA3	Zm	VIMS	06-19
		NA4	Zm	VIMS	06-19
		OA4	Zm,Rm	VIMS	06-19
124	CB2	DB2	Rm,Zm	Cit.	07-02
	BB2	CB2	Rm,Zm	Cit.	07-02
	DB4	AB4	Zm	Cit.	07-02
	EB2	BB2	Zm	Cit.	07-02
	GB3	FB3	Zm	Cit.	07-02

Quad	1992 Bed	1993 Bed	Species**	Surveyor***	1993 Survey Date
124	HB3	GB2	Rm,Zm	Cit.	07-02
	IB2	HB2	Rm,Zm	Cit.	07-02
	KB3	JB2	Rm,Zm	Cit.	07-02
131	DA4	DA4	Zm	VIMS	06-29
	HA4	GA4 (east)	Zm	VIMS	06-01
	HA4	GA4 (west)	Zm	VIMS	06-01
	OA4	OA4 (north)	Rm,Zm	VIMS	06-01
	OA4	OA4 (south)	Rm,Zm	VIMS	06-01
	QA4	QA4	Rm,Zm	VIMS	06-01
	NB2	JB4	Rm,Zm	VIMS	06-05
	FB2	BB2	Rm,Zm	VIMS	06-05
	JB4	FB4	Zm	VIMS	06-05
			Zm	VIMS	06-05
132	GA3	GA3	Rm,Zm	Cit.	05-29
	CA4	CA4	Rm,Zm	VIMS	06-01
	CA4	CA4	Zm	VIMS	06-01
	QA1	PA1	Zm	VIMS	06-20
	RA4	QA4	Zm	VIMS	06-20
	OA4	NA4	Zm	VIMS	06-20
	OA4	NA4	Zm	VIMS	06-20
			Zm	VIMS	06-20
			Zm	VIMS	JULY
			Zm	VIMS	06-30
139		CA1	Zm	VIMS	06-30
		BA2	Zm	VIMS	05-06
	AA2	AA1	Zm	VIMS	05-06
	BA2	FA1	Zm	VIMS	06-29
	CA2	EA2	Zm	VIMS	06-29
	DA3	DA2	Zm	VIMS	06-29
			Zm	VIMS	06-29
140	SA4	UA4	Rm,Zm	VIMS	06-20
	UA4	XA4	Rm,Zm	VIMS	06-30
	WA2	ZA2	Rm,Zm	VIMS	06-30
		AB1 (east)	Zm	VIMS	06-30
		AB1 (west)	Zm	VIMS	06-30

Quad	1992 Bed	1993 Bed	Species**	Surveyor***	1993 Survey Date	
140	XA4	BB1 CB4	Zm Zm	VIMS VIMS	06-29 06-29	
	CA2	BA3 DA2 BA3 AA2 The Narrows#	Zm Rm Rm Rm Rm	Cit. Cit. Cit. Cit. Cit.	06-07 07-02 07-02 07-02 07-02	
159		Sand & Gravel Pits# Sand & Gravel Pits# Spyglass Is. # Trailer Park# Back Channel# Mill Cr. - Rt. 4# Mill Cr. # Green Landing# Leon# Bristol Landing# Railroad Cr. # opp. Charles Branch# Charles Branch# Western Branch# Marlboro Speedway# House Creek#	Cd Cd Cd Cd,Pe,Ec,Per Ngu,Hv,Nm,Per,Va,Cd,Ec,Ppu Ec,Cd,Ngu,Nm,Per,Va,Hv Cd,Ec,Ngu,Nm Nm,Cd,Ngu Nm,Ec,Cd,Ngu,Per Nm,Cd,Ec,Per Nm,Cd,Ec Nm Per Cd,Nm Cd,Ec,Nm Nm,Per	PARK PARK PARK PARK PARK PARK PARK PARK PARK PARK PARK PARK PARK PARK PARK PARK	07-08 07-08 07-08 07-08 07-08 07-08 07-08 07-08 07-08 07-08 07-08 07-08 07-14 07-14 07-14 07-14	
	AA4		Hv,Ms,Cd	USGS	07-20-22	
	166	67th Street DA2 CA1	DA1	Rm,Zm Rm Rm,Zm	Cit. Cit. Cit.	07-27 07-27 07-27
		AA2 BA2	Horse Is. # *	Rm Rm	Cit. Cit.	07-27 08-02
		CA2	AA1 BA1 *	Rm Rm Rm	Cit. Cit. Cit.	08-02 08-02 08-02

Quad	1992 Bed	1993 Bed	Species**	Surveyor***	1993 Survey Date	
166		Ocean City Exprwy.#	Rm	Pines	07-27	
		Water tank#	Rm	Pines	07-27	
		S. of CA1 #	Rm	Pines	07-27	
		CA1	Rm	Pines	07-27	
		CA1	Rm	Pines	07-27	
		CA1	Rm	Pines	07-27	
		Devil Is.#	Rm	Pines	07-27	
		Devil Is.#	Rm	Pines	07-27	
		Devil Is.#	Rm	Pines	07-27	
167	CA4	GA3	Zm	Cit.	09-28	
	DA2	BA4	Zm	Cit.	10-01	
168	DA3	DA2	Zm	Cit.	09-28	
	AA3	AA4	Rm,Zm	Cit.	10-01	
		Water tank#	Rm	Pines	08-02	
		Filtration Plant#	Rm	Pines	08-02	
170		IA2	Zm	Cit.	06-11	
		Tidal Flat#	Rm,Zm	Cit.	09-28	
	OA3	UA2	Rm,Rm	Cit.	09-28	
	KA3	QA2, PA3	Rm,Zm	Cit.	09-28	
	MA3	TA2, SA2	Rm,Zm	Cit.	09-28	
	IA3	LA3	Rm,Zm	Cit.	10-01	
	JA4	MA2	Rm,Zm	Cit.	10-01	
	DA4	DA4	Rm,Zm	Cit.	10-04	
	CA3	CA3	Zm,Rm	Cit.	10-04	
	BA2	*	Zm,Rm	Cit.	10-04	
	FA2	DA4	Rm,Zm	Cit.	10-07	
	EA2	EA4	Rm	Cit.	10-07	
	172	LA3	MA2	Zm	Cit.	10-04
		KA4	LA4, NA2	Zm	Cit.	10-05
		JAI	KA1	Rm,Zm	Cit.	10-05
		IA2	IA2	Zm	Cit.	10-05
HA3		IA2	Zm	Cit.	10-05	

Quad	1992 Bed	1993 Bed	Species**	Surveyor***	1993 Survey Date
172	GA4	HA4	Zm	Cit.	10-05
	AA4	BA4, AA2	Rm, Zm	Cit.	10-06
	BA3	CA2	Zm	Cit.	10-06
	CA2	DA2	Zm	Cit.	10-06
	DA3	EA2	Rm, Zm	Cit.	10-06
	FA3	GA2	Rm, Zm	Cit.	10-06
173	JA2	IA2	Rm, Zm	Cit.	10-04
	HA3	GA3	Rm, Zm	Cit.	10-04
	GA3	FA2	Rm, Zm	Cit.	10-04
	EA4	CA4	Zm	Cit.	10-05
	AA1	AA1	Zm	Cit.	10-05
	DA2	DA2	Rm, Zm	Cit.	10-05
	FA2	EA3	Rm, Zm	Cit.	10-05
	CA1	BA1	Rm, Zm	Cit.	10-05
					10-07
175	EA3	FA2	Rm, Zm	Cit.	07-06
	AA4	AA4	Rm, Zm	Cit.	07-06





Virginia Institute of Marine Science
School of Marine Science
The College of William & Mary
Gloucester Point, Virginia 23062

