

D. Miller
Permanent Loan
LIBRARY
ENVIRON. PROT. AGENCY
MONTICELLO MARYLAND 21761

NRI
SPECIAL
REPORT
No. 4


OCTOBER 1971

**FISH EGGS
AND
LARVAE
OF THE
UPPER CHESAPEAKE BAY**



NATURAL
RESOURCES
INSTITUTE

UNIVERSITY OF MARYLAND



FOREWORD

The biological wealth of a highly productive estuary like Chesapeake Bay is readily apparent to those who enjoy its bounty of oysters, crabs, clams, and fish. About 500,000,000 pounds of these various seafoods are landed at bay ports each year. Thousands of people are directly or indirectly involved in the annual harvest—for many it is their livelihood, for many more it is wonderful outdoor fun.

Part of the wealth and greatness of the bay is, however, invisible to most of us. We never see or are usually unaware of the vast numbers of the immature stages of the species we know only as adults. This tremendous volume of life constitutes the source of larger forms. Each species plays an important role in the drama of life and each requires very special environmental circumstances for success.

William Dovel presents here a new and useful summary and analysis of the use which fish make of one of the world's richest estuarine nursery areas—the upper Chesapeake Bay and portions of its tributaries. He shows that nearly fifty species occur in this zone as eggs or larvae; that there are strong seasonal rhythms in growth and movement but that some young stages are present at almost all times; that users of the zone may originate in fresh water, the estuary, or the sea; and that suitable sampling and study of these valuable eggs and larvae are possible despite the large inherent variation in their abundance and the notable complexity of the estuarine environment.

Mr. Dovel emphasizes the importance of a characteristic of estuaries. The low-salinity portion, very frequently near densely populated areas, is a critical region in the life cycles of many species; but it is here also where many cities dump their wastes. Water quality and other environmental conditions in the nursery areas must be favorable — not merely tolerable — for the many small fish and associated organisms that develop there. If such quality is not protected, rich resources may be lost through effects most of us cannot even see.

Here, then, is valuable information garnered during an eleven year period—a catalogue of species, a summary of patterns, and identification of some of the essential areas necessary for the propagation and growth of extraordinarily important fish eggs and larvae in one of the world's greatest estuaries. Further research is required for optimal management, but Mr. Dovel has made a significant and valuable contribution.

L. EUGENE CRONIN
Director

October 1971

FISH EGGS AND LARVAE OF THE UPPER CHESAPEAKE BAY

NRI SPECIAL REPORT NO. 4

William L. Dovel

Contribution No. 460, Natural Resources Institute, University of Maryland

TABLE OF CONTENTS

| | | |
|----|---|----|
| 1. | INTRODUCTION | 1 |
| | Areas of Investigation | 1 |
| | Patuxent River | 2 |
| | Magothy River | 2 |
| | Upper Chesapeake | 2 |
| | Chesapeake and Delaware Canal | 2 |
| | Susquehanna River | 2 |
| 2. | METHODS | 2 |
| 3. | RESULTS | 3 |
| 4. | DISCUSSION | 4 |
| | Freshwater Spawners | 4 |
| | Family Clupeidae - herring; <i>Alosa</i> , <i>Brevoortia</i> , <i>Dorosoma</i> | 4 |
| | White perch, <i>Morone americana</i> | 5 |
| | Striped bass, <i>Morone saxatilis</i> | 5 |
| | Yellow perch, <i>Perca flavescens</i> | 6 |
| | Estuarine Spawners | 6 |
| | Bay anchovy, <i>Anchoa mitchilli</i> | 6 |
| | Northern pipefish, <i>Syngnathus fuscus</i> | 10 |
| | Naked goby, <i>Gobiosoma boscii</i> | 10 |
| | Rough silverside, <i>Membras martinica</i> | 10 |
| | Tidewater silverside, <i>Menidia beryllina</i> | 10 |
| | Atlantic silverside, <i>Menidia menidia</i> | 10 |
| | Winter flounder, <i>Pseudopleuronectes americanus</i> | 10 |

| | | |
|----|---|-------------------|
| | Hogchoker, <i>Trinectes maculatus</i> | 11 |
| | Skilletfish, <i>Gobiosox strumosus</i> | 11 |
| | Marine Species | 11 |
| | Atlantic menhaden, <i>Brevoortia tyrannus</i> | 11 |
| | Atlantic croaker, <i>Micropogon undulatus</i> | 12 |
| | Kingfish, <i>Menticirrhus americanus</i> | 12 |
| | Seaboard goby, <i>Gobiosoma ginsburgi</i> | 12 |
| 5. | SUMMARY | 12 |
| | Salinity Correlations | 13 |
| | Freshwater Spawners | 13 |
| | Estuarine Spawners | 13 |
| | Marine Spawners | 13 |
| | A Common Estuarine Nursery | 14 |
| | Movement | 14 |
| | Stratification | 15 |
| | Survival | 15 |
| | Rapid and Rhythmical | 15 |
| | Temperature Correlations | 16 |
| | Growth | 16 |
| 6. | LITERATURE CITED | 19 |
| 7. | APPENDIX — LIST OF TABLES | 21 |
| 8. | ACKNOWLEDGMENTS | Inside Back Cover |

INTRODUCTION

During the past eleven years, the author has been engaged in numerous investigations of the fish eggs and larvae prevalent in different areas of Chesapeake Bay. This report presents a synoptic characterization of the data collected during that period.

In the spring of 1960, the author, under the direction of the late Dr. Romeo J. Mansueti, began an intensive effort to gain a better knowledge of the biological characteristics of pelagic fish eggs and larvae that occur in channel areas of the lower Patuxent River (Fig. 1). No effort was made to investigate the eggs and larvae of shoreline areas. The area under investigation was later expanded to include additional areas of Chesapeake Bay (Table 1, Fig. 1). From 1960 to 1962, effort was devoted to designing appropriate field sampling techniques suitable for this estuarine system. In 1961, a one-year pilot survey was initiated over a 3-mile section at the mouth of the Patuxent River to determine the suitability of various types of gear and to gain familiarization with the ichthyoplankton found there. A weekly sampling program was initiated in January 1963 to determine seasonal biological characteristics of species present throughout the entire estuarine portion of the Patuxent River. This project continued for three years but with reduced effort during 1964 and 1965. From March 19, 1965 to May 24, 1966, a similar investigation was conducted in the Magothy River, Maryland.

Sampling was extended to the upper Chesapeake Bay on November 9, 1965 as part of an investigation to determine the effects of dredging and spoil disposal on the geology and biological communities of that area. This project was supported by the Philadelphia District, U. S. Army Corps of Engineers and the Bureau of Sport Fisheries and Wildlife, U. S. Department of the Interior, under Contract #14-16-005-2096, and the Natural Resources Institute, University of Maryland.

An analysis of the fish egg and larvae information obtained from all areas listed above indicated the feasibility of formulating a general characterization to describe organism abundance, distribution and movement in physically different but ecologically related areas of Chesapeake Bay.

The purpose of this report is to present the general characterization, the data from which it was derived, and to discuss in detail those species for which the nature of the data warrants elaboration.

Areas of Investigation

Chesapeake Bay is a large Atlantic coast estuarine system containing many tributaries which, together, drain an area of 67,505 square miles (Wells, Bailey, and Henderson, 1929) of New York, Pennsylvania, Maryland, Delaware, West Virginia, and Virginia. Each tributary is, in a sense, an individual estuary, differing in some respects from the




Figure 1. Fish egg and larvae sampling stations (●) in upper Chesapeake Bay 1960-1968. (Patuxent River to the Susquehanna and Delaware Rivers).

dominant branch, the bay proper. A description of the study areas for this investigation must therefore describe the bay region in general and salinity gradients of the Patuxent, Magothy, and Susquehanna, in particular.

The Patuxent River (Fig. 1) is situated entirely in Maryland and drains 963 square miles (Corps of Engineers, 1930). The river is approximately 80 miles long (Mansueti, 1961) and empties into Chesapeake Bay about 25 miles north of the mouth of the Potomac. Tidewater extends about 56 miles upstream. From Lower Marlboro south, the river increases in width from 0.1 to 2.0 miles. Mean tidal amplitude varies from 1.2 feet at Solomons to 2.5 feet at Nottingham, about 40 miles upstream. Channel depth generally increases downstream from Lower Marlboro to Benedict Bridge, ranging from 9 to 40 feet. From Benedict Bridge south, depths range from 33 to 120 feet. The lower section of the river possesses much sandy shoal area. The soil on the watershed consists of sand and clay with some limestone (Nash, 1947). Water velocity increases from about 0.9 feet per second at the mouth to about 1.6 feet per second near its upper limits. The lower Patuxent River is a typical two-layered estuary with net upstream movement of subsurface, high saline waters and a net down-

stream movement of fresher, surface waters (Pritchard, 1951, 1955). The upper river is bordered by many freshwater marshes. Further descriptions of this river appear in Nash (1947), Mansueti (1961), Heidel and Frenier (1965), Stross and Stottlemeyer (1965), and Herman et al. (1968).

The Magothy River (Fig. 1) is 7.0 miles long and drains an area of only 31.1 square miles. Most of the river is less than 20 feet deep; however, a constricted entrance to Chesapeake Bay aids tidal currents in maintaining a short 40-ft deep channel where the river and the bay meet. Freshwater inflow into the Magothy originates from Lake Waterford and direct watershed runoff. This flow is very small as compared with the flow into other tributaries of Chesapeake Bay. Salinity of the Magothy is greatly affected by the flow from the Susquehanna River, which drains into Chesapeake Bay approximately 40 miles north of the mouth of the Magothy. At times during spring runoff, lighter, less saline surface waters from the head of the bay flow into the Magothy River, creating a condition contrary to the normal estuarine chemical and circulation patterns. Salinity at the mouth of the river may be less than salinities found upriver (Pritchard and Bunce, 1959). Due to the limited input of fresh water and the shortness of the river, the salinity range for the entire river is small. Normally salinities between upriver and downriver stations do not vary more than several parts per thousand (‰). During the 15 months of investigation in this river, salinities at the uppermost sampling station ranged from 5 ‰ in late spring to 14 ‰ in late fall. A description of the Magothy River also appears in a report of the fish eggs and larvae of that river (Dovel, 1967).

Sampling in the upper Chesapeake Bay was performed along a 29-mile section from off Tolchester, Maryland, to Chesapeake City in the Chesapeake and Delaware Canal, plus the 8-mile channel section from Turkey Point to Havre de Grace in the Susquehanna River (Fig. 1). The 29-mile section is part of the main navigational channel of the upper Chesapeake and ranges in depth from about 55 feet off Tolchester to 35 feet in the C & D Canal. Water depth from Turkey Point to Havre de Grace fluctuates from 35 feet in the channel off Turkey Point to 20 feet for most of the 8-mile distance, then drops to 60 feet in the immediate vicinity of Havre de Grace.

Salinities in the upper bay fluctuate seasonally according to rainfall, but generally range from about 12.0 ‰ off Tolchester to 0.0 ‰ off Turkey Point (Stroup and Lynn, 1963). Water between Turkey Point and Havre de Grace is fresh and originates mostly in the Susquehanna River. This river accounts for slightly more than 50% of the fresh water contribution to Chesapeake Bay. The upper Chesapeake, leading into the Susquehanna River at Havre de Grace, also exhibits the two-layered circulation system described for the Patuxent River and will be referred to as the Susquehanna gradient.

The 27-mile Chesapeake and Delaware Canal, however,

presents a unique pattern of water circulation between two large bodies of water. Differing tidal cycles and amplitudes in Chesapeake Bay and Delaware Bay, plus water storage capacities of several reservoir-like areas located along the canal, serve to complicate water movement. Water salinity within the canal remained consistently between 0.0 and 4.0 ‰ during April, May and June of 1966 and 1967. These waters appear to be fairly well-mixed vertically. Haight, Finnegan, and Anderson (1930) provide the best account of water velocities and movements in the canal for the period immediately following the completion of the canal in 1927. At present, water current velocities through the canal reach a calculated maximum of 2.5 knots (U. S. Coast and Geodetic Survey, 1967), with an average velocity of about 1.9 knots. Gray (1967) and Burgess (1965) document many physical changes that have taken place in the canal area since 1764, when the first surveys were made. The texture of the canal bottom resembles, by visual inspection, that found further south in Chesapeake Bay in areas of comparable depths. Biggs (1967) states that in geological records, deepwater Chesapeake Bay sediments would probably be characterized as homogeneous gray-green organic shales containing pyrite and scattered shell and sand layers.

The Susquehanna River drains an area of 27,501 square miles (Carlson, 1968) of New York, Pennsylvania, and Maryland. The large fresh water contribution from the river has profound influence on the biota of a large area of the upper Chesapeake. This influence is a primary concern when investigating fish eggs and larvae of the upper bay. The industrial use of Susquehanna River water, plus the construction of many dams, has altered the physical, chemical, and biological characteristics of that river and the waters flowing into the upper Chesapeake. Many of the physical changes that have taken place in the river are discussed by Whitney (1961). Some biological implications resulting from these changes are discussed by Mansueti and Kolb (1953). Specific changes in the patterns of reproductive activities and commercial harvest of striped bass, *Morone saxatilis*, also attributable to physical changes in the lower Susquehanna, are discussed by Dovel and Edmunds (1971).

METHODS

The tools and techniques developed for a comprehensive investigation of fish eggs and larvae of the large Chesapeake estuarine system are documented (Dovel, 1964). The basic approach and recent modifications are discussed only briefly here. Sampling at 13 main channel locations along the Patuxent River from Lower Marlboro to Hoopers Island Light in Chesapeake Bay was begun in January, 1963. Meter net collections were made at surface, mid-depth, and bottom on a weekly frequency, using netting fabric with 0.4 x 0.6 millimeter (mm) apertures. A complete sampling run of 43 miles required about 15 hours of working time and was usually completed in a single day.

The processing of samples emphasized discarding or excluding extraneous material in the field (see filtering device; Dovel, 1964). The system made it possible to reduce most samples to less than 30 ounces in volume without the loss of desired organisms. Plankton net contents were concentrated on a filter of netting fabric, then placed in a 30-oz wide-mouthed jar containing about 10 oz of 4% formalin. Permanent, white plastic-tape labels on the exterior of the jar provided a reusable surface on which field data were recorded in pencil.

Sampling of the Patuxent River during 1964 and 1965 was reduced to a biweekly frequency at one-half the number of original stations (Fig. 1) after a review of 1963 data indicated that reduced frequency would be adequate to produce basic information. Techniques used to sample the Magothy River and upper Chesapeake during 1965-68 were comparable to those used in the Patuxent during 1963-65 with one slight exception. Upper bay sampling stations were divided about equally between shallow (10ft.) areas and deeper (20-60 ft) channel areas. Sampling stations for all areas were located at approximately equal intervals along the gradient with some alterations necessary for locating areas free of obstacles which might damage the plankton sled. All channel tows were made in a downstream direction at speeds ranging from 1-3 knots. Data from all tows were standardized to a 10-minute duration. Temperature and salinity readings were made with an induction conductivity salinometer.

Variations in sample size necessitated some aliquoting. Aliquots from samples containing small fish ranged from 1/4 to 1/10 of the original sample. Aliquots from samples containing mostly fish eggs were usually 1 to 10 milliliters (ml) in size.

RESULTS

Results are presented in two forms. First, data for individual species, although collected from several locations in the Chesapeake, are combined to provide basic abundance, distribution, and movement patterns for each species in the upper Chesapeake estuarine system. Second, data for some species are discussed in detail where subtle differences in biological patterns seem to exist but could be obscured by a broad summarization, or where additional comments seem warranted.

An evaluation of the ecological significance of the patterns of Chesapeake ichthyoplankton can only be made by correlating biological data with the estuarine environment. Since an estuary is characterized by a salt concentration gradient to which fish are known to respond, the salinity gradient appears to be the logical link between the biological data and the environment. The variation in the salinity gradient from year to year, as a result of varying freshwater input, precludes the assignment of a fixed salinity value for

a given geographical location. The validity of the major conclusions drawn in this report is subject to the acceptance of the salinity gradient as an adequate yardstick for the estuary.

The eggs and/or larvae (5,148,596) representing 48 species of fishes were collected and processed during the period January, 1960 to November 22, 1968 from all areas sampled (Tables 1, 2, 3, 4, 5, and 6.). Less than 0.1% of the total number of fishes collected could not be identified. This group, for the most part, is made up of common fishes which were found in a deteriorated condition when identification was attempted and will not be discussed in this report.

Although this investigation was directed primarily toward fish egg and larval stages, data for all fish collected which were not fully mature are included in this report. This criterion is necessary since larvae, juveniles, and adults are collected with differing degrees of efficiency as hydrographical conditions, mostly water temperature, and physiological responses change. Some juvenile and adult fish (many white perch), which usually would not be taken in warmer months, may appear in samples taken when fish activity is reduced during colder periods. The inclusion of data from these larger, almost mature fish which are still found in the nursery area is necessary in order to make a more complete determination of early life history movements of these species. The use of two calendar years in Table 6 assures the inclusion of all recorded captures of the young of a species during a complete year following the first occurrence of that species in the estuarine sampling area.

Table 7 presents numbers of bay anchovy larvae collected at different salinities. The combined distribution for all years (Grand Total) is used as generally indicative of organism distribution by salinity in the Chesapeake area.

Tables 8 and 9 are constructed after the pattern used in Table 7 and show the distribution and density of fish eggs and larvae in the upper Chesapeake Bay area as related to the salinity gradient. The patterns of abundance and distribution shown in these tables reflect contributions of a few dominant estuarine species. For example, 51.4% of all eggs came from waters with a 4 o/oo (12-15) salinity spread, with the bay anchovy and hogchoker accounting for 99% of the eggs in that range.

Tables 10 and 11 show the distribution of eggs and larvae as related to a water temperature gradient. Again, these tables reflect the predominance of eggs and larvae of a few dominant estuarine species. Eighty-three percent of fish larvae and juveniles were present during an 8-degree (22-29 C) temperature spread. This temperature range represents the period approaching maximum summer conditions. Table 12, Column A, presents the major species collected from the Patuxent River ranked by order of abundance for the period 1963-65. Column B provides numbers of the same fishes collected from the upper bay during 1966-67.

DISCUSSION

Species for which data were abundant were considered for discussion in detail in this section. Species for which biological patterns appear to be sufficiently interpretable through tabular presentation (Tables 1-12) are not discussed.

The abundance of fish eggs and larvae of some species that spawn in fresh water is directly correlated to individual species characteristics of egg deposition and attachment. The location of egg deposition in a tributary, plus the transporting effect of water currents, may suggest patterns of abundance not in keeping with actual biological production. For example, herring eggs which normally adhere to the substrate upstream in fresh water may become dislodged and move downstream. Collections made downstream may therefore represent only those eggs which have been transported away from their natural habitat and may not actually indicate species productivity in the area of occurrence. Movement in the opposite direction, where larvae from eggs deposited in the marine environment appear in brackish water samples, will also be discussed under appropriate species.

The eggs of many species which spawn in brackish water adhere to the smooth surfaces of oyster shells and other hard, submerged objects in shallow water areas. The resulting larvae apparently remain close to the spawning area and may not appear in samples collected only a short distance away in deeper water. This pattern of distribution is advantageous in estuarine areas where dissolved oxygen levels are reduced in summer in depths below about 50 feet and food organisms are fewer in number.

Species discussed in this section will be grouped according to the salt content of the environment where spawning takes place. This will provide a framework through which the reader should be able to grasp the concept of broad patterns of activity which are correlated with specific freshwater, estuarine, and marine environments.

Freshwater spawners

Family Clupeidae - herring

Clupeid eggs and larvae were present in all collecting areas but not necessarily at the same times or in comparable numbers (Tables 3-12). The eggs collected resulted from deposition upstream, dislodgment, and transport downstream by freshwater runoff. Four species are listed (Table 3); however, the great percentage of these eggs belong to two species: blueback herring (*Alosa aestivalis*) and alewife (*A. pseudoharengus*). Egg condition (state of decomposition), the accumulation of detritus on the egg membrane, and general morphological similarities between these two species prohibit positive identifications. The eggs of *A. mediocris* and *A. sapidissima* are somewhat larger and should not be confused with *A. aestivalis* and *A. pseudoharengus* which

are smaller and both about 1.0 mm in diameter (Mansueti and Hardy, 1967).

Clupeid larvae, possibly belonging to four species of *Alosa*, and one each of *Brevoortia (tyrannus)* and *Dorosoma¹ (cepedianum)*, were present in all areas. Although some differences exist in size at hatching and during comparable stages of larval development, size range overlap and similar morphological indexes prohibited positive identification of the majority of the newly-hatched specimens of the genus *Alosa*. Peritoneum color normally serves to distinguish juvenile blueback herring from the alewife. Hybridization and possibly the physiological state of the organism at time of preservation, as it affects the contractibility of melanophores, makes a color differentiation between dark and dusky sometimes very difficult. The 17,377 clupeid fish (Table 4) all possessed slightly pigmented peritoneums. It seems probable that both *A. pseudoharengus* and *A. aestivalis* are present in this group. Table 13 provides length frequency data for blueback herring, *Alosa aestivalis*, collected from two areas of the Chesapeake and possessing pronounced blueback characteristics. Data suggest the presence in the upper estuary of two distinct age groups, young-of-the-year and fish about 1 year old. Fish almost 1 year of age are present there in early spring, but were not found in later collections. Young-of-the-year and older fish were both found in the same area of low salinity water. Close inspection of organism size, as compared with water salinity at collection locations, suggests movement of maturing organisms toward higher salinity areas. While developing fish should show the tendency to spread throughout the estuary, it is also true that the salinity gradient varies with the season. Salt water penetrates further upstream during late summer and fall. The location of capture could conceivably remain the same while water salinities at a given location vary appreciably according to season.

Numbers of blueback herring (*A. aestivalis*) larvae and juveniles are insufficient for determining a definitive growth rate; however an approximation of this rate could be derived from the data in Table 13. This data would suggest that fish one year of age are about 80.0 mm total length (TL) or slightly larger and about 65.0 mm when six months old, using April 15 as the hatching date.

The scarcity of larvae less than 30 mm TL in Table 13 resulted from the inability to distinguish between the early developmental stages of different species of the genus *Alosa*, as mentioned above. The lack of specimens in collections taken in low salinity areas during periods of extreme cold raises the question as to whether these fish (0-year-class) left the upper estuary or became inaccessible for the period November through February. The latter seems logical as these fish were present in the same low salinity area the following spring before apparently moving toward areas of higher salinities.

¹Not otherwise listed in this report.

In summary, it would appear that, following spring hatching upstream in fresh water, young blueback herring and other members of the genus *Alosa* move slightly downstream but remain in the upper estuary until about 1 year of age, at which time they become less susceptible to plankton gear and move toward the lower estuary and the ocean. Movements through the estuary appear to be rapid. Fish appear to congregate and move in schools; major movement taking place in June or early July.

All *Alosa* larvae and juveniles were found in fresh water or waters of 12 o/oo or less salinity.

White perch, *Morone americana*

The abundance and distributions of eggs and larvae of the white perch are subject to some of the same conditions described for the freshwater clupeids. Mansueti (1964) states that great variation occurs in the attachment rate of detritus to eggs (or substrata) during different developmental stages. This factor, plus variations in water currents in the upper estuary, precludes a determination of the proportion of eggs that are transported downstream. The relatively large number of eggs recorded in Table 3 represents only eggs that have been transported away from the area of deposition but suggests a higher reproduction potential for this species than for many others in the Chesapeake estuarine system.

Eggs were found in waters having a temperature as low as 5 C; however, most occurred when temperatures ranged between 8 and 16 C, with some occurrence when temperatures were as high as 24 C (Table 10). Egg occurrences, as determined for this study, varied slightly from that documented in a comprehensive description of the eggs, larvae, and young of this species in the Patuxent River (Mansueti, 1964).

Larvae, which hatched upstream in fresh water, were present downstream in the sampling areas of all tributaries sampled. Mansueti (1964) suggests that low catches of larvae in the upper Patuxent River in fresh water were due to the demersal behavior of these fish and their swift transport downstream. Most larvae were found in fresh waters or waters having a very low salt content. This pattern was observed for the Susquehanna River during 1966 and 1967 when large numbers of white perch larvae moved out of this river and into the upper bay.

Following hatching in late March, April, and May, young white perch moved, mostly as planktonic larvae, toward brackish waters. Larvae remained abundant in low salinity waters until winter, when their numbers began to decline. From the following spring, when these fish were 1 year old, until fall, when about 18 months old, this year-class was present in the same area but in ever decreasing numbers. Their absence from collections was probably due to decreased efficiency of collecting gear as these fish grew, plus dispersal away from the nursery area. It would appear that at least some juvenile white perch remain in the most critical nursery environment for about one year.

Although spawning takes place over a rather short period, the variations in hatching dates contribute to the presence of a large size-range of individuals of the same year-class at any given time during the first summer. Table 14 provides length frequency data for white perch collected by and subject to the selectivity of the plankton gear described. It would appear that these data provide a fair representation of the growth for the first year of life. It was noticed, during the first winter of this investigation, that young-of-the-year fish and representatives of earlier year-classes were present concurrently on the nursery ground. The possibility that developing white perch return to the nursery ground during winter for several years should be investigated. Dovel et al. (1969) describe this type of movement for the hogchoker from the same area. Mansueti (1961, 1964) discusses the life history and some movements of this species in detail.

Striped bass, *Morone saxatilis*

Eggs of this species were present in low saline areas of the Patuxent River and upper Chesapeake Bay (Tables 3 and 8), but were not found in the Magothy River (Dovel, 1967). Striped bass eggs are normally deposited in fresh waters not far above brackish areas in Chesapeake Bay. There is some evidence² suggesting that spawning of this species may occur in low salinity waters of some Chesapeake tributaries. Tidal actions, freshwater runoff and turbulence produced in the vicinity of the salt-fresh water interface, the transitional zone where fresh and salt water meet, maintain necessary egg suspension. Since sampling was performed in the vicinity of the interface, the numbers of eggs collected should be correlated with total abundance. Eggs were collected from waters with an 11 to 24 C range in temperatures. Temperatures above 12 and 13 C appear conducive to unlimited spawning when water temperatures follow a gradual increase (Table 10). Eggs were abundant during the months of April, May and June (Table 15). Close inspection of the data indicates that although the period of egg occurrence varies according to season, generally the peak production period extends from mid-April to the last week of May, a period of about 6 weeks. Very few eggs were found after the first of June of any year. No appreciable number of eggs were found in areas where salinities were greater than 3 o/oo. Large numbers were found in the Chesapeake and Delaware Canal (Table 16) where salinities were greater than 0.1 o/oo, but less than 4.0 o/oo. No eggs were found in the 8-mile section of fresh water from Turkey Point to Havre de Grace during the spring of 1966. Forty-four eggs were collected in the same area during 1967 (Table 16).

Determinations of vertical distributions of eggs were attempted without the benefit of an opening and closing net. The numbers from bottom samples listed in Table 16 are therefore subject to vertical contamination as the plankton

²Chesapeake and Delaware Canal study now in progress, and Joe Boone, pers. comm.

sled passes through surface waters where suspended eggs occur. Although the extent of vertical contamination has not been determined, live eggs do not appear to be stratified. They appear fairly well dispersed throughout the water column as a result of the mixing forces of the estuary.

Larvae of this species were generally most abundant during the month of May. This was to be expected as the incubation period under normal conditions is less than two days. Larvae were found in greatest density in low saline waters (< 2.0 ‰, Table 9). Areas of egg and larvae occurrence appear to be, at most, only slightly separated. This pattern appears to be characteristic of most upper Chesapeake tributaries but differs from patterns found in North Carolina and California estuaries.

Length frequency data collected for striped bass are not sufficient to calculate a growth rate for young fish but do provide an indication of the length of time developing striped bass could be collected with plankton gear in the nursery area. Generally speaking, young-of-the-year striped bass were not collected in appreciable numbers in the low salinity nursery area after June or July in any area sampled. Twenty-nine fish ranging in size from 64.2 to 119.9 TL (average 84.6 TL) were collected from a low saline (0.0 to 11.0 ‰) area of the upper Chesapeake Bay during January through March, 1967. This information suggests that at least some small young-of-the-year fish will be found in the nursery area when almost one year old. It is not certain whether these fish remain there constantly during this time or move about in the estuary. Two fish, 32.0 and 35.0 mm TL, were collected from waters of 16.0 ‰ as early as mid-June, 1963.

A special attempt was made to determine the ratio of young striped bass to white perch on the nursery ground in upper Chesapeake Bay during 1966. Only fish possessing distinguishing characteristics were used. Newly-hatched white perch and striped bass up to about 6.0 mm TL can be separated with ease (Mansueti, 1958; 1964). Fish from about 6.0 to 12.0 mm TL cannot be separated by the usual meristic and morphometric characters. Fish larger than about 12.0 mm TL normally possess a full complement of spines and fin rays which can be used to separate species. Fish with twelve or less total spines and rays in the anal fin were classified as white perch (Mansueti, 1958; 1964). Those with 14 or more were identified as striped bass. Fish with 13 anal rays and spines were not used, as these fish could belong to either species.

A total of 3,834 fish were used for anal fin counts. Of these, 1,116 were eliminated due to fin condition or fin counts of 13. Of the remaining 2,718 fish, 2,606 or 68.0% were identified as white perch. This produced a ratio of 1 striped bass for every 23 white perch of comparable stages of development in the upper Chesapeake. In all probability, the true identification of the 1,116 unused specimens would alter this ratio only slightly.

Dovel and Edmunds (1971) discuss changes that have

taken place in striped bass activity in the upper Chesapeake Bay in recent years. The abstract of that paper is presented here.

"Recent changes in striped bass (*Morone saxatilis*) spawning sites and commercial fishing areas in upper Chesapeake Bay; possible influencing factors."

Abstract

Chesapeake Bay annually contributes a large percentage of all the striped bass produced in North America. For many years, the lower Susquehanna River was considered the major reproductive area for the entire bay. Recent evidence suggests that the areas of greatest egg abundance, as well as commercial fishing for the striped bass in the upper bay now occur along the main channel from Worten Point to Chesapeake City on the Chesapeake and Delaware Canal. The shift has apparently resulted from major environmental alterations which have occurred in the last 90 years. The construction of the C & D Canal, at the head of Chesapeake Bay, has increased the total area suitable for species propagation and commercial fishing activities.

Yellow perch, *Perca flavescens*

Larvae of the yellow perch were most abundant in very low salinity waters (Table 9). This species, like the freshwater herring group and the white perch, spawns far upstream in fresh water. Their larvae then move downstream toward brackish waters. Yolk-sac absorption was complete by the time larvae were found in the low salinity nursery area by early April. Their growth rate is suggested in Table 17. Water temperatures, at times when larvae were abundant, ranged from 8 to 17 C (Table 11).

Estuarine Spawners

Bay anchovy, *Anchoa mitchilli*

Anchovy eggs were abundant in upper Chesapeake Bay from April to September of every year during 1963-67. Eggs were present as early as April 22 (1963) and as late as September 27 (1965, Fig. 2). Peak production occurred during July (Table 18, Fig. 2). Some eggs were found over a wide range of salinities (1 to 22 ‰) but the area of greatest abundance occurred in waters with salinities of 13 to 15 ‰. This area contributed 47.6% of all eggs collected. Patterns of egg distributions along the Patuxent salinity gradient for 1964 and 1965 indicated large numbers of eggs at 4 and 6 ‰ (Figs. 3 and 4). These concentrations were exceptions rather than the rule. Most eggs were present with water temperatures in excess of 20 C (Table 10, Fig. 2). A table of egg sizes (Table 19) indicates that eggs collected

from more saline areas were smaller than eggs which came from low saline waters.




Figure 2. Seasonal occurrence of the eggs of the bay anchovy (*Anchoa mitchilli*) plotted against total hours of daylight and average surface water temperature at Solomons, Maryland, 1963-1965.

Larvae and juveniles were abundant throughout most of the year over much of the salinity gradients sampled (Tables 7, 9, and 20). Juveniles were most dense in low salinity areas upstream (Table 9). Seventy-two percent of all fish collected were found in waters of 3 to 7 o/oo. Vertically, the larvae appeared to be concentrated in surface waters (Table 20), at least during 1963. Some larvae or juveniles were collected in waters at all temperatures up to 31 C. Eighty-eight percent of the total was found when temperatures ranged between 23 and 27 C. Most of these fish were abundant in shallow water areas during summer (June to September). In colder months (October to March), fish were not as plentiful in samples and those present were found in deeper waters. Anchovies were abundant in low salinity waters at the head of Chesapeake Bay just south of Turkey Point and in the C & D Canal. Fish were present but not plentiful in fresh water in the Susquehanna River west of Turkey Point toward Havre de Grace.




Figure 3. Distribution and abundance of the eggs and larvae of the bay anchovy (*Anchoa mitchilli*) in the Patuxent River in 1964 by salinity (o/oo) at location of capture.

Anchovies of the family Engraulidae are an important source of fresh fish, live bait, fish meal and a substitute for sardines along the coastal areas of most of the world's

oceans. The north Atlantic coast of America is one of the few major coastal areas where anchovies are not utilized commercially (Hildebrand, 1943; Baxter, 1967). Anchovies of middle Atlantic Bight estuaries belong to two genera, *Anchoa* and *Anchoviella*. The bay anchovy, *Anchoa mitchilli*, is very abundant in the upper Chesapeake Bay but rarely exceeds 97 millimeters in length (Hildebrand and Schroeder, 1927). This species extends from Maine to Yucatan, Mexico (Mansueti and Hardy, 1967), and is abundant along most of the middle Atlantic coastal region of North America. Richards (1959) stated that numbers of anchovy eggs dominated the total abundance of fish eggs in the Long Island Sound area. Massmann (McHugh, 1967) found the anchovy to be "the most abundant species in Chesapeake Bay." Many investigators have recorded biological data relative to the bay anchovy, but very little information exists concerning movements of the early life history stages of this species in estuaries. Kuntz (1914), Hildebrand and Cable (1930), Hildebrand (1963b), Richards (1959), Wheatland (1956),




Figure 4. Distribution and abundance of the eggs and larvae of the bay anchovy (*Anchoa mitchilli*) in the Patuxent River in 1965 by salinity (o/oo) at location of capture.

and Herman (1958) provide information on the embryology of this species. Distribution information is available from many sources. Juveniles and adults are found primarily in estuaries and coastal waters in all types of environments; from muddy coves (Hildebrand and Schroeder, 1927), in grassy areas (Hildebrand, 1963b), in bayous (Springer and Woodburn, 1960), off sandy beaches (Reid, 1954; Kilby, 1955), and in deep offshore waters (Hildebrand, 1963b). Juveniles have ascended rivers at least 40 miles above brackish water (Massmann, 1954). Massmann (1962) records juvenile and adult anchovies in the York River System, Virginia, concentrated in two distinct locations, in fresh water and areas with 12 to 32 o/oo salinity. The data of Stevenson (1958) and Massmann (1953, 1954), the latter summarized in McHugh (1967), is perhaps the best available information on the abundance and distribution of this species in estuaries of the middle Atlantic region.

Egg and larval stages of the bay anchovy were very abundant in the upper Chesapeake area in low to intermediate salinities (0 to 21 o/oo). A single 10-minute plankton tow produced 141,440 eggs from 330 m³ of water. This

species shows some life history aspects similar to those of the hogchoker (Dovel et al., 1969). These two species provided the majority of the free-floating fish eggs collected in this investigation.

The anchovy, like the hogchoker, spawns throughout much of the lower estuary during midsummer conditions when water temperatures are at a maximum (Fig. 2). This information agrees with Stevenson (1958), who determined egg maturity for anchovies collected in Delaware Bay. The deposited eggs possess no oil globules and apparently occur throughout the water column. Vertical sampling contamination obviously accounts for an exaggerated egg distribution pattern, as well as an unreal range in water temperatures at which eggs are found. For example, eggs recorded as coming from a bottom sample where water temperature was 9.7 C (1963 data) may actually have occurred in an intermediate or surface sample where the temperature was as much as 4 degrees warmer.

Anchovy eggs collected in the lower salinity waters of the Patuxent River were appreciably larger than eggs of this species collected in ocean waters off Beaufort, North Carolina. Kuntz (1914) gives the length of the major axis of anchovy eggs collected at Beaufort at 0.65 to 0.75 millimeters with the minor axis 0.1 to 0.3 mm less. Eggs from the Chesapeake show a definite size-salinity correlation which could account for the differences between Chesapeake and Beaufort egg sizes. The incubation period is about 24 hours (Mansueti and Hardy, 1967) which would not permit the eggs to be carried by tidal currents many miles from the area of deposition. The distribution of anchovy eggs in the Chesapeake should, therefore, be generally indicative of the spawning area of this species.

Newly-hatched larvae were not abundant in samples taken from any area of the estuary. Slightly older larvae and juveniles, > 5.0 mm, were abundant and found upstream in low saline areas suggesting a migration to an upstream feeding or nursery area. The general trend of movement is shown in Figs. 3 (1964) and 4 (1965) but is somewhat complicated by the presence of what appear to be isolated groups of eggs located in salinities lower than those where eggs were most abundant. This movement was most obvious when viewing data collected with a weekly sampling frequency in the Patuxent River during 1963 (Fig. 5). An attempt to monitor movement of the larvae and juveniles as they moved upstream did not succeed. Small, recently-hatched larvae were found almost simultaneously in both upstream and downstream areas. Since most of the eggs are deposited in areas with salinities of 9 ‰ or greater, the occurrence of recently-hatched larvae upstream in low salinities suggests rapid movement of fish upstream toward the nursery ground.

Young anchovies were found predominantly in surface waters (Table 20) from about the first of May to the middle of October. These fish apparently moved out of surface waters when water temperatures dropped below about 11 C.




Figure 5. Distribution and abundance of the eggs and larvae of the bay anchovy (*Anchoa mitchilli*) in the Patuxent River in 1963 by salinity (‰) at location of capture.

During the colder period, December to May, young-of-the-year fish were collected in limited numbers in low salinity areas. During April until July, juvenile fish about one year of age and recently-hatched young-of-the-year were present concurrently on the nursery ground and could be separated according to size (Table 21). Fish collected during August through November included both young-of-the-year and older fish but could not be separated with certainty on the basis of length frequency (Table 21).

By late fall or winter, only young-of-the-year fish remain on the nursery ground. Fish of the previous year-class, now about 15 months old, appear to move from the area or become less vulnerable to plankton gear as water temperatures drop with the onset of colder weather. Young-of-the-year will be found in the nursery areas the following spring.

The upstream limits of the anchovy nursery area could not be determined from collections made in the Patuxent River. A plot of total number of fish collected during 1963 by location of capture (Fig. 6) showed the greatest abundance of fish (67.4%) at the sampling station (0) closest to freshwater. No determination could be made as to whether areas slightly further upstream in fresh water of




Figure 6. Distribution and abundance of bay anchovy (*Anchoa mitchilli*) by collection station in the Patuxent River during 1963.

the Patuxent would have contributed more or less organisms.

Sampling at the head of Chesapeake Bay provided data helpful in evaluating the importance of fresh water as a

nursery environment. Sampling there was performed along two routes. One route extended along the navigational channel from off Tolchester to Havre de Grace in the Susquehanna River (Fig. 1). This route extended into fresh water by at least 8 miles at all times of the year. The other route extended along the main navigational channel from off Tolchester to Chesapeake City on the C & D Canal. Both ends of the canal route were located in low saline waters at all times of the year. Water in the short section from Old Courthouse Point to slightly south of Turkey Point was usually fresh. Developing anchovies were more abundant in the canal area than the lower Susquehanna River area (Havre de Grace to Turkey Point, Fig. 7). A plot of collections by station (Fig. 7) and salinity for 1967 (Fig. 8) indicated an abundance of larval stages in low salinity areas. The lack of fish at freshwater stations IE and Old Courthouse coincides with reduced water salinities in those areas. Havre de Grace and Fishing Battery stations, located in fresh water (Fig. 7) produced less than 1.0% of the total fish collected during either year.




Figure 7. Distribution and abundance of bay anchovies (*Anchoa mitchilli*) by collection station and water salinity in upper Chesapeake Bay during 1966 and 1967.

HG = Havre de Grace
 FB = Fishing Battery
 CC = Chesapeake City
 OC = Old Courthouse

I-E } Channel stations
 II-D } between
 III-C } Turkey Point
 IV-D } and Tolchester
 V-F }

A plot of the distribution of both eggs and larvae from the salinity gradient leading into the canal (to Chesapeake City) or Susquehanna River (to Havre de Grace) for 1967 (Fig. 8) suggests a pattern of upstream migration or movement of larvae and juveniles similar to such a movement in the Patuxent River (Figs. 3, 4, 5, and 6). Close inspection of the data shows that many of the fish recorded as occurring in fresh water actually occurred in water of 0.5 to 0.9 ‰ salinity. It is therefore apparent that in reference to a salinity gradient leading into the Susquehanna River, some few juvenile anchovies may be found upstream in fresh water, but the majority of these fish congregated in brackish




Figure 8. Distribution and abundance of the eggs and larvae of the bay anchovy (*Anchoa mitchilli*) by salinity in upper Chesapeake Bay during 1967. Larvae data are plotted along two sampling routes leading into the Susquehanna River and the Chesapeake and Delaware Canal.

waters very close to the salt-fresh water interface. A comparison of the concentrations of fish in the Patuxent and Susquehanna salinity gradients indicated a slightly different distribution pattern (Table 7). This difference appears to have been due to sampling effort as related to a shifting salinity gradient. An upstream shift of salt waters of the Patuxent during fall months precluded equal and constant sampling of very low salinity waters since sampling locations remained constant. Sampling effort for the Susquehanna adequately monitored the low salinity end of the gradient and adjacent areas of fresh water. This information would suggest that the distribution pattern determined for the Susquehanna gradient, i.e., concentration in brackish areas very close to the salt-fresh water interface, may be typical for other areas in the upper Chesapeake. There is, of course, some movement of a few fish well upstream, and spillover of many into fresh water bordering brackish areas.

The anchovies of the family Engraulidae reach sexual maturity at an early age. Hildebrand and Cable (1930) concluded that *A. mitchilli* may spawn when 21½ and 3 months old. Information presented in Table 21 tends to substantiate the claim of maturation in this short duration if maturity is attained by the time fishes reach 40 mm TL. Stevenson (1958) states that the bay anchovy reaches a maximum standard length (SL) of 75-80 mm in its second year. The largest fish recorded in Delaware Bay by Stevenson (1958) was 86 mm SL.

Table 21 indicates the presence of at least two year-classes of this species in the upper reaches of Chesapeake tributaries during early spring and summer. This information is based on collections made with plankton nets. Trawl catches of adult fish³ from part of the same area (Patuxent River) and period do not indicate the presence of large numbers of older fish. All available information suggests that this

³A. J. McErlean, pers. comm.

species has a short life span in Chesapeake estuaries. We will assume that all bay anchovies are mature when one year of age. The rapid maturation and prolonged spawning season of this species in Chesapeake Bay contributes to a large total size range for young-of-the-year fish (Table 21). Perlmutter (1939) records 0-year-class fish from Long Island Sound as 20-55 mm. The potential for spawning late during the same year in which they were hatched may contribute to distinct early and late season spawners of this species, as mentioned by Stevenson (1958).

In summary, the bay anchovy produces large numbers of pelagic eggs and larvae in upper Chesapeake Bay tributaries and the bay proper. Spawning takes place along most of the salinity gradient with large concentrations of resulting larvae found in low salinity nursery areas. This species matures quickly, possibly before one year of age, and apparently has a short life span. The characteristics of spawning in one location followed by movement of larvae toward a low salinity nursery area is similar to the early life history movements of the hogchoker (Dovel et al., 1969).

At present, man utilizes the anchovy only indirectly in the Chesapeake Bay area. Here, as a forage fish, it constitutes a major portion of the diet of the striped bass, *Morone saxatilis* (Raney, 1952; Hollis, 1952; Dovel, 1968). We cannot overlook the possibility that this fish could also be an important food item for waterfowl and other animals. The abundance of anchovy larvae and juveniles in accessible estuaries makes this organism vulnerable to exploitation as a possible untapped natural resource. The prolific nature of this species, its rapid maturation and short life span favor exploitation. Its small size may be the primary reason why this anchovy is not used commercially, as are other larger members of the family Engraulidae. Any future plans for exploiting the bay anchovy must, however, take into consideration the true value of this species as a forage fish. In addition, the selective or unselective characteristics⁴ of potential harvesting gear must be considered in relation to adverse effects such gear might have on other important species.

Northern pipefish, *Syngnathus fuscus*

Adults of the northern pipefish are common during warmer months in vegetated areas in the vicinity of Solomons. Recently-born larvae of 10-12.0 mm TL usually appear about the middle of May of each year. Individual or combined collections made during the summers of 1961-67 seem to indicate two spawning peaks about early May and late July (Tables 22 and 23). Growth appears more rapid following the second hatching about July, when water temperatures are approaching their maximum for the year. Young of this species appeared over most of the salinity range sampled for this investigation (Table 9).

⁴E. Dunnington, Pers. comm.

Naked goby, *Gobiosoma bosci*

Demersal and attached eggs of this goby are common in oyster shells in shallow areas in the vicinity of Solomons, Maryland. Larvae were apparently restricted close to egg deposition areas since few were collected in deeper channel areas in the lower Patuxent River with intensive sampling effort. Larvae of this goby were more abundant than similar stages of other gobies collected during this investigation (Tables 9, 11, and 12). In fact, naked goby larvae accounted for 55.2% of all fish larvae collected. Young gobies were most dense in low saline areas of 1 to 12 o/oo (Table 9). Some were collected as early as June 4 (1963) with large numbers present in most areas soon after mid-June of each year. Size data in Table 24 suggest that spawning takes place for a 2- to 3-month period. Larvae were generally abundant until about October or during the warmest period of the year. During 1963, 94.0% of the larvae collected occurred when temperatures were between 22 and 29 C with the first occurrence at 18.9 C.

Recently-hatched larvae < 12mm TL appear to be efficiently collected by the plankton gear used for this study (Table 24).

A total of 149 larvae were found west of Turkey Point (at Fishing Battery and Havre de Grace) in fresh water of upper Chesapeake Bay. Large numbers were collected from low salinity waters of the C & D Canal. No appreciable numbers of larvae were found in waters having salinities over 18 o/oo. Sampling effort was, however, reduced in these waters.

| | Silversides | |
|--------------------------|--------------------------|------------------------|
| | Tidewater | Atlantic |
| Rough | | |
| <i>Membras martinica</i> | <i>Menidia beryllina</i> | <i>Menidia menidia</i> |

The Maryland portion of the Chesapeake possesses the species listed above, all of which have demersal and attached eggs and morphometrically similar newly-hatched larvae. Positive identifications were made for fish which possessed a full complement of fin rays and scales. The anal fin count, relative positions of the dorsal and anal fins (Hildebrand and Schroeder, 1927) and scale counts (Bayliff, 1950) were used to differentiate species. All three species were common in low saline areas (Table 9) during the period April to December. Temperatures during this time ranged from about 12 C to maximum summer conditions approaching 30 C. Larvae were most common in surface waters. During 1966 and 1967, young silversides were present at upper bay stations located both in fresh water west of Turkey Point and in low saline waters in the C & D Canal.

Winter flounder, *Pseudopleuronectes americanus*

A single winter flounder egg was collected off Cove Point in 20 feet of water on March 14, 1966. Water temperature and salinity were 3.0 C and 20.7 o/oo respectively. The

collection of this egg was unusual as eggs of this species adhere to the substrata and are not available to plankton gear. The egg was apparently scraped from the bottom of the bay by the apron attached to the front of the plankton sled (Dovel, 1964).

Larvae were present in areas where water salinities ranged up to 20 o/oo (Table 9), during the period February to May. Areas of peak abundance were found only where salinities ranged between 6 and 15 o/oo. Larvae were most abundant for a 1-month period of each year, usually during the last half of March and first half of April. Water temperatures during this time ranged from 3 to 18 C with most larvae present when temperatures ranged between 4 and 13 C (Table 11).

An inspection of data by vertical location of capture did not suggest definite vertical stratification. Samples from the Magothy River during 1966, however, indicated most larvae were found close to the bottom. This characteristic may be partly attributed to the descent of larvae that swim to the surface, then while resting, sink for a short period of time. Any descent in the shallow Magothy River (less than 20 feet) brings these fish close to the bottom. Samples from the Patuxent River (35 to 50 feet depth) did not suggest concentrations at either surface, mid, or bottom depths in 1963. The water depth, sinking rate of the larvae, lengths of their resting periods and response to illumination may thus affect the level at which these organisms are found.

All but 2 of the recently-hatched winter flounders collected were less than 11.0 mm TL (Table 25). It would appear that collecting methods employed in this study efficiently captured larvae up to about 8.0 mm TL. Fig. 9 provides an approximation of winter flounder growth during the period February to May for four different years. No migration of newly-hatched larvae from hatching area to a separate nursery area was noted.




Figure 9. Length frequencies of winter flounders, (*Pseudopleuronectes americanus*), in different locations of upper Chesapeake Bay (1963-1966), showing similarity of seasonal growth for different year classes.

Hogchoker, *Trinectes maculatus*

Dovel et al. (1959) provide an account of some aspects of the early life history of this species summarizing all information collected during the period 1963-68. The abstract of that paper is presented here.

"Some life history aspects of the hogchoker, *Trinectes maculatus*, in the Patuxent River estuary, Maryland."

Abstract

Information is presented on the abundance, distribution, and seasonality of hogchoker life history stages in the Patuxent River estuary. Egg collections indicate that the spawning area is located in the lower river in salinities greater than 9.0 o/oo. Following hatching during July and August, the larvae move upstream and congregate in a low salinity nursery area close to the salt-freshwater interface where they remain during winter. As spring approaches, the juveniles move toward the spawning area. These two distinct movements, upstream toward the nursery area in fall and downstream toward the spawning area in spring, apparently continue at least through the fourth year. As the fish mature, they progressively increase their range of travel away from the nursery ground toward higher salinities. Life history activities were determined by monitoring an abundant 1963 year-class using three sampling techniques: meter nets, shallow water trawls and deep water trawls. The importance of various segments of the estuary for completion of life history cycles is emphasized.

Skilletfish (clingfish), *Gobiosox strumosus*

Skilletfish, like gobies, are abundant in the lower Patuxent River. This species is associated with the oyster community where its eggs are deposited on hard substrate, mostly oyster shells. This species did not appear in very low salinity waters (Table 9). In 1963, larvae were present from May 13 to September 5, mostly in waters with salinities of 4 to 19 o/oo. Water temperatures ranged from 17 to 27 C. A description of the early development of this species in the Solomons area may be found in Runyan (1962) and Dovel (1963).

Marine Species

Atlantic menhaden, *Brevoortia tyrannus*

Menhaden eggs were found in the vicinity of Solomons for the first time in the spring (May and June) of 1963. They have since been collected there also in the fall (October and November) of 1967 and 1968. Water temperatures ranged from 13 to 20 C with salinities of 10 to 22 ppt.

The possibility exists that these eggs may have been transported a short distance upstream by a subsurface water current described by Pritchard (1951). A short incubation period (ca. 48 hrs at 22 C; Mansueti and Hardy, 1967) precludes movement over any appreciable distance. Although no adults in spawning condition have ever been reported from the upper portion of Chesapeake Bay, it is apparent that some eggs are occasionally, if not regularly, deposited in this area. These eggs have not been found north of the mouth of the Patuxent River in Chesapeake Bay. Young menhaden appeared in upper estuarine, low salinity areas (Table 9) in upper Chesapeake Bay tributaries during the period late March to late June. The majority of the 2,322 fish recorded in Table 4 ranged in size from 20 to 40 mm TL. Many of these fish were at least 25 mm TL, when they first appeared in collections in early spring (Table 26). We must conclude that these fish hatched and entered the upper estuary the previous fall or winter.

Only 25 specimens less than 20.0 mm TL were collected. Twenty-four of these (14.4 to 18.4 mm TL) were found in the C & D Canal on June 5, 1967. We can only wonder what significance there may have been in the collection of slightly smaller than normal fish in the canal which affords direct and fast access to the Atlantic Ocean through Delaware Bay. A specimen, 10.4 mm TL (Table 26, A) was found off Cove Point on November 7, 1967. This fish may have been on its way toward an upstream nursery area after hatching in the lower bay or in the Atlantic Ocean. Limited data (Table 26) suggest that these fish do not grow appreciably during winter or early spring but show accelerated growth during May and June before disappearing from plankton samples. Their disappearance is probably due to increased organism activity, gear avoidance and movement away from nursery areas where organisms are densely congregated. Accelerated growth during May is also characteristic of most other species recorded in this report.

Atlantic croaker, *Micropogon undulatus*

Dovel (1968) discusses aspects of the early life history of this species as related to a predator-prey relationship involving adult striped bass during winter months in mid and upper Chesapeake Bay areas. The information presented in that article was collected as part of this study. The abstract of that paper is presented here.

"Predation by striped bass as a possible influence on population size of the Atlantic croaker."

Abstract

Recent data collected from the upper Chesapeake Bay, Maryland and a review of commercial fishery statistics suggest that predation by striped bass on post larvae and juveniles of the Atlantic croaker, during periods of low water temperatures, may be a significant factor influencing

the recent decline of the croaker on the northeastern coast of North America. The life history of the croaker, stomach analysis of adult striped bass, and commercial fishery statistics for both species during the period 1930-1965 are discussed in this connection.

Southern Kingfish, *Menticirrhus americanus*

Eggs probably of a sciaenid species and tentatively identified as *Menticirrhus americanus* (Table 8) were found in bottom samples collected along the channel of Chesapeake Bay from south of the Chesapeake Bay Bridge to Hoopers Island Light. The estuarine circulation pattern could affect the distribution of these eggs, as mentioned for menhaden eggs.

Eggs were found during May and June of 1965, 1966, and 1967. Salinities ranged from 15.9 to 19.5 ‰ with a water temperature range of 12.5 to 23.8 C. Eggs collected in May 1966 were maintained live and allowed to hatch. Resulting larvae survived for only 7 days, but were helpful in making the tentative identification. Larvae (Table 9) were collected in field samples in May 1966, September 1963 and September 1965.

Seaboard goby, *Gobiosoma ginsburgi*

One hundred sixty-six specimens were found in Chesapeake Bay channel locations between the Chesapeake Bay Bridge and Hoopers Island Light. Salinities ranged from about 17 to 22 ‰ (Table 9) during June to December when larvae were present. *Gobiosoma ginsburgi* was found concurrently with *G. bosci* but was apparently restricted to higher salinity areas, whereas *G. bosci* was very abundant throughout much of the upper bay in low salinity areas (0 to 17 ‰). Young gobies, < 10.0mm TL, of the genus *Gobiosoma* were present in waters with salinities greater than 20 ‰ but could not be identified to species due to similar morphometric characteristics (Hildebrand and Cable, 1938). Fish, > 10.0mm TL, possessed distinguishing characteristics and were identified to species (Hildebrand and Cable, 1938; Massmann et al., 1963).

SUMMARY

The broad area of this investigation, upper Chesapeake Bay, is generally typical of the upper reaches of many Coastal Plain estuaries. The fauna is typically estuarine but should exhibit unique Chesapeake Bay characteristics. The following discussion attempts to present a synoptic view of the basic biological pattern of abundant early developmental stages of fishes found in upper Chesapeake Bay.

The biological information obtained during this investigation has been correlated with the estuarine salinity gradient, the temperature gradient, water depths, and seasons of oc-

currence in an attempt to ascertain life history patterns. An analysis of this information with special emphasis on the patterns of organism movement has made possible an evaluation of the nursery potential of the Chesapeake estuarine environment for fishes.

Salinity Correlations

Major emphasis has been directed toward correlations with the salinity gradient which serves as a yardstick to the estuary. Progressive changes in the location of an organism in relation to the salt content of the estuarine environment are assumed to indicate important migrations or movements, primarily between spawning and feeding areas with different salinities. The fishes represented here can be conveniently grouped in three main categories according to the salinity of the environment in which spawning takes place. These categories are freshwater, estuarine and marine.

Freshwater Spawners

The species listed here spawn close to the fresh-saltwater interface (striped bass), or many miles upstream (white and yellow perch and some of the herring groups). All possess yolk-sac food supplies and apparently do not depend on the nutritional capacities of their home tributaries for a short period following hatching. During this time larvae are planktonic and are carried downstream by freshwater runoff. Active feeding apparently starts for most of these fish as they approach low salinity waters. Larvae of freshwater fishes were abundant in spring in low salinity waters (< 13 o/oo) of the Patuxent and Magothy Rivers and upper Chesapeake Bay. This salinity range covers about 10 miles in the upper Patuxent River. Herman et al. (1968) state that the richest area of zooplankton in that river in spring is located near Trueman Point or about the middle of this 10-mile section. Numbers of adult zooplankton reach a maximum there in March and April or about the time larvae of freshwater fishes are hatching in the upper estuary. During May, zooplankton abundance declines due partly to feeding by large numbers of recently-hatched fishes and young-of-the-year fishes still on the nursery ground. Early developmental stages of several species collected during this investigation show no appreciable growth prior to the first of May but show accelerated growth from May to late summer. As summer approaches, numbers of zooplankton decrease and maturing fishes are less abundant in plankton samples collected in the low salinity area. By fall, many of the maturing fishes have moved toward the lower estuary.

In summary, developing fishes of the freshwater group hatch upstream and move downstream where they feed on phyto- and zooplankton produced in brackish water areas. Some fishes move farther downstream at the same time upper estuarine zooplankton abundance decreases and their food requirements change (e.g., striped bass and white perch).

Some fishes either remain in or return to the low salinity nursery area for several years.

Estuarine Spawners

Estuarine fishes spawn and mature in brackish areas. Some hatch and remain in the same area while others, like the bay anchovy and hogchoker, spawn over much of the estuarine salinity gradient but move, as larvae, to low salinity nursery grounds upstream in summer and fall. The hogchoker, *Trinectes maculatus* (Dovel et al., 1969), and the bay anchovy, *Anchoa mitchilli*, are ideal species for showing upstream migration of larvae, since larvae were abundant in areas where eggs were scarce or nonexistent. The spawning grounds for many of these estuarine species apparently extend into the marine environment.

Fishes of the oyster community, gobies, blennies, toadfish, and skillettfish, plus silversides, killifish, pipefish, sticklebacks, and winter flounder spawn over much of the middle estuary but show no apparent movement toward a different nursery area.

In general, the larvae of the estuarine group of fishes either remain in the area where they were hatched or move upstream to the vicinity of the salt-freshwater interface as critical early growth begins. Immature stages of some fishes of this group exhibit cyclic movement within the estuary during early development stages (e.g., hogchoker).

Marine Spawners

The marine category consists of fishes whose larvae hatch in the marine environment and show varying degrees of dependency upon the estuary as a nursery area. This group includes the menhaden, American eel, spot, weakfish, and Atlantic croaker which show strong upstream migrations. Most larvae of these species generally hatch over a several-month period and appear in Chesapeake estuaries during late summer to early winter. Migrations carry most of these fishes upstream to the vicinity of the salt-freshwater interface and some into fresh water. The larvae and juvenile stages of these fishes were not found in abundance in the lower Patuxent and adjacent Chesapeake Bay. It would appear that species which accumulate in the estuary over a period of several months are most abundant where these fishes congregate to feed. Larvae and juveniles of all species mentioned so far in this category were found congregated upstream in low salinity areas. In addition, four young Atlantic herring and four darter gobies were also found in the same area. This pattern would suggest that the sampling methods employed were more efficient in determining the presence of ichthyoplankton once congregated on the low salinity nursery grounds than while in transport to these areas.

Post larvae and juveniles of the seaboard goby, kingfish, feather blenny, blackcheek tonguefish, harvestfish, and

northern puffer were found in waters of intermediate salinities (11 to 22 o/oo). The absence of eggs of these marine spawners, with the possible exception of those of the kingfish, suggests upstream movement of larvae. The absence of larvae and juveniles from waters with salinities less than 11 o/oo would suggest that either the inshore movements of these forms were not pronounced, or that the presence of these forms was not sufficiently monitored by sampling gear.

A Common Estuarine Nursery

Early developmental stages of fishes from all three groups listed above utilize common low salinity feeding or nursery areas in the upper Chesapeake Bay. Each gradient or major tributary, which possesses a transition zone between fresh and salt waters, provides this nursery potential. These areas, as determined in this study, may be roughly described as water area with a 0 to 11 o/oo salinity range. Ninety-five percent of all fish larvae collected during this investigation were found in this salinity range (Table 27).

but before their full significance can be realized a critical comparison of maturation under both fresh and brackish water conditions must be made. The nursery area emphasized here is synonymous with the "critical zone" of estuaries described by Massman (1963) and referred to as having salinities of 1 to 15 or 5 to 15 o/oo (Massmann, 1964). There is probably no appreciable difference between the values of areas here designated as having salinities of 0 to 11, 1 to 15, and 5 to 15 o/oo. It seems probable that environmental differences, random movements of the organisms, and varying gear efficiency contribute to different salinity designations.

The "critical zones" are relatively small in Chesapeake tributaries; they must possess high turnover rates of food organisms to support large numbers of maturing fishes, especially during spring and summer months. Herman et al. (1968) describe the high production of copepods in the low salinity area of the Patuxent during spring. Heinle (1970) shows an inverse correlation between water temperature and the turnover time for the dominant copepods in




Figure 10. Schematic diagram of movements of estuarine-dependent fish larvae and juveniles toward a common low salinity nursery area. Numbers represent approximate salinity in parts per thousand (o/oo).

Movement

Critical Zone. Figure 10 depicts movement of developing fishes from different environments toward these common nursery grounds in the upper Chesapeake estuarine system. This figure suggests the importance of the low salinity estuarine environment as a nursery for fishes listed in this report. These brackish water nursery areas are obviously important,

that area as summer approaches. A turnover time of about one month during December and January decreases to about one day during midsummer.

Figure 11 emphasizes the specific and intricate movements of one important estuarine species, the hogchoker, in relation to the broad concept of larval and juvenile fish movements in the estuary.




Figure 11. Schematic diagram of movements of major groups of developing estuarine-dependent fishes with a breakdown of the inter-estuarine activities of a single dominant species, the hogchoker (*Trinectes maculatus*).

The young stages of most species collected during this study go through some variation of the cyclic movement pattern depicted for the hogchoker. An understanding of these complex movement patterns for estuarine-dependent fishes is essential before we can gain a working knowledge of appropriate ecological relationships.

Stratification. Correlations with depth show some stratification, either deliberate or accidental, but conducive to the movement of larvae and juveniles toward nursery areas, as described above. Much of the movement described here is passive rather than deliberate but is beneficial to the organism's development. Marine and estuarine fishes utilize a subsurface, net upstream water current to move upstream, while freshwater species utilize fresh water runoff to move downstream. The increasing ability to swim against water currents makes it impossible at this time to determine the extent to which fish determine their own route of travel.

Survival. There is evidence to suggest that downstream movement of some newly-hatched larvae from freshwater spawners is not only beneficial but necessary. Conte et al., (1966), showed that as juvenile *Oncorhynchus kisutch* develop, they exhibit an increasing ability to tolerate high salinity waters. In the case of the Coho salmon this increased tolerance precedes the downstream migration by about

6 months. Baggerman (1960) found that the fry of *Oncorhynchus keta* could not live for long periods in fresh water. Larvae hatched in fresh water in March would begin to die by June if restricted to that environment. Further restriction would result in the death of all fish by November. Larvae, when presented with a salinity choice situation, generally preferred fresh water at the end of the yolk-sac period but within about 4 weeks, there was a change to a salt water preference. It may be assumed that downstream movement of young fishes in the Chesapeake estuarine system is not entirely accidental but is correlated with changing salinity tolerances or preferences and has been beneficial to the evolution of those species.

Rapid and rhythmical. The movements of some estuarine-dependent fish larvae, whether passive or active, appear to be rapid and predictably rhythmical. Rapid movement suggests the need for frequent sampling in the estuary in order to accurately monitor the presence of nektonic animals. Some movement is apparently so rapid that even a weekly frequency is insufficient to follow movement along the gradient (Dovel et al., 1969). A weekly sampling frequency does, however, provide an indication of the general areas of organism concentration. A less frequent sampling schedule may allow organisms to move in and out of an area undetected.

Temperature Correlations

Correlations of egg and larvae presence with environmental water temperatures did not suggest the existence of distinct categories associated with different temperature ranges, although each species has an optimal temperature range for reproduction. Generally, some fish eggs and larvae were present in the estuary when water temperatures were between 3 and 31 C (Table 11). Ninety percent of all eggs collected were found with temperatures > 19 C. Temperatures > 27 C accounted for only 1.0%. An eight degree spread, 20-27 C, accounted for almost 90% of the total eggs collected. This reflects production of the two dominant estuarine fishes, the anchovy and hogchoker.

Eighty-three percent of all larvae and juveniles occurred with temperatures of 22 to 29 C. Most of these forms were larvae representing a few dominant estuarine species and were present during or just prior to the period of maximum summer water temperatures. Many juveniles were also present in low salinity nursery areas during periods of minimal water temperatures when a slight movement downstream would place these fish in warmer waters. Subjection of larvae and juveniles to the coldest possible water temperatures in the upper estuary during the first years of life is characteristic for many estuarine-dependent species. This temperature-growth relationship should be thoroughly investigated.

doubtedly correlated with a high growth potential in that environment. Although each species will follow a specific pattern of growth, it may be assumed that all estuarine-dependent forms gain nutritional benefits from the rich and productive coastal environment. Calculated patterns of growth will vary according to the abundance of different developmental stages, which in turn are affected by the characteristics and efficiency of collecting gear and sampling methods employed.

In consideration of inter-specific variations, the author planned to present individual growth curves for some species. A comprehensive examination of all raw data available suggested many problems in attempting to fit either a line or curve to the data in such a manner as to accurately portray the characteristics of early growth. It should be noted here that preliminary calculations produced line fits with high statistical correlation coefficients ($r = 0.8$). It is the author's opinion that while a high coefficient denotes good fit, this approach does not sufficiently convey the complex nature of the early growth of estuarine-dependent fishes.

In lieu of a single line fit, the author has developed a schematic diagram (Fig. 12) which will facilitate an understanding of larval and juvenile fish growth under estuarine conditions. It should be kept in mind that interpretations conveyed through this diagram have been derived from data collected with a one-meter net sampling approach (Dovel,




Figure 12. Schematic diagram of the generalized early growth pattern of estuarine-dependent fishes, derived from data collected with one-meter plankton nets.

Growth

This report documents the utilization of a common estuarine feeding area by fishes that originate in fresh water, the sea, and the estuary. Utilization of the estuary is un-

1964). The schematic was developed primarily from bay anchovy data but accurately conveys the general trend of growth for most species listed in this report.

The determination of an accurate pattern of growth for estuarine fishes requires the availability of at least some data

on several developmental stages from hatching until the organism approaches maturity.

The pattern of growth calculated from these stages could probably be depicted by a simple sigmoid curve were it not for changes in the environment, sampling efficiency, and the biological characteristics of the species which interrupt smooth growth.

Due to a short life span in the estuary, this critical period of growth prior to maturity, during which changes occur, will usually extend through the organism's first year to a year-and-one-half of growth. Data for this period are available for species such as the bay anchovy, white perch, yellow perch, northern pipefish, hogchoker, and several others. Fragmentary information is available for gobies, blennies, winter flounder, menhaden, and others listed in tables 2, 3, and 4. In some cases, large numbers of larvae were collected but represented only a small size-range or period of growth in the total life history (Tables 24, 25, and 26). In cases of the naked goby or winter flounder, for example, only larvae less than 10 mm were efficiently captured. In the case of menhaden, fish 25 to 60 mm were available. Regardless of the number of larvae available, an insufficient coverage of the total size-range of the immature stages in cases such as these precluded a determination of a growth pattern for these species.

The following discussion outlines, in a general way, the growth of the early developmental stages of fishes that are dependent upon factors associated with low salt content of estuarine waters.

The production of fish eggs and consequently larvae of dominant species in the estuary approximates a normal distribution (Figs. 2 and 12; Dovel, 1969, Fig. 2). This distribution is suggested in Fig. 12 by using a normally distributed frequency of a generalized sigmoid growth curve. Each line theoretically represents one individual of the population. Mortalities could be built into this diagram by terminating the lines or the growth of individuals at appropriate stages of development, if such points could be determined. Growth of early-hatched fishes accelerates rapidly so that by early fall, gear that efficiently collects eggs and small larvae no longer effectively monitors the population. This point of diminishing returns may be attributed in part to the organism's ability to evade collecting gear, mortality, and reduced availability due to emigration away from the critical nursery area. By mid-winter, when water temperatures have dropped almost to freezing, small, late-hatched organisms are present but not abundant in the nursery area. Larger individuals, with the exception of white perch (in this study) are absent and remain so until about April when the water temperature rises above 10 C. No appreciable growth occurs during cold months. This period is represented by the jagged break in the diagram (Fig. 12).

As water temperatures rise in the spring, the slower-growing fish which have remained close to the nursery are joined by larger individuals which were absent during the


cold months. The year-class now has a wide range of individual sizes. By August or late summer, faster-growing organisms of a new year-class attain a size comparable to the slow growers of the previous year-class. This period of size-overlap complicates the separation of year-classes on the basis of size and is attributed, in part, to prolonged spawning periods which result in varying rates of growth, as water temperatures fluctuate in the estuary. It may be assumed that if the organisms of the previous year-class were abundant during the period of size-overlap, and that both year-classes were sampled efficiently, the two age groups would show a definite bi-modal size distribution. Because of sampling limitations and the biological characteristics of estuarine fishes, members of a newly-hatched year-class are monitored far more accurately than older fish.

The developmental stages of many of the species listed in this report appear to be catchable in estuarine low-salinity areas during the first year to year-and-one-half of life. Their absence from samples after this time may be attributed to a combination of factors, including mortality, emigration, avoidance, and reduced gear efficiency. There is a point of no return in size for each species past which plankton sampling gear is not effective in sampling the population. This point may vary from 10 mm for the goby to something over 100 mm for several other species. For estuarine-dependent species that originate outside the estuary, growth patterns differ somewhat but suggest the same benefits from estuarine environmental factors conducive to maximum growth of early developmental stages of fishes.

It is obvious that the shape of a curve depicting periodic mean size of the population (Fig. 12, heavy black dots) will be substantially influenced by the distribution or pattern of occurrence of individuals. When a few larvae first appear in samples, their mean size (Fig. 12,A) may be slightly larger than the average size for newly-hatched forms. This pattern may result from the random chance of collecting representatives of the first larvae hatched, some of which may have hatched days before but did not appear in samples as a result of the sampling frequency used. As the bulk of new year-class begins to appear, the mean size may decrease slightly due to the influence of the concentration of newly-hatched (smallest possible) individuals. The calculation of a pattern of mean growth for estuarine fishes thus results from monitoring the peak activity of populations. With the onset of winter, the peak activity of a population may give way to the activity of the late hatch, or runts, which will reduce size estimates deflecting the upper end of the growth curve downward.

Fish eggs, recently-hatched larvae, or juveniles were present somewhere in the estuary at almost all times of the year. Species composition, relative abundance or density, distribution and movement patterns are constantly changing. Fig. 13 is symbolic of the complex activity that occurs as early developmental stages of estuarine-dependent fishes move through the estuary for brief intervals. It is here that some live out their lives entirely and many others, having

come from opposite directions, receive an important start in life.



W.L. Dovel '70

Figure 13. Stylized illustration of the movements of marine, freshwater and estuarine fishes through the critical nursery zone in an estuary.

LITERATURE CITED

- Baggermann, B. 1960. Salinity preference, thyroid activity and the seaward migration of four species of Pacific salmon. *Fish. Res. Bd. Can.* 17:296-322.
- Baxter, J. L. 1967. Introduction to symposium on anchovies, Genus *Engraulis*. *Calif. Corp. Oceanic Fish. Invest.* Vol. XI:28.
- Bayliff, W., Jr. 1950. The life history of the silverside, *Menidia menidia* (Linnaeus). *Chesapeake Biol. Lab. Publ.* No. 90.
- Biggs, R. B. 1967. The sediments of Chesapeake Bay estuaries. In *Estuaries*. George H. Lauff (Ed.), *Amer. Assoc. Advancement Sci. Publ.* No. 83.
- Burgess, R. H. 1965. Chesapeake circle. Cornell Maritime Press, Inc., Cambridge, Md.
- Carlson, F. T. 1968. Report on the biological findings. In Suitability of the Susquehanna River for Restoration of Shad. *U. S. Govt. Prtg. Off., Wash., D. C.* pp. 1-60.
- Coast and Geodetic Survey. 1967. Tidal current tables, Atlantic Coast of North America. *U. S. Dept. Int., Environm. Sci. Ser. Admn., Wash., D. C.* 199 p.
- Conte, F. P., H. H. Wagner, J. Fessler, and G. Grose. 1966. Development of osmotic and ionic regulation in juvenile Coho salmon, *Oncorhynchus kisutch*. *Comp. Biochem. Physiol.* 18, 1-15.
- Corps of Engineers. 1930. Patuxent River, Maryland. *U. S. Army House Doc. 463, 71st Congr., 2nd Sess.* 1-142.
- Dovel, W. L. 1963. Larval development of clingfish, *Gobiosox strumosus*, 4.0 to 12.0 millimeters total length. *Chesapeake Sci.* 4(4):161-6.
- . 1964. An approach to sampling estuarine macroplankton. *Chesapeake Sci.* 5(1-2):77-90.
- . 1967. Fish eggs and larvae of the Magothy River, Maryland. *Chesapeake Sci.* 8(2):125-29.
- . 1968. Predation by striped bass as a possible influence on population size of the Atlantic croaker. *Trans. Amer. Fish. Soc.* 97(4):313-9.
- , and J. R. Edmunds IV. 1971. Recent changes in striped bass (*Morone saxatilis*) spawning sites and commercial fishing areas in upper Chesapeake Bay. *Chesapeake Sci.* 12(1):33-39.
- , J. A. Mihursky, and A. J. McErlean. 1969. Life history aspects of the hogchoker, *Trinectes maculatus*, in the Patuxent River estuary, Maryland. *Chesapeake Sci.* 10(2):104-19.
- Gray, R. D. 1967. The national waterway, a history of the Chesapeake and Delaware Canal, 1769-1965. *U. Ill. Press, Library of Congr. Cat. Card* No. 67-12989
- Haight, F. J., H. E. Finnegan, and G. L. Anderson. 1930. Tides and currents in Chesapeake Bay and tributaries. *U. S. Dept. Comm. Spec. Publ.* No. 162:1-145.
- Heidel, S. G., and W. W. Frenier. 1965. Chemical quality of water and trace elements in the Patuxent River Basin. *Md. Geol. Surv. Rept. of Invest.* 1:1-40.
- Herman, S. S. 1958. The planktonic fish eggs and larvae of Narragansett Bay. M. S. Thesis, Univ. Rhode Island. 65 p.
- , J. A. Mihursky, and A. J. McErlean. 1968. Zooplankton and environment characteristics of the Patuxent River estuary, 1963-65. *Chesapeake Sci.* 9(2):67-82.
- Hildebrand, S. F. 1943. A review of the American anchovies (family Engraulidae). *Bing. Ocean. Coll. Bull.* 8(2):1-165.
- . 1963b. Family Engraulidae. In *Fishes of the Western North Atlantic*. *Sears Found. Mar. Res. Mem.* 1(3):152-249.
- , and Louella E. Cable. 1930. Development and life history of fourteen teleostean fishes at Beaufort, North Carolina. *Bull. U. S. Bur. Fish.* 46:383-488.
- . 1938. Further notes on the development and life history of some teleosts at Beaufort, North Carolina. *Bull. U. S. Bur. Fish.* 48(24):505-642.
- Hildebrand, S. F., and W. C. Schroeder. 1927. Fishes of Chesapeake Bay. *Bull. U. S. Bur. Fish.* XLIII, Part I:1-366.
- Hollis, E. H. 1952. Variations in the feeding habits of the striped bass, *Roccus saxatilis* (Walbaum), in the Chesapeake Bay. *Bull. Bing. Ocean. Coll.* 14(1):111-31.
- Kilby, J. D. 1955. The fishes of two Gulf coastal Marsh areas in Florida. *Tulane Stud. in Zool.* 2(8):175-247.
- Kuntz, A. 1914. The embryology and larval development of *Bairdiella chrysura* and *Anchovia mitchilli*. *Bull. U. S. Bur. Fish.* (1913) 33:3-19.
- McHugh, J. L. 1967. Estuarine nekton. *Amer. Ass. Adv. Sci. Publ.* 83:581-620.
- Mansueti, Alice J., and J. D. Hardy, Jr. 1967. Development of fishes of the Chesapeake Bay region, Part I. An atlas of egg, larval, and juvenile stages. Edited by E. E. Deubler, Jr. *Nat. Res. Inst., U. of Md.* 202 p.
- Mansueti, R. J. 1958. Eggs, larvae and young of the striped bass, *Roccus saxatilis*. *Md. Dept. Res. & Educ. Contr. No.* 112:35p.
- . 1961. Movements, reproduction, and mortality of the white perch, *Roccus americanus*, in the Patuxent estuary, Maryland. *Chesapeake Sci.* 2(3-4):142-205.
- . 1964. Eggs, larvae and young of the white perch, *Roccus americanus*, with comments on its ecology in the estuary. *Chesapeake Sci.* 5(1-2):3-45.
- , and H. Kolb. 1953. A historical review of the shad fisheries of North America. *Md. Dept. Res. & Educ. Publ.* 97:293 p.
- Massmann, W. H. 1954. Marine fishes in fresh and brackish waters of Virginia rivers. *Ecology* 35(1):75-8.
- . 1962. Water temperatures, salinities, and fishes collected during trawl surveys of Chesapeake Bay and York and Pamunkey Rivers, 1956-1959. *Va. Inst. Mar. Sci. Spec. Sci. Rept.* 27:1-3.
- . 1963. The "critical zone" in estuaries. *Bull. Sport Fish. Inst.* 141.
- . 1964. A positive approach to coastal sport fishery problems. *18th Proc., Oct. S. E. Ass. Game & Fish. Comm.*
- , J. J. Norcross, and E. B. Joseph. 1963. Distribution of larvae of the naked goby, *Gobiosoma boscii*, in the York River. *Chesapeake Sci.* 4(3):120-5.
- Nash, C. B. 1947. Environmental characteristics of a river estuary. *Md. Dept. Res. & Educ. Publ.* No. 64.
- Perlmutter, A. 1939. I. An ecological survey of a young fish and eggs identified from tow-net collections. In *A Biological Survey of the Salt Water of Long Island, 1938*. *Bull. N. Y. Cons. Dept., 28th Ann. Rept. Suppl.* (2):11-71.
- Pritchard, D. W. 1951. The physical hydrography of estuaries and some applications to biological problems. *Trans. 16th N. Am. Wildl. Conf. March*, 1951:368-76.
- . 1955. Estuarine circulation patterns. *Proc. Amer. Soc. Civ. Eng.* 81:Sep. No. 717.
- , and R. E. Bunce. 1959. Physical and chemical hydrography of the Magothy River. *Tech. Rept. XVII, Chesapeake Bay Inst., Johns Hopkins Univ., Ref.* 59-2.

- Reid, G. F., Jr. 1954. An ecological study of the Gulf of Mexico fishes in the vicinity of Cedar Keys, Florida. *Bull. Mar. Sci. Gulf and Carib.* 4(1):1-94.
- Richards, Sarah W. 1959. Oceanography of Long Island Sound. VI. Pelagic fish eggs and larvae of Long Island Sound. *Bull. Bing. Ocean. Coll.* 17(1):95-124.
- Runyan, Suzanne. 1961. Early development of the clingfish, *Gobiosox strumosus*, Cope. *Chesapeake Sci.* 2(3-4):113-41.
- Springer, V. G., and K. D. Woodburn. 1960. An ecological study of the fishes of the Tampa Bay area. *Fla. St. Bd. of Cons. Mar. Lab. Prof. Paper Ser.* 1:1-104.
- Stevenson, R. A., Jr. 1958. A biology of the anchovies, *Anchoa mitchilli mitchilli* Cuvier and Valenciennes, 1848, and *Anchoa hepsetus hepsetus*, 1758. M. S. Thesis, Univ. Dela. 56 p.
- Stross, R. G., and J. R. Stottlemeyer. 1965. Primary production in the Patuxent River. *Chesapeake Sci.* 6(3):125-40.
- Stroup, E. D., and R. J. Lynn. 1963. Atlas of salinity and temperature distributions in Chesapeake Bay 1952-61 and seasonal averages 1949-61. *Chesapeake Bay Inst., Johns Hopkins U., Graphic Summ. Rept.* 2, Ref. 63-1.
- Wells, R. C., R. K. Bailey, and E. P. Henderson. 1929. Salinity of the waters of Chesapeake Bay. *U. S. Geol. Surv. Prof. Paper Ser.* 154-C:105-52.
- Wheatland, Sarah D., 1956. Oceanography of Long Island Sound, 1952-54. VII. Pelagic fish eggs and larvae. *Bull. Bing. Ocean. Coll.* 15:234-314.
- Whitney, R. R. 1961. The Susquehanna fishery study, 1957-1960; A report of a study on the desirability and feasibility of passing fish at Conowingo Dam. *Md. Dept. Res. & Educ. Contr.* No. (169):1-81.

LIST OF TABLES

- Table 1. Fish egg and larvae sampling effort in the upper Chesapeake Bay (1960-1968).
- Table 2. List of common and scientific names of fishes appearing in this report.
- Table 3. Numbers of fish eggs collected at various locations in the upper Chesapeake Bay during 1963-67.
- Table 4. Numbers of fish larvae collected at various locations in the upper Chesapeake Bay during 1963-67.
- Table 5. Seasonal occurrence of fish eggs collected by meter nets in upper Chesapeake Bay (1963-67).
- Table 6. Seasonal occurrence of fish larvae and juveniles collected by meter nets in upper Chesapeake Bay (1963-67).
- Table 7. Distribution of the larvae of the bay anchovy, Anchoa mitchilli, by water salinity. Collections made in the Patuxent and Susquehanna River gradients (1963-67).
- Table 8. Distribution of fish eggs by water salinity in upper Chesapeake Bay (1963-67).
- Table 9. Distribution of fish larvae and juveniles by water salinity in upper Chesapeake Bay (1963-67).
- Table 10. Distribution of fish eggs by water temperature (°C) in upper Chesapeake Bay (1963-67).
- Table 11. Distribution of fish larvae and juveniles by water temperature (°C) in upper Chesapeake Bay (1963-67).
- Table 12. Species represented by larval stages in collections from upper Chesapeake Bay (1963-67).

- Table 13. Length frequencies of blueback herring, Alosa aestivalis, larvae collected in upper Chesapeake Bay.
- Table 14. Length frequencies of the white perch, Roccus americanus, larvae collected in upper Chesapeake Bay.
- Table 15. Monthly abundance of striped bass eggs from upper Chesapeake Bay, 1963-67.
- Table 16. Vertical distribution of the eggs of the striped bass (Morone saxatilis) by date and location of collection in upper Chesapeake Bay during 1966 and 1967.
- Table 17. Length frequencies of yellow perch, Perca flavescens, larvae collected from upper Chesapeake Bay during 1966 and 1967.
- Table 18. Monthly abundance of the eggs of the bay anchovy (Anchoa mitchilli) from the Chesapeake Bay area for the period 1963-67.
- Table 19. Relative sizes of bay anchovy eggs collected at different salinities.
- Table 20. Frequency distribution of the larvae of the bay anchovy, Anchoa mitchilli, by month of collection, water salinity, and vertical distribution in the Patuxent River during 1963.
- Table 21. Length frequency of bay anchovy larvae and juveniles collected in upper Chesapeake Bay.
- Table 22. Length frequencies of northern pipefish, Syngnathus fuscus, collected at Solomons, Maryland from 17 July 1961 to 25 July 1962.
- Table 23. Length frequencies of northern pipefish, Syngnathus fuscus, collected in upper Chesapeake Bay.
- Table 24. Length frequencies of naked goby, Gobiosoma bosci, larvae collected from the Patuxent River during 1963.
- Table 25. Length frequencies of winter flounder, Pseudopleuronectes americanus, larvae collected in upper Chesapeake Bay.

Table 26. Length frequencies of menhaden, Brevoortia tyrannus, collected in upper Chesapeake Bay, 1963-67.

Table 27. Cumulative percentages of fish larvae collected from different salinity ranges in Chesapeake Bay.

Table 1. Fish egg and larvae sampling effort in the upper Chesapeake Bay (1960-1968).

| Area Sampled | Date | Purpose |
|---------------------------------------|-----------|------------------------------|
| Mouth, Patuxent R. | 1960-1962 | Designing Field Techniques |
| " " " | 1961-1962 | Pilot Study, Familiarization |
| Patuxent River (Estuarine Portion) | 1963 | Weekly Sampling |
| " " | 1964 | Biweekly Sampling |
| " " | 1965 | " " |
| Magothy River | 1965-1966 | " " |
| Upper Chesapeake Bay | 1965-1968 | " " |

Table 2. List of Common and Scientific Names of Fishes Appearing in this Report.

| Common name | Scientific name |
|-------------------------|--------------------------------------|
| Blueback herring | <u>Alosa aestivalis</u> |
| Hickory shad | <u>Alosa mediocris</u> |
| Alewife | <u>Alosa pseudoharengus</u> |
| American shad | <u>Alosa sapidissima</u> |
| Atlantic menhaden | <u>Brevoortia tyrannus</u> |
| Atlantic herring | <u>Clupea harengus</u> |
| Bay anchovy | <u>Anchoa mitchilli</u> |
| Golden shiner | <u>Notemigonus crysoleucas</u> |
| Spottail shiner | <u>Notropis hudsonius</u> |
| White catfish | <u>Ictalurus catus</u> |
| Channel catfish | <u>Ictalurus punctatus</u> |
| American eel | <u>Anguilla rostrata</u> |
| Atlantic needlefish | <u>Strongylura marina</u> |
| Halfbeak | <u>Hyporhamphus unifasciatus</u> |
| Killifish | Family Cyprinodontidae |
| Rainwater killifish | <u>Lucania parva</u> |
| Fourspine stickleback | <u>Apeltes quadracus</u> |
| Threespine stickleback | <u>Gasterosteus aculeatus</u> |
| Spotted seahorse | <u>Hippocampus erectus</u> |
| Northern pipefish | <u>Syngnathus fuscus</u> |
| White perch | <u>Morone americana</u> |
| Striped bass | <u>Morone saxatilis</u> |
| Pumpkinseed | <u>Lepomis gibbosus</u> |
| Johnny darter | <u>Etheostoma nigrum</u> |
| Yellow perch | <u>Perca flavescens</u> |
| Silver perch | <u>Bairdiella chrysura</u> |
| Weakfish | <u>Cynoscion regalis</u> |
| Spot | <u>Leiostomus xanthurus</u> |
| Southern kingfish | <u>Menticirrhus americanus</u> |
| Atlantic croaker | <u>Micropogon undulatus</u> |
| Red drum | <u>Sciaenops ocellata</u> |
| Darter goby | <u>Gobionellus boleosoma</u> |
| Naked goby | <u>Gobiosoma bosci</u> |
| Seaboard goby | <u>Gobiosoma ginsburgi</u> |
| Green goby | <u>Microgobius thalassinus</u> |
| Striped blenny | <u>Chasmodes bosquianus</u> |
| Feather blenny | <u>Hypsoblennius hentzi</u> |
| Southern harvestfish | <u>Peprilus alepidotus</u> |
| Rough silverside | <u>Membras martinica</u> |
| Tidewater silverside | <u>Menidia beryllina</u> |
| Atlantic silverside | <u>Menidia menidia</u> |
| Winter flounder | <u>Pseudopleuronectes americanus</u> |
| Hogchoker | <u>Trinectes maculatus</u> |
| Blackcheek tonguefish | <u>Symphurus plagiosa</u> |
| Skilletfish (clingfish) | <u>Gobiesox strumosus</u> |
| Northern puffer | <u>Sphaeroides maculatus</u> |
| Striped burrfish | <u>Chilomycterus schoepfi</u> |
| Oyster toadfish | <u>Opsanus tau</u> |

Table 3. Numbers of fish eggs collected at various locations in the upper Chesapeake Bay during 1963-67.

| Species | Patuxent River | | | Magothy River | Upper Bay | | Total |
|---|----------------|------|------|---------------|-----------|------|-----------|
| | 1963 | 1964 | 1965 | 1965 | 1966 | 1967 | |
| <u>Fresh</u> | | | | | | | |
| Blueback herring <u>Alosa aestivalis</u> | | | | x | x | x | 4,394 |
| Hickory shad <u>Alosa mediocris</u> | | | | x | | | 1 |
| Alewife <u>Alosa pseudoharengus</u> | x | | | | x | x | 1,933 |
| American shad <u>Alosa sapidissima</u> | | | | | x | x | 308 |
| Clupeidae sp. | x | | | | x | x | 39,915 |
| White perch <u>Roccus americanus</u> | x | x | x | x | x | x | 59,736 |
| Striped bass <u>Roccus saxatilis</u> | x | x | x | | x | x | 38,643 |
| <u>Brackish</u> | | | | | | | |
| Bay anchovy <u>Anchoa mitchilli</u> | x | x | x | x | x | x | 3,310,258 |
| Winter flounder <u>Pseudopleuronectes americanus</u> | | | | | | x | 1 |
| Hogchoker <u>Trinectes maculatus</u> | x | x | x | x | x | x | 411,954 |
| <u>Marine</u> | | | | | | | |
| Atlantic menhaden <u>Brevoortia tyrannus</u> | x | x | x | | x | x | 9,942 |
| Southern kingfish <u>Menticirrhus americanus</u> | | | | | x | x | 2,393 |
| | | | | | Total | | 3,879,477 |

Table 4. Numbers of fish larvae collected at various locations in the upper Chesapeake Bay during 1963-67.

| Species | Patuxent River | | Magothy River | | Bay | Upper Bay | | Total |
|---|----------------|------|---------------|------|------|-----------|------|---------|
| | 1963 | 1964 | 1965 | 1965 | 1965 | 1966 | 1967 | |
| <u>Fresh</u> | | | | | | | | |
| Blueback herring <u>Alosa aestivalis</u> | x | | | x | | | x) | 17,337 |
| Alewife | x | x | | x | | | x) | |
| <u>Alosa pseudoharengus</u> | | | | | | |) | |
| American shad <u>Alosa sapidissima</u> | | | | | | x | | 236 |
| Clupeidae sp. | x | x | x | x | | x | x | 78,919 |
| Golden shiner | x | | | | | | | 2 |
| <u>Notemigonus crysoleucas</u> | | | | | | | | |
| Spottail shiner | x | x | | | | | | 32 |
| <u>Notropis hudsonius</u> | | | | | | | | |
| White catfish <u>Ictalurus catus</u> | | | | | | | x | 12 |
| Channel catfish <u>Ictalurus punctatus</u> | | | | | | x | x | 139 |
| Atlantic needlesifh <u>Strongylura marina</u> | x | | | x | | x | x | 19 |
| Halfbeak <u>Hyporhamphus unifasciatus</u> | | x | x | | | x | | 25 |
| White perch <u>Roccus americanus</u> | x | x | x | x | | x | x | 51,275 |
| Striped bass <u>Roccus saxatilis</u> | x | x | x | | | x | x | 10,022 |
| Pumpkinseed <u>Lepomis gibbosus</u> | | | | x | | | x | 5 |
| Johnny darter <u>Etheostoma nigrum</u> | | | | | | x | x | 80 |
| Yellow perch <u>Perca flavescens</u> | x | x | x | | | x | x | 23,634 |
| <u>Brackish</u> | | | | | | | | |
| Bay anchovy <u>Anchoa mitchilli</u> | x | x | x | x | | x | x | 239,116 |
| Fourspine stickleback <u>Apeltes quadracus</u> | x | x | | | | | | 30 |
| Threespine stickleback <u>Gasterosteus aculeatus</u> | | | | | | | x | 1 |
| Northern pipefish <u>Syngnathus fuscus</u> | x | x | x | x | | x | x | 529 |
| Naked goby <u>Gobiosoma bosci</u> | x | x | x | x | | x | x | 700,222 |
| Green goby <u>Microgobius thalassinus</u> | | | x | x | | x | x | 135 |
| Striped blenny <u>Chasmodes bosquianus</u> | x | x | x | x | | | | 1,008 |

Table 4. (Continued)

| Species | Patuxent River | | | Magothy River | Bay | Upper Bay | Total | |
|---|----------------|------|------|---------------|------|-----------|-------|-----------|
| | 1963 | 1964 | 1965 | 1965 | 1965 | 1966 | | 1967 |
| Rough silverside <u>Membras martinica</u> | x | x | x | | | x | x | 884 |
| Tidewater silverside <u>Menidia beryllina</u> | x | x | | | | | x | 1,019 |
| Atlantic silverside <u>Menidia menidia</u> | x | x | x | | | | x | 2,345 |
| <u>Menidia sp.</u> | x | x | x | x | | x | x | 138,325 |
| Winter flounder <u>Pseudopleuronectes americanus</u> | x | x | x | x | | x | x | 14,303 |
| Hogchoker <u>Trinectes maculatus</u> | x | x | x | x | | x | x | 2,374 |
| Skilletfish (clingfish) <u>Gobiesox strumosus</u> | x | x | x | x | | | | 683 |
| Oyster toadfish <u>Opsanus tau</u> | x | x | x | | | x | | 9 |
| <u>Marine</u> | | | | | | | | |
| Atlantic menhaden <u>Brevoortia tyrannus</u> | x | x | | x | | x | | 2,322 |
| Atlantic herring <u>Clupea harengus harengus</u> | x | | | | | | | 4 |
| American eel <u>Anquilla rostrata</u> | x | | | | | x | x | 160 |
| Spotted seahorse <u>Hippocampus erectus</u> | x | | | | | | | 3 |
| Silver perch <u>Bairdiella chrysura</u> | | | | | | | x | 8 |
| Weakfish <u>Cynoscion regalis</u> | x | | | | | x | x | 368 |
| Spot <u>Leiostomus xanthurus</u> | x | | | | x | x | | 41 |
| Southern kingfish <u>Menticirrhus americanus</u> | x | | | | x | x | | 3 |
| Atlantic croaker <u>Micropogon undulatus</u> | x | x | x | | x | x | x | 468 |
| Red drum <u>Sciaenops ocellata</u> | x | | | | | | | 1 |
| Darter goby <u>Gobionellus boleosoma</u> | | x | | | | | | 4 |
| Seaboard goby <u>Gobiosoma ginsburgi</u> | x | | | | x | x | x | 166 |
| Feather blenny <u>Hypsoblennius hentzi</u> | | | | | x | | | 42 |
| Southern harvestfish <u>Peprilus alepidotus</u> | x | x | | | | | | 101 |
| Blackcheek tonguefish <u>Symphurus plagiusa</u> | | | | | x | | | 10 |
| Northern puffer <u>Sphaeroides maculatus</u> | x | | | | | | | 34 |
| | | | | | | | | 1,286,455 |

Table 5. Seasonal occurrence of fish eggs collected by meter nets in upper Chesapeake Bay (1963-67).

| Species | Monthly Occurrence | | | | | | | | | | | |
|--------------------------|--------------------|------|------|------|-----|------|------|------|-------|------|------|------|
| | Jan. | Feb. | Mar. | Apr. | May | June | July | Aug. | Sept. | Oct. | Nov. | Dec. |
| <u>Fresh</u> | | | | | | | | | | | | |
| <u>A. aestivalis</u> | | | | — | | | | | | | | |
| <u>A. mediocris</u> | | | | — | — | | | | | | | |
| <u>A. pseudoharengus</u> | | | — | — | — | | | | | | | |
| <u>A. sapidissima</u> | | | | — | — | | | | | | | |
| <u>Clupeidae sp.</u> | | | | — | — | — | | | | | | |
| <u>R. americanus</u> | | | — | — | — | | | | | | | |
| <u>M. saxatilis</u> | | | | — | — | — | | | | | | |
| <u>Brackish</u> | | | | | | | | | | | | |
| <u>A. mitchilli</u> | | | | | — | — | — | — | — | | | |
| <u>P. americanus</u> | | | ■ | | | | | | | | | |
| <u>T. maculatus</u> | | | | | — | — | — | — | | | | |
| <u>Marine</u> | | | | | | | | | | | | |
| <u>B. tyrannus</u> | | | | | — | — | | | | | — | |
| <u>M. americanus</u> | | | | | | — | | | | | | |

Table 6. Seasonal occurrence of fish larvae and juveniles collected by meter nets in upper Chesapeake Bay (1963-67). Bars represent recorded occurrence during organism's first year of growth.

| Species | Recently hatched | | | | | | | | | | | | | |
|------------------------|------------------|------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|--|--|
| | Jan. | Feb. | Mar. | Apr. | May | June | July | Aug. | Sept. | Oct. | Nov. | Dec. | | |
| <u>Fresh</u> | | | | | | | | | | | | | | |
| Clupeidae sp. | | | | ————— | | | | | | | | | | |
| <u>N. cryoleucas</u> | | | | ————— | | | | | | | | | | |
| <u>N. hudsonius</u> | | | | ————— | | | | | | | | | | |
| <u>I. catus</u> | | | | | | | ■ | | | | ■ | | | |
| <u>I. punctatus</u> | | | | | | | ————— | | | ————— | | | | |
| <u>S. marina</u> | | | | | | ————— | | | | | | | | |
| <u>H. unifasciatus</u> | | | | | ————— | | | | | | | | | |
| <u>R. americanus</u> | | | | ————— | | | | | | | | | | |
| <u>M. saxatilis</u> | | | | ————— | | | | | | | | | | |
| <u>L. gibbosus</u> | | | | ■ | | | ■ | | | | | | | |
| <u>E. nigrum</u> | | | | | | | ————— | | | | ————— | | | |
| <u>P. flavescens</u> | | | | ————— | | | | | | | | | | |
| <u>Brackish</u> | | | | | | | | | | | | | | |
| <u>A. mitchilli</u> | | | | | ————— | | | | | | | | | |
| <u>A. quadracus</u> | | | | ————— | | | | | | | | | | |
| <u>G. aculeatus</u> | | | ■ | | | | | | | | | | | |
| <u>S. fuscus</u> | | | | | ————— | | | | | | | | | |
| <u>G. bosci</u> | | | | | ————— | | | | | | | | | |
| <u>M. thalassinus</u> | | | | | ————— | | | | | | | | | |
| <u>C. bosquianus</u> | | | | | ————— | | | | | | | | | |
| <u>M. martinica</u> | | | | | ————— | | | | | | | | | |
| <u>M. beryllina</u> | | | | | ————— | | | | | | | | | |
| <u>M. menidia</u> | | | | | ————— | | | | | | | | | |
| Menidia sp. | | | | ————— | | | | | | | | | | |
| <u>P. americanus</u> | | | ————— | | | | | | | | | | | |
| <u>T. maculatus</u> | | | | ————— | | | | | | | | | | |
| <u>G. strumosus</u> | | | | ————— | | | | | | | | | | |
| <u>O. tau</u> | | | | | | | | ————— | | | | ————— | | |
| <u>Marine</u> | | | | | | | | | | | | | | |
| <u>B. tyrannus</u> | | | | | ————— | | | | | | | | | |
| <u>C. harengus</u> | | | | | | | | | | | ■ | | | |
| <u> harengus</u> | | | | | | | | | | | | | | |
| <u>A. rostrata</u> | | | | | | | | | | | | | | |
| <u>H. erectus</u> | | | | | | | ■ | | ————— | | | | | |
| <u>B. chrysur</u> | | | | | | | | | | ■ | | | | |
| <u>C. regalis</u> | | | | | ■ | | ————— | | | | | | | |
| <u>L. xanthurus</u> | | | | | ————— | | | | | | | | | |
| <u>M. americanus</u> | | | | | ■ | | | | | ■ | | | | |
| <u>M. undulatus</u> | | | | | | | | | ————— | | | | | |
| <u>S. ocellata</u> | | | | | | | | | ————— | | | | | |
| <u>G. boleosoma</u> | | | | ■ | | | | | | ■ | | | | |
| <u>G. ginsburgi</u> | | | | | ————— | | | | | | | | | |
| <u>H. hentzi</u> | | | | | ————— | | | | | | | | | |
| <u>P. alepidotus</u> | | | | | | | ■ | | | ■ | | ■ | | |
| <u>S. plagiusa</u> | | | | | | | | | ————— | | | | | |
| <u>S. maculatus</u> | | | | | ————— | | | | | | | | | |
| <u>C. schoepfi</u> | | | | | ————— | | | | | | | | | |

Table 6. Seasonal occurrence of fish larvae and juveniles collected by meter nets in upper Chesapeake Bay (1963-67). Bars represent recorded occurrence during organism's first year of growth.

| Species | Jan. | Feb. | Mar. | Apr. | May | June | July | Aug. | Sept. | Oct. | Nov. | Dec. |
|------------------------|------|------|------|------|-----|------|------|------|-------|------|------|------|
| <u>Fresh</u> | | | | | | | | | | | | |
| <u>Clupeidae sp.</u> | | | | — | | | | | | | | |
| <u>N. cryoleucas</u> | | | | | | ■ | | | | | | |
| <u>N. hudsonius</u> | | — | — | | | ■ | | | | | | |
| <u>I. catus</u> | | | | | | | | | | | | |
| <u>I. punctatus</u> | — | | | — | | | | | | | | |
| <u>S. marina</u> | — | | | | | | | | | | | |
| <u>H. unifasciatus</u> | | | | | | | | | | | | |
| <u>R. americanus</u> | — | | — | | | | | | | | | |
| <u>M. saxatilis</u> | — | | — | | | | | | | | | |
| <u>L. gibbosus</u> | — | | | | | | | | | | | |
| <u>E. nigrum</u> | — | | — | | | | | | | | | |
| <u>P. flavescens</u> | ■ | | ■ | | | | | | | | | |
| <u>Brackish</u> | | | | | | | | | | | | |
| <u>A. mitchilli</u> | — | | — | — | — | | | | | | | |
| <u>A. quadracus</u> | | — | — | | | | | | | | | |
| <u>G. aculeatus</u> | | | | | | | | | | | | |
| <u>S. fuscus</u> | — | — | — | — | — | | | | | | | |
| <u>G. bosci</u> | — | — | — | — | — | | | | | | | |
| <u>M. thalassinus</u> | | | | — | — | | | | | | | |
| <u>C. bosquianus</u> | | | | | | | | | | | | |
| <u>M. martinica</u> | | | | | — | — | — | | | | | |
| <u>M. beryllina</u> | | | | — | — | — | — | | | | | |
| <u>M. menidia</u> | | ■ | | — | — | | | | | | | |
| <u>Menidia sp.</u> | | | — | — | — | | | | | | | |
| <u>P. americanus</u> | | | — | — | — | | | | | | | |
| <u>T. maculatus</u> | ■ | | — | — | — | — | — | | | | | |
| <u>G. strumosus</u> | | | | | | | | | | | | |
| <u>O. tau</u> | | | | | | ■ | | | | | | |
| <u>Marine</u> | | | | | | | | | | | | |
| <u>B. tyrannus</u> | | | | — | — | | | | | | | |
| <u>C. harengus</u> | | | | | | | | | | | | |
| <u> harengus</u> | | | — | | | | | | | | | |
| <u>A. rostrata</u> | | — | — | — | — | — | — | — | — | | — | — |
| <u>H. erectus</u> | | — | — | — | — | — | — | — | — | | | |
| <u>B. chrysur</u> | | | | | | | | | | | | |
| <u>C. regalis</u> | | | | | | | | | | | | |
| <u>L. xanthurus</u> | | | | | | | | | | | | |
| <u>M. americanus</u> | | | | | | | | | | | | |
| <u>M. undulatus</u> | — | | | | | | | | | | | |
| <u>S. ocellata</u> | | | | | | | | | | | | |
| <u>G. boleosoma</u> | | | | | | | | | | | | |
| <u>G. ginsburgi</u> | | | | | | | | | | | | |
| <u>H. hentzi</u> | | | | | | | | | | | | |
| <u>P. alepidotus</u> | | | | | | | | | | | | |
| <u>S. plagiusa</u> | | | | | | | | | | | | |
| <u>S. maculatus</u> | | | | | | | | | | | | |
| <u>G. schoepfi</u> | | | | | | | | | | | | |

Table 7. Distribution of larvae of the bay anchovy, *Anchoa mitchilli*, by water salinity. Collections made in the Patuxent and Susquehanna River gradients - 1963-67.

| Sal. o/oo | Patuxent River | | | | | Upper Chesapeake Bay | | | | Grand | |
|--------------|----------------|-------|-------|---------|------|----------------------|-------|--------|------|---------|------|
| | 1963 | 1964 | 1965 | Totals | % | 1966 | 1967 | Totals | % | Total | % |
| 23 | 26 | | | 26 | + | | | | | 26 | + |
| 22 | | | | | | | | | | | |
| 21 | 156 | | | 156 | + | | | | | 156 | + |
| 20 | 971 | | 2 | 973 | 0.5 | | | | | 973 | 0.5 |
| 19 | 48 | 2 | | 50 | + | | | | | 50 | + |
| 18 | 66 | | | 66 | + | | | | | 66 | + |
| 17 | 18 | 1 | 1 | 20 | + | | | | | 20 | + |
| 16 | 210 | 3 | 3 | 216 | 0.1 | 27 | 4 | 31 | 0.2 | 247 | 0.1 |
| 15 | 1,744 | 2 | 49 | 1,795 | 1.0 | 8 | 2 | 10 | + | 1,805 | 0.9 |
| 14 | 1,201 | 24 | 231 | 1,456 | 0.8 | 46 | | 46 | 0.4 | 1,502 | 0.8 |
| 13 | 2,787 | 94 | 76 | 2,957 | 1.6 | 25 | | 25 | 0.2 | 2,982 | 1.5 |
| 12 | 8,531 | 97 | 214 | 8,842 | 4.8 | 117 | 4 | 121 | 1.0 | 8,963 | 4.6 |
| 11 | 11,997 | 186 | 298 | 12,481 | 6.8 | 23 | 298 | 321 | 2.7 | 12,802 | 6.6 |
| 10 | 6,613 | 176 | 498 | 7,287 | 4.0 | 15 | 46 | 61 | 0.5 | 7,348 | 3.8 |
| 9 | 479 | 18 | 46 | 543 | 0.3 | 13 | 24 | 37 | 0.3 | 480 | 0.3 |
| 8 | 5,706 | 805 | 472 | 6,983 | 3.8 | 24 | 20 | 44 | 0.4 | 7,027 | 3.6 |
| 7 | 24,134 | 214 | 1,472 | 25,820 | 14.1 | 74 | 13 | 87 | 0.7 | 25,907 | 13.3 |
| 6 | 38,950 | 64 | 684 | 39,698 | 21.7 | 78 | 1,209 | 1,287 | 10.8 | 40,985 | 21.0 |
| 5 | 14,082 | 260 | 586 | 14,928 | 8.2 | 14 | 51 | 65 | 0.5 | 14,993 | 7.7 |
| 4 | 19,776 | 856 | 2,234 | 22,866 | 12.5 | 421 | 5,232 | 5,653 | 47.6 | 28,519 | 14.6 |
| 3 | 26,536 | 716 | 1,628 | 28,880 | 15.8 | 375 | 714 | 1,089 | 9.2 | 29,969 | 15.4 |
| 2 | 5,080 | 92 | 314 | 5,486 | 3.0 | 389 | 467 | 856 | 7.2 | 6,432 | 3.3 |
| 1 | 1,224 | 12 | 14 | 1,250 | 0.7 | 229 | 22 | 251 | 2.1 | 1,501 | 0.8 |
| 0 | 152 | 4 | 86 | 242 | 0.1 | 32 | 1,857 | 1,889 | 15.9 | 2,131 | 1.1 |
| Total | 170,487 | 3,626 | 8,908 | 183,021 | 99.8 | 1,910 | 9,963 | 11,873 | 99.8 | 194,896 | 99.9 |

Table 8. Distribution of fish eggs by water salinity in upper Chesapeake Bay (1963-67). Numbers represent total number of eggs collected from various locations during several years.

| Salinity o/oo | Species | | | |
|------------------|--------------------------------|-------------------------------|------------------------------------|---------------------------------|
| | <u>A.</u> <u>aestivalis</u> | <u>A.</u> <u>mediocris</u> | <u>A.</u> <u>pseudoharengus</u> | <u>A.</u> <u>sapidissima</u> |
| 23 | | | | |
| 22 | | | | |
| 21 | | | | |
| 20 | | | | |
| 19 | | | | |
| 18 | | | | |
| 17 | | | | |
| 16 | | | | |
| 15 | | | | |
| 14 | | | | |
| 13 | | | | |
| 12 | | | | |
| 11 | | | | |
| 10 | | | | |
| 9 | | | | |
| 8 | | | | |
| 7 | | 1 | | |
| 6 | | | | |
| 5 | | | | |
| 4 | | | | |
| 3 | | | 2 | |
| 2 | 16 | | 6 | |
| 1 | | | | 152 |
| 0 | 3,089 | | 1,010 | 156 |
| Total | 3,105 | 1 | 1,018 | 308 |

Table 8 . Distribution of fish eggs by water salinity in upper Chesapeake (cont.) Bay (1963-67). Numbers represent total number of eggs collected from various locations during several years.

| Salinity o/oo | Species | | | | |
|------------------|--------------------|------------------|---------------------|----------------------|---------------------|
| | <u>B. tyrannus</u> | <u>Clupeidae</u> | <u>A. mitchilli</u> | <u>R. americanus</u> | <u>R. saxatilis</u> |
| 23 | | | | | |
| 22 | 3,968 | | 1,773 | | |
| 21 | | | 92,783 | | |
| 20 | 1,792 | | 22,720 | | |
| 19 | 13 | | 211,096 | | |
| 18 | 322 | | 103,955 | | |
| 17 | 9 | | 52,288 | | |
| 16 | 27 | | 48,055 | | |
| 15 | 70 | | 340,794 | | |
| 14 | 139 | | 356,654 | | |
| 13 | 81 | | 449,826 | 1 | |
| 12 | 56 | | 216,909 | | 6 |
| 11 | | | 132,788 | 30 | |
| 10 | 24 | | 15,724 | | 18 |
| 9 | | | 114,979 | 8 | |
| 8 | | | 23,894 | | 12 |
| 7 | | | 35,811 | 28 | 22 |
| 6 | | | 151,836 | 31 | 10 |
| 5 | | 3 | 6,288 | 10 | 289 |
| 4 | | | 28,320 | | 105 |
| 3 | | 2 | 133 | 16 | 2,635 |
| 2 | | 6,617 | 1,459 | 83 | 4,521 |
| 1 | | 2,651 | 56 | 401 | 8,358 |
| 0 | | 29,884 | | 20,110 | 20,612 |
| Total | 6,501 | 39,157 | 2,408,141 | 20,718 | 36,588 |

Table 8. Distribution of fish eggs by water salinity in upper Chesapeake Bay (1963-67). Numbers represent total number of eggs collected from various locations during several years.

| Salinity o/oo | Species | | | Total |
|------------------|----------------------|----------------------|---------------------|-----------|
| | <u>M. americanus</u> | <u>P. americanus</u> | <u>T. maculatus</u> | |
| 23 | | | 388 | 2,158 |
| 22 | | | 155 | 5,896 |
| 21 | | | 53,050 | 145,833 |
| 20 | | 1 | 7,986 | 32,499 |
| 19 | 358 | | 15,959 | 227,426 |
| 18 | 269 | | 90,378 | 194,924 |
| 17 | | | 43,315 | 95,612 |
| 16 | 117 | | 47,261 | 95,460 |
| 15 | 247 | | 47,912 | 389,023 |
| 14 | | | 46,686 | 403,479 |
| 13 | | | 22,979 | 472,887 |
| 12 | | | 17,797 | 234,768 |
| 11 | | | 1,793 | 134,611 |
| 10 | | | 10 | 15,776 |
| 9 | | | 24 | 115,011 |
| 8 | | | 40 | 23,946 |
| 7 | | | 9 | 35,871 |
| 6 | | | | 151,877 |
| 5 | | | | 6,590 |
| 4 | | | | 28,425 |
| 3 | | | 4 | 2,792 |
| 2 | | | | 12,702 |
| 1 | | | | 11,618 |
| 0 | | | | 74,861 |
| Total | 991 | 1 | 395,746 | 2,914,045 |

Table 9. Distribution of fish larvae and juveniles by water salinity in upper Chesapeake Bay (1963-67). Numbers represent total number of fish collected at various locations during several years.

| Salinity o/oo | <u>Alosa</u> <u>aestivalis</u> | <u>A. pseudo-</u> <u>harengus</u> | <u>A. sapi-</u> <u>dissima</u> | <u>Brevoortia</u> <u>tyrannus</u> | <u>Clupea</u> <u>harengus</u> | <u>Anchoa</u> <u>mitchilli</u> | <u>Notemigonus</u> <u>crysoleucas</u> |
|------------------|-----------------------------------|--------------------------------------|-----------------------------------|--------------------------------------|----------------------------------|-----------------------------------|--|
| 23 | | | | | | 26 | |
| 22 | 1 | | | | | | |
| 21 | | | | | | 156 | |
| 20 | | | | | | 971 | |
| 19 | | | | | | 50 | |
| 18 | | | | | | 66 | |
| 17 | | | | | | 20 | |
| 16 | | | | | | 247 | |
| 15 | | | | | | 1,805 | |
| 14 | | | | | | 1,502 | |
| 13 | | | | | | 2,982 | |
| 12 | | | | | | 8,963 | |
| 11 | 98 | | | 4 | 2 | 12,802 | |
| 10 | | | | | 2 | 7,348 | |
| 9 | | | | 10 | | 580 | |
| 8 | | 2 | | 68 | | 7,027 | |
| 7 | 2 | | | 44 | | 25,907 | |
| 6 | 14 | | | 42 | | 40,985 | |
| 5 | | 2 | | 336 | | 14,993 | |
| 4 | 2 | 4 | | 70 | | 28,519 | |
| 3 | 42 | 60 | | 334 | | 29,969 | 2 |
| 2 | 16 | 25 | | 124 | | 6,342 | |
| 1 | 2 | 48 | | 1,090 | | 1,501 | |
| 0 | 3,089 | 662 | 88 | 173 | | 2,131 | |
| Total | 3,266 | 803 | 88 | 2,295 | 4 | 194,892 | 2 |

Table 9 . Distribution of fish larvae and juveniles by water salinity in
 (Cont.) upper Chesapeake Bay (1963-67). Numbers represent total number
 of fish collected at various locations during several years.

| Salinity | <u>Etheostoma</u> <u>nigrum</u> | <u>Perca</u> <u>flavescens</u> | <u>Bairdiella</u> <u>chrysur</u> | <u>Cynoscion</u> <u>regalis</u> | <u>Leiostomus</u> <u>xanthurus</u> | <u>Menticirrhus</u> <u>americanus</u> | <u>M.</u> <u>undulatus</u> |
|----------|------------------------------------|-----------------------------------|-------------------------------------|------------------------------------|---------------------------------------|--|-------------------------------|
| 23 | | | | | | | |
| 22 | | | | 5 | | | |
| 21 | | | | 9 | | | 5 |
| 20 | | | | | | | |
| 19 | | | | | | 3 | 3 |
| 18 | | | | | | | 5 |
| 17 | | | | 1 | | | |
| 16 | | | | 1 | | | 38 |
| 15 | | | | 1 | | | 17 |
| 14 | | | | | 2 | | 4 |
| 13 | | 24 | | 15 | 1 | | 50 |
| 12 | | | | | 1 | | 23 |
| 11 | | 43 | | | | | 14 |
| 10 | | 44 | | | | | 6 |
| 9 | | 59 | | | | | 18 |
| 8 | | 9 | | | 14 | | 44 |
| 7 | 24 | 79 | | 2 | 4 | | 45 |
| 6 | | 38 | | 6 | | | 19 |
| 5 | 29 | 200 | 1 | 1 | | | 16 |
| 4 | 20 | 29 | 1 | | | | 23 |
| 3 | | 355 | | 2 | | | 2 |
| 2 | | 942 | 6 | | 2 | | 1 |
| 1 | | 1,041 | | | 13 | | 11 |
| 0 | 2 | 9,571 | | | 4 | | 1 |
| Total | 75 | 12,434 | 8 | 43 | 41 | 3 | 345 |

Table 9. Distribution of fish larvae and juveniles by water salinity in upper Chesapeake Bay (1963-67). Numbers represent total number of fish collected at various locations during several years.

| Salinity o/oo | <u>Notropis</u> <u>hudsonius</u> | <u>Ictalurus</u> <u>catus</u> | <u>Ictalurus</u> <u>punctatus</u> | <u>Anguilla</u> <u>rostrata</u> | <u>Strongylura</u> <u>marina</u> | <u>H. unifas-</u> <u>ciatus</u> | <u>Apeltes</u> <u>quadracus</u> |
|------------------|-------------------------------------|----------------------------------|--------------------------------------|------------------------------------|-------------------------------------|------------------------------------|------------------------------------|
| 23 | | | | 1 | | | |
| 22 | | | | 5 | | | 1 |
| 21 | | | | | | | |
| 20 | | | | 5 | | | |
| 19 | | | | 4 | | | |
| 18 | | | | | | | 3 |
| 17 | | | | | | | 5 |
| 16 | | | | | | | 1 |
| 15 | | | | 1 | | | 5 |
| 14 | | | | 3 | 1 | | 2 |
| 13 | | | | | | | |
| 12 | | | | 4 | 3 | 1 | 3 |
| 11 | | | | | | 1 | |
| 10 | | | | | | | 1 |
| 9 | | | | | | | |
| 8 | | | | 4 | | | 1 |
| 7 | | | | 14 | | | 4 |
| 6 | | | | 6 | | 18 | |
| 5 | | | | | | | |
| 4 | 2 | | | 4 | | | |
| 3 | | | | 14 | | | |
| 2 | | | | | | | |
| 1 | 12 | | | 4 | | | |
| 0 | 18 | 8 | 46 | | | 2 | |
| Total | 32 | 8 | 46 | 69 | 4 | 22 | 26 |

Table 9. Distribution of fish larvae and juveniles by water salinity in upper Chesapeake Bay (1963-67). Numbers represent total number of fish collected at various locations during several years.

| Salinity o/oo | <u>Gasterosteus</u> <u>aculeatus</u> | <u>Hippocampus</u> <u>erectus</u> | <u>Syngnathus</u> <u>fuscus</u> | <u>Roccus</u> <u>americanus</u> | <u>Morone</u> <u>saxatilis</u> | <u>Lepomis</u> <u>gibbosus</u> |
|------------------|---|--------------------------------------|------------------------------------|------------------------------------|-----------------------------------|-----------------------------------|
| 23 | | | | | | |
| 22 | | | 1 | | | |
| 21 | | 1 | 12 | | | |
| 20 | | | 5 | | | |
| 19 | | 1 | 48 | | | |
| 18 | | | 8 | | | |
| 17 | | | 22 | | | |
| 16 | | | 42 | | 1 | |
| 15 | | | 50 | | | |
| 14 | | | 45 | | | |
| 13 | | 1 | 42 | 1 | | |
| 12 | | | 71 | 9 | | |
| 11 | | | 30 | 143 | 2 | |
| 10 | | | 6 | 63 | | |
| 9 | | | 12 | 16 | 18 | |
| 8 | | | 10 | 150 | 4 | |
| 7 | | | 6 | 161 | 9 | |
| 6 | | | | 226 | 2 | |
| 5 | | | 4 | 516 | 55 | |
| 4 | | | 2 | 55 | 5 | |
| 3 | | | 1 | 815 | 104 | |
| 2 | | | 2 | 730 | 87 | |
| 1 | 1 | | | 8,422 | 4,657 | |
| 0 | | | | 80,985 | 2,475 | 2 |
| Total | 1 | 3 | 419 | 92,292 | 7,419 | 2 |

Table 9. Distribution of fish larvae and juveniles by water salinity in (Cont.) upper Chesapeake Bay (1963-67). Numbers represent total number of fish collected at various locations during several years.

| Salinity o/oo | <u>Sciaenops</u> <u>ocellata</u> | <u>Gobionellus</u> <u>boleosoma</u> | <u>Gobiosoma</u> <u>bosci</u> | <u>G. gins-</u> <u>burgi</u> | <u>Microgobius</u> <u>thallasinus</u> | <u>C. bosqui-</u> <u>anus</u> | <u>Hypsob.</u> <u>hentzi</u> |
|------------------|-------------------------------------|--|----------------------------------|---------------------------------|--|----------------------------------|---------------------------------|
| 23 | | | 2 | | | | |
| 22 | | | | | | | |
| 21 | | | 14 | | | | 2 |
| 20 | | | 2 | | | | |
| 19 | 1 | | 5 | 166 | | | |
| 18 | | | 588 | | | | |
| 17 | | | 11 | | 1 | | |
| 16 | | | 91 | | 35 | 44 | |
| 15 | | | 318 | | 9 | 29 | |
| 14 | | | 1,309 | | 35 | 186 | |
| 13 | | | 3,987 | | 8 | 124 | |
| 12 | | | 13,327 | | 14 | 330 | |
| 11 | | 4 | 51,446 | | 2 | 178 | 40 |
| 10 | | | 20,244 | | | 63 | |
| 9 | | | 12,516 | | 8 | 17 | |
| 8 | | | 13,770 | | | 4 | |
| 7 | | | 23,642 | | | | |
| 6 | | | 20,287 | | | | |
| 5 | | | 51,641 | | | 16 | |
| 4 | | | 34,797 | | | | |
| 3 | | | 116,434 | | | | |
| 2 | | | 135,658 | | | | |
| 1 | | | 48,580 | | | | |
| 0 | | | 1,201 | | | | |
| Total | 1 | 4 | 549,870 | 166 | 112 | 991 | 42 |

Table 9. Distribution of fish larvae and juveniles by water salinity in (Cont.) upper Chesapeake Bay (1963-67). Numbers represent total number of fish collected at various locations during several years.

| Salinity o/oo | <u>Peprilus</u> <u>alepidotus</u> | <u>Membras</u> <u>martinica</u> | <u>Menidia</u> <u>beryllina</u> | <u>Menidia</u> <u>menidia</u> | <u>Pseudopleuro-</u> <u>nectes americanus</u> | <u>Trinectes</u> <u>maculatus</u> |
|------------------|--------------------------------------|------------------------------------|------------------------------------|----------------------------------|--|--------------------------------------|
| 23 | | | | | | 32 |
| 22 | 2 | | | | | 1 |
| 21 | 1 | | | | | 36 |
| 20 | | | | | 1 | 47 |
| 19 | 1 | | | | | 37 |
| 18 | | | | | | 3 |
| 17 | 13 | | | | 33 | 17 |
| 16 | 13 | | | | 104 | 33 |
| 15 | 15 | 1 | 1 | | 385 | 86 |
| 14 | 3 | 11 | | 4 | 1,035 | 101 |
| 13 | 42 | 1 | | 14 | 440 | 28 |
| 12 | 16 | 2 | | 78 | 1,000 | 58 |
| 11 | | 16 | 1 | 28 | 1,949 | 30 |
| 10 | | 52 | 4 | 33 | 418 | 26 |
| 9 | | | | 66 | 660 | 54 |
| 8 | | 16 | 22 | 147 | 3,590 | 122 |
| 7 | | 16 | 6 | 448 | 39 | 22 |
| 6 | | 128 | 75 | 38 | 2,043 | 28 |
| 5 | | 2 | 17 | 44 | 112 | 74 |
| 4 | | 122 | 102 | 56 | | 90 |
| 3 | | | 20 | 40 | 2 | 57 |
| 2 | | 41 | 69 | 10 | 19 | 11 |
| 1 | | | 27 | 138 | 1 | 840 |
| 0 | | 21 | 33 | | 3 | 214 |
| Total | 106 | 429 | 377 | 1,144 | 11,834 | 2,047 |

Table 9.. Distribution of fish larvae and juveniles by water salinity in
(Cont.) upper Chesapeake Bay (1963-67). Numbers represent total number
of fish collected at various locations during several years.

| Salinity o/oo | <u>Symphurus</u> <u>plagiusa</u> | <u>Gobiesox</u> <u>strumosus</u> | <u>Sphaeroides</u> <u>maculatus</u> | <u>Chilomycterus</u> <u>schoepfi</u> | <u>Opsanus</u> <u>tau</u> | Totals |
|------------------|-------------------------------------|-------------------------------------|--|---|------------------------------|---------|
| 23 | | | | | | 61 |
| 22 | | | | | | 16 |
| 21 | 10 | | 2 | | | 250 |
| 20 | | | 1 | | | 1,032 |
| 19 | | 1 | 2 | | | 322 |
| 18 | | | 1 | | | 674 |
| 17 | | 1 | 1 | | 2 | 127 |
| 16 | | 2 | 4 | | | 656 |
| 15 | | 1 | 3 | | 5 | 2,732 |
| 14 | | 156 | 17 | | 1 | 4,417 |
| 13 | | 9 | 2 | | | 7,771 |
| 12 | | 255 | 1 | 2 | | 24,161 |
| 11 | | 66 | | | | 66,899 |
| 10 | | 25 | | | | 28,335 |
| 9 | | 6 | | | | 14,040 |
| 8 | | | | | | 25,004 |
| 7 | | | | | | 50,474 |
| 6 | | 12 | | | | 63,967 |
| 5 | | 5 | | | | 68,064 |
| 4 | | 2 | | | | 63,905 |
| 3 | | | | | | 148,253 |
| 2 | | | | | | 144,085 |
| 1 | | | | | | 66,729 |
| 0 | | | | | | 100,729 |
| Total | 10 | 541 | 34 | 2 | 8 | 882,703 |

Table 10. Distribution of fish eggs by water temperature (°C) in upper Chesapeake Bay (1963-67). Numbers represent total numbers of eggs collected at various locations during several years.

| Temp. °C. | Species | | | | |
|---------------|--------------------------------|-------------------------------|------------------------------------|---------------------------------|------------------------------|
| | <u>A.</u> <u>aestivalis</u> | <u>A.</u> <u>mediocris</u> | <u>A.</u> <u>pseudoharengus</u> | <u>A.</u> <u>sapidissima</u> | <u>B.</u> <u>tyrannus</u> |
| 31 | | | | | |
| 30 | | | | | |
| 29 | | | | | |
| 28 | | | | | |
| 27 | | | | | |
| 26 | | | | | |
| 25 | | | | | |
| 24 | | | | | |
| 23 | | | | | |
| 22 | | | | | 3,425 |
| 21 | | | | 38 | |
| 20 | | | | | |
| 19 | | 1 | | | 28 |
| 18 | | | | | 66 |
| 17 | | | | 48 | 70 |
| 16 | | | | | 67 |
| 15 | | 1 | | | 184 |
| 14 | 2,788 | | 184 | 44 | 5,963 |
| 13 | | | 308 | 176 | 65 |
| 12 | | | 218 | 2 | 58 |
| 11 | | | 1 | | |
| 10 | | | 98 | | |
| 9 | 219 | | 9 | | |
| 8 | 16 | | | | |
| 7 | 82 | | 212 | | |
| 6 | | | | | |
| 5 | | | | | |
| 4 | | | | | |
| 3 | | | | | |
| 2 | | | | | |
| 1 | | | | | |
| 0 | | | | | |
| Totals | 3,105 | 2 | 1,030 | 308 | 9,926 |

Table 10. Distribution of fish eggs by water temperature (°C) in upper Chesapeake Bay (1963-67). Numbers represent total numbers of eggs collected at various locations during several years.

| Temp. °C | Species | | | | |
|-------------|----------------------|-------------------------------|--------------------------------|-------------------------------|--------------------------------|
| | <u>Clupeidae</u> sp. | <u>A.</u> <u>mitchilli</u> | <u>R.</u> <u>americanus</u> | <u>R.</u> <u>saxatilis</u> | <u>M.</u> <u>americanus</u> |
| 31 | | 25 | | | |
| 30 | | 2,049 | | | |
| 29 | | 5,775 | | | |
| 28 | | 6,578 | | | |
| 27 | | 175,464 | | | |
| 26 | 8 | 474,390 | | | |
| 25 | | 438,427 | | 10 | |
| 24 | | 292,535 | 6 | 8 | |
| 23 | | 493,390 | | | |
| 22 | | 209,463 | 72 | 20 | |
| 21 | 561 | 148,634 | 88 | 1,989 | |
| 20 | 1,763 | 128,467 | | 2,104 | |
| 19 | 77 | 52,924 | 48 | 437 | |
| 18 | 64 | 31,981 | | 1,332 | |
| 17 | | 6,345 | 1 | 400 | |
| 16 | 13,273 | 13,245 | 12 | 444 | |
| 15 | 24 | 4,471 | 990 | 3,032 | |
| 14 | 6,188 | 13,158 | 1,808 | 7,111 | 627 |
| 13 | 11,376 | 11,435 | 2,773 | 16,279 | |
| 12 | 2,342 | | 9,601 | 2,401 | 364 |
| 11 | 765 | 13 | 548 | 1,020 | |
| 10 | 2,193 | | 4,650 | | |
| 9 | 523 | 2 | 1,722 | | |
| 8 | | | 862 | 1 | |
| 7 | | | 50 | | |
| 6 | | | | | |
| 5 | | | 1 | | |
| 4 | | | | | |
| 3 | | | | | |
| 2 | | | | | |
| 1 | | | | | |
| 0 | | | | | |
| Totals | 39,157 | 2,508,771 | 23,232 | 36,588 | 991 |

Table 10. Distribution of fish eggs by water temperature (°C) in upper (Cont.) Chesapeake Bay (1963-67). Numbers represent total numbers of eggs collected at various locations during several years.

| Species | | | |
|---------------|----------------------|---------------------|-----------|
| Temp. °C | <u>P. americanus</u> | <u>T. maculatus</u> | Totals |
| 31 | | | 25 |
| 30 | | 50 | 2,099 |
| 29 | | 264 | 6,039 |
| 28 | | 2,260 | 8,838 |
| 27 | | 1,355 | 176,819 |
| 26 | | 55,496 | 529,894 |
| 25 | | 19,212 | 457,649 |
| 24 | | 49,053 | 341,602 |
| 23 | | 46,405 | 539,795 |
| 22 | | 106,403 | 319,383 |
| 21 | | 33,315 | 184,625 |
| 20 | | 28,766 | 161,100 |
| 19 | | 46,184 | 99,699 |
| 18 | | 609 | 34,052 |
| 17 | | 233 | 7,097 |
| 16 | | 16 | 27,057 |
| 15 | | | 8,702 |
| 14 | | | 37,871 |
| 13 | | | 42,412 |
| 12 | | | 14,986 |
| 11 | | | 2,347 |
| 10 | | | 6,941 |
| 9 | | | 2,475 |
| 8 | | | 879 |
| 7 | | | 344 |
| 6 | | | |
| 5 | | | 1 |
| 4 | | | |
| 3 | 1 | | 1 |
| 2 | | | |
| 1 | | | |
| 0 | | | |
| Totals | 1 | 389,621 | 3,012,732 |

Table 11. Distribution of fish larvae and juveniles by water temperature (°C) in upper Chesapeake Bay (1963-67). Numbers represent total numbers of fish collected at various locations during several years.

| Temp. °C | Species | | | | | |
|-------------|--|-------------------------------------|----------------------------------|--------------------------------------|------------------------------------|-------------------------------------|
| | <u>Notemigonus</u> <u>crysoleucas</u> | <u>Notropis</u> <u>hudsonius</u> | <u>Ictalurus</u> <u>catus</u> | <u>Ictalurus</u> <u>punctatus</u> | <u>Anguilla</u> <u>rostrata</u> | <u>Strongylura</u> <u>marina</u> |
| 31 | | | | | | |
| 30 | | | | | | |
| 29 | | | | | | 4 |
| 28 | | | | 1 | 4 | |
| 27 | | | | 1 | 3 | 3 |
| 26 | | | | 4 | | 3 |
| 25 | | | 8 | | 9 | |
| 24 | 2 | 2 | | 30 | 16 | 3 |
| 23 | | | | | 13 | |
| 22 | | | | | | 2 |
| 21 | | | | | | |
| 20 | | | | | 6 | 1 |
| 19 | | | | | | |
| 18 | | | | | | |
| 17 | | | | | | |
| 16 | | | | | | |
| 15 | | | | | 10 | |
| 14 | | | | | 20 | |
| 13 | | | | | 4 | |
| 12 | | | | | 6 | |
| 11 | | | | | 3 | |
| 10 | | | | 1 | 13 | |
| 9 | | 6 | | | 3 | |
| 8 | | 8 | | 2 | 6 | |
| 7 | | 2 | 4 | 7 | 4 | |
| 6 | | | | | 2 | |
| 5 | | 6 | | 2 | 13 | |
| 4 | | | | 2 | 6 | |
| 3 | | 4 | | | 4 | |
| 2 | | | | 2 | 4 | |
| 1 | | 2 | | 2 | 2 | |
| 0 | | | | | 2 | |
| Totals | 2 | 30 | 12 | 54 | 153 | 16 |

Table 11. Distribution of fish larvae and juveniles by water temperature (°C) in upper Chesapeake Bay (1963-67). Numbers represent total numbers of fish collected at various locations during several years.

| Temp. °C | Species | | | | | |
|-------------|-----------------------------------|---------------------------------------|------------------------------------|--------------------------------------|----------------------------------|-----------------------------------|
| | <u>Alosa</u> <u>aestivalis</u> | <u>Alosa</u> <u>pseudoharengus</u> | <u>Alosa</u> <u>sapidissima</u> | <u>Brevoortia</u> <u>tyrannus</u> | <u>Clupea</u> <u>harengus</u> | <u>Anchoa</u> <u>mitchilli</u> |
| 31 | | | | | | 940 |
| 30 | | | | | | 922 |
| 29 | | | | | | 3,402 |
| 28 | 6 | | | | | 5,728 |
| 27 | 540 | 56 | | 29 | | 60,550 |
| 26 | 36 | 22 | | | | 40,780 |
| 25 | | 16,148 | 12 | 10 | | 39,556 |
| 24 | 12 | 173 | | 218 | | 16,708 |
| 23 | 108 | 50 | | 738 | | 15,092 |
| 22 | | 1 | | 257 | | 646 |
| 21 | | 2 | 200 | 178 | | 1,184 |
| 20 | | | | 286 | | 1,674 |
| 19 | | | | 66 | | 216 |
| 18 | | | | | | 784 |
| 17 | 2 | | | 48 | | 5,992 |
| 16 | | | | 62 | | 155 |
| 15 | | | | | | 7 |
| 14 | 2 | 2 | | 70 | | 414 |
| 13 | 12 | 1 | | 13 | | 19 |
| 12 | | | | 5 | | 609 |
| 11 | 6 | | | | | 42 |
| 10 | | | 3 | | | 24 |
| 9 | | | | 8 | | 90 |
| 8 | | | | 2 | | 8 |
| 7 | | | | | 2 | 137 |
| 6 | | | | | | 1 |
| 5 | 6 | | | | | 40 |
| 4 | | 1 | | 1 | 2 | 25 |
| 3 | 2 | | | | | 1 |
| 2 | | | | | | 14 |
| 1 | | | | | | 1 |
| 0 | | | | | | 6 |
| Totals | 732 | 16, | 215 | 1,991 | 4 | 197,629 |

Table 11. Distribution of fish larvae and juveniles by water temperature (°C) (Cont.) in upper Chesapeake Bay (1963-67). Numbers represent total numbers of fish collected at various locations during several years.

| Temp. °C | Species | | | | | |
|-------------|--|------------------------------------|---|--------------------------------------|------------------------------------|------------------------------------|
| | <u>Hyporhamphus</u> <u>unifasciatus</u> | <u>Apeltes</u> <u>quadracus</u> | <u>Gasterosteus</u> <u>aculeatus</u> | <u>Hippocampus</u> <u>erectus</u> | <u>Syngnathus</u> <u>fuscus</u> | <u>Roccus</u> <u>americanus</u> |
| 31 | | | | | | |
| 30 | | | | | | |
| 29 | | | | | 2 | |
| 28 | | | | | 2 | 166 |
| 27 | | | | | 9 | 320 |
| 26 | 2 | | | | 34 | 175 |
| 25 | | | | 1 | 47 | 8,259 |
| 24 | 1 | | | 1 | 70 | 793 |
| 23 | | | | 1 | 23 | 333 |
| 22 | 18 | | | | 36 | 229 |
| 21 | | | | | 37 | 877 |
| 20 | 1 | 4 | | | 14 | 30 |
| 19 | | | | | 61 | 6,621 |
| 18 | | 1 | | | 107 | 8,549 |
| 17 | | | | | 20 | 5,478 |
| 16 | | 1 | | | 4 | 106 |
| 15 | | | | | 2 | 1,758 |
| 14 | | 1 | | | 3 | 19,611 |
| 13 | | 3 | | | 4 | 31,672 |
| 12 | | | | | | 15,214 |
| 11 | | | | | 5 | 1,879 |
| 10 | | | | | | 254 |
| 9 | | | | | 2 | 120 |
| 8 | | | | | 2 | 1,888 |
| 7 | | | | | 1 | 15 |
| 6 | | | | | | |
| 5 | | | | | 1 | 9 |
| 4 | | | 1 | | | 32 |
| 3 | | 1 | | | | 30 |
| 2 | | | | | | 31 |
| 1 | | | | | | 9 |
| 0 | | 14 | | | 3 | 2 |
| Totals | 22 | 25 | 1 | 3 | 489 | 104,460 |

Table 11. Distribution of fish larvae and juveniles by water temperature (°C) in upper Chesapeake Bay (1963-67). Numbers represent total numbers of fish collected at various locations during several years.

| Temp. °C | Species | | | | | |
|-------------|-----------------------------------|-----------------------------------|------------------------------------|-----------------------------------|-------------------------------------|------------------------------------|
| | <u>Morone</u> <u>saxatilis</u> | <u>Lepomis</u> <u>gibbosus</u> | <u>Etheostoma</u> <u>nigrum</u> | <u>Perca</u> <u>flavescens</u> | <u>Bairdiella</u> <u>chrysur</u> | <u>Cynoscion</u> <u>regalis</u> |
| 31 | | | | | | |
| 30 | | | | | | |
| 29 | | | | | | 90 |
| 28 | | 2 | | | | 52 |
| 27 | 1 | | | | | 138 |
| 26 | 2 | | | 15 | | 29 |
| 25 | 4 | | 8 | 4 | | 12 |
| 24 | 94 | | | 1 | | 16 |
| 23 | 2 | | | | | |
| 22 | 2 | | | | | |
| 21 | 283 | | | 28 | 3 | 2 |
| 20 | 176 | | | 4 | | |
| 19 | 20 | | | 1 | 5 | 26 |
| 18 | 1,000 | | | | | 4 |
| 17 | 3,532 | | | 956 | | |
| 16 | 2 | | | 7 | | |
| 15 | 28 | | | 16 | | |
| 14 | 332 | | | 2,099 | | |
| 13 | 1,272 | | | 4,488 | | |
| 12 | 447 | | | 6,634 | | |
| 11 | | 3 | | 1,260 | | |
| 10 | | | | 268 | | |
| 9 | | | | 148 | | |
| 8 | | | | 2,009 | | |
| 7 | 1 | | 24 | 26 | | |
| 6 | | | | 66 | | |
| 5 | | | 24 | 1 | | |
| 4 | 4 | | 21 | 3 | | |
| 3 | 5 | | | | | |
| 2 | 3 | | | | | |
| 1 | 1 | | | | | |
| 0 | | | | | | |
| Totals | 7,211 | 5 | 77 | 18,034 | 8 | 369 |

Table 11. Distribution of fish larvae and juveniles by water temperature (Cont.) (°C) in upper Chesapeake Bay (1963-67). Numbers represent total numbers of fish collected at various locations during several years.

| Temp. °C | Species | | | | | |
|-------------|---------------------------------------|--|---------------------------------------|-------------------------------------|--|----------------------------------|
| | <u>Leiostomus</u> <u>xanthurus</u> | <u>Menticirrhus</u> <u>americanus</u> | <u>Micropogon</u> <u>undulatus</u> | <u>Sciaenops</u> <u>ocellata</u> | <u>Gobionellus</u> <u>boleosoma</u> | <u>Gobiosoma</u> <u>bosci</u> |
| 31 | | | | | | 24 |
| 30 | | | | | | 2,572 |
| 29 | | | | | | 19,905 |
| 28 | 3 | | | | | 122,715 |
| 27 | | | | | | 101,998 |
| 26 | | | | | | 54,828 |
| 25 | | | | | 4 | 117,094 |
| 24 | 2 | | 2 | | | 48,176 |
| 23 | | 2 | 3 | 1 | | 72,039 |
| 22 | | | | | | 18,567 |
| 21 | 5 | | | | | 1,860 |
| 20 | | | | | | 70 |
| 19 | 14 | | 4 | | | 842 |
| 18 | 1 | | 1 | | | 130 |
| 17 | | | | | | 152 |
| 16 | | | | | | 20 |
| 15 | 15 | | 3 | | | 30 |
| 14 | | 1 | 2 | | | 1 |
| 13 | | | 2 | | | 20 |
| 12 | 1 | | 30 | | | |
| 11 | | | 36 | | | 6 |
| 10 | | | 5 | | | |
| 9 | | | 28 | | | |
| 8 | | | | | | |
| 7 | | | 34 | | | |
| 6 | | | 2 | | | |
| 5 | | | 19 | | | |
| 4 | | | 39 | | | |
| 3 | | | 18 | | | |
| 2 | | | 36 | | | |
| 1 | | | 5 | | | |
| 0 | | | 1 | | | |
| Totals | 41 | 3 | 270 | 1 | 4 | 561,049 |

Table 11. Distribution of fish larvae and juveniles by water temperature (°C) (Cont.) in upper Chesapeake Bay (1963-67). Numbers represent total numbers of fish collected at various locations during several years.

| Temp. °C | Species | | | | | |
|-------------|--------------------------------------|--|---------------------------------------|---------------------------------------|--------------------------------------|------------------------------------|
| | <u>Gobiosoma</u> <u>ginsburgi</u> | <u>Microbobius</u> <u>thalassinus</u> | <u>Chasmodes</u> <u>bosquianus</u> | <u>Hypsoblennius</u> <u>hentzi</u> | <u>Peprilus</u> <u>alepidotus</u> | <u>Membras</u> <u>martinica</u> |
| 31 | | | 3 | | | |
| 30 | | 2 | 5 | | | |
| 29 | | 2 | 13 | | | |
| 28 | | | 6 | | | 58 |
| 27 | | 19 | 72 | | 1 | 72 |
| 26 | 1 | 8 | 298 | | 7 | 71 |
| 25 | | 31 | 167 | 40 | 21 | 105 |
| 24 | 1 | 49 | 248 | | 36 | 405 |
| 23 | | | 60 | | 26 | 150 |
| 22 | 28 | | 152 | | | 18 |
| 21 | 3 | | 64 | | | |
| 20 | 8 | | 118 | | | |
| 19 | 1 | | 1 | | 4 | |
| 18 | 2 | | 2 | | | 32 |
| 17 | 3 | | | | | 13 |
| 16 | | | | | 1 | |
| 15 | | | | | | |
| 14 | | 1 | | | | 10 |
| 13 | 1 | | | | | |
| 12 | | | | | | |
| 11 | 2 | | | 2 | | |
| 10 | | | | | | |
| 9 | 2 | | | | | |
| 8 | | | | | | |
| 7 | | | | | | |
| 6 | | | | | | |
| 5 | | | | | | |
| 4 | | | | | | |
| 3 | | | | | | |
| 2 | | | | | | |
| 1 | | | | | | |
| 0 | | | | | | |
| Totals | 52 | 112 | 1,209 | 42 | 96 | 934 |

Table 11. Distribution of fish larvae and juveniles by water temperature (°C) (Cont.) in upper Chesapeake Bay (1963-67). Numbers represent total numbers of fish collected at various locations during several years.

| Temp. °C | Species | | | | | |
|---------------|------------------------------|----------------------------|--|--------------------------------|-------------------------------|-------------------------------|
| | <u>Menidia beryllina</u> | <u>Menidia menidia</u> | <u>Pseudopleuronectes americanus</u> | <u>Trinectes maculatus</u> | <u>Symphurus plagiusa</u> | <u>Gobiesox strumosus</u> |
| 31 | | 12 | | | | |
| 30 | | 30 | | | | |
| 29 | 6 | 23 | | 14 | | |
| 28 | 32 | 35 | | 35 | | |
| 27 | 70 | 33 | | 62 | | 2 |
| 26 | 8 | 48 | | 148 | | 31 |
| 25 | 83 | 58 | | 176 | | 27 |
| 24 | 133 | 908 | | 193 | | 2 |
| 23 | 44 | 21 | | 103 | | 10 |
| 22 | 1 | 13 | | 8 | | 71 |
| 21 | 13 | 114 | | 6 | | 2 |
| 20 | | 482 | | 26 | | 37 |
| 19 | | 26 | | 23 | | 8 |
| 18 | | 77 | | 2 | | 336 |
| 17 | | | 2 | 22 | | 1 |
| 16 | | 20 | 11 | 8 | | |
| 15 | | 1 | 25 | 1 | | |
| 14 | 1 | | 66 | 81 | | |
| 13 | 2 | | 317 | 66 | | |
| 12 | 1 | 2 | 314 | 17 | | |
| 11 | | | 2,066 | 34 | 1 | |
| 10 | | 2 | 2,389 | 6 | | |
| 9 | 2 | | 653 | | | |
| 8 | 2 | | 1,990 | 10 | | |
| 7 | 24 | 2 | 741 | 36 | | |
| 6 | | | 89 | 846 | | |
| 5 | | | 3,169 | 73 | | |
| 4 | 1 | | 535 | 8 | | |
| 3 | | 4 | 41 | | | |
| 2 | | | | | | |
| 1 | | | | | | |
| 0 | | | | | | |
| Totals | 423 | 1,911 | 12,408 | 2,004 | 1 | 527 |

Table 11. Distribution of fish larvae and juveniles by water temperature (°C) (Cont.) in upper Chesapeake Bay (1963-67). Numbers represent numbers of fish collected at various locations during several years.

| Temp. °C | Species | | | Total |
|-------------|----------------------------------|-----------------------------------|------------------------|---------|
| | <u>Sphaeroides maculatus</u> | <u>Chilomycterus schoepfi</u> | <u>Opsanus tau</u> | |
| 31 | | | | 979 |
| 30 | | | | 3,531 |
| 29 | | | | 23,461 |
| 28 | | | | 128,845 |
| 27 | | | 2 | 163,981 |
| 26 | 3 | 2 | 2 | 96,557 |
| 25 | 2 | | 1 | 181,887 |
| 24 | 17 | | 1 | 68,313 |
| 23 | 5 | | | 88,824 |
| 22 | 1 | | | 20,050 |
| 21 | | | | 4,861 |
| 20 | 1 | | | 2,938 |
| 19 | 1 | | | 7,940 |
| 18 | 2 | | | 11,366 |
| 17 | | | | 16,221 |
| 16 | 2 | | | 399 |
| 15 | | | 1 | 1,897 |
| 14 | | | | 22,717 |
| 13 | | | | 37,896 |
| 12 | | | | 23,280 |
| 11 | | | 2 | 5,347 |
| 10 | | | | 2,965 |
| 9 | | | | 1,062 |
| 8 | | | | 5,927 |
| 7 | | | | 1,060 |
| 6 | | | | 1,006 |
| 5 | | | | 3,363 |
| 4 | | | | 681 |
| 3 | | | | 110 |
| 2 | | | | 90 |
| 1 | | | 1 | 23 |
| 0 | | | | 28 |
| Totals | 34 | 2 | 10 | 927,267 |

Table 12. Species represented by larval stages in collections from upper Chesapeake Bay (1963-67). Species arranged by order of abundance in the Patuxent River (A) and the upper Bay (C).

| Species | Location of Collection | | | Totals |
|-----------------------------|---|-----------------------|---|---------|
| | <u>A</u> Patuxent R. in order of abundance | <u>B</u> Upper Bay | <u>C</u> Upper Bay in order of abundance | |
| <u>Gobiosoma bosci</u> | 505,606 | 32,264 | 2 | 537,870 |
| <u>Anchoa mitchilli</u> | 179,674 | 15,218 | 3 | 194,892 |
| <u>Roccus americanus</u> | 32,243 | 59,319 | 1 | 91,562 |
| <u>Perca flavescens</u> | 9,195 | 3,239 | 4 | 12,434 |
| <u>Pseudo. americanus</u> | 8,750 | 3,084 | 6 | 11,834 |
| <u>Clupeidae sp.</u> | 5,968 | - | - | 5,968 |
| <u>Menidia sp.</u> | 5,889 | - | - | 5,889 |
| <u>Roccus saxatilis</u> | 6,272 | 1,147 | 7 | 7,419 |
| <u>Brevoortia tyrannus</u> | 2,282 | 13 | 21 | 2,295 |
| <u>Trinectes maculatus</u> | 1,661 | 386 | 10 | 2,047 |
| <u>Menidia menidia</u> | 936 | 208 | 14 | 1,144 |
| <u>Gobiesox strumosus</u> | 500 | 41 | 16 | 541 |
| <u>Chasmodes bosquianus</u> | 464 | 527 | 9 | 991 |
| <u>A. pseudoharengus</u> | 262 | 541 | 8 | 803 |
| <u>Syngnathus fuscus</u> | 218 | 201 | 15 | 419 |
| <u>Micropogon undulatus</u> | 122 | 223 | 13 | 345 |
| <u>Alosa aestivalis</u> | 160 | 3,106 | 5 | 3,266 |
| <u>Membras martinica</u> | 155 | 314 | 11 | 469 |
| <u>Menidia beryllina</u> | 86 | 291 | 12 | 377 |
| <u>Peprilus alepidotus</u> | 76 | 30 | 17 | 106 |
| <u>Anguilla rostrata</u> | 46 | 23 | 18 | 69 |
| <u>Notropis hudsonius</u> | 32 | - | - | 32 |
| <u>Cynoscion regalis</u> | 25 | 18 | 20 | 43 |
| <u>Cyprinodontidae sp.</u> | 10 | - | - | 10 |
| <u>Apeltes quadracus</u> | 7 | 19 | 19 | 26 |
| <u>Cyprinidae sp.</u> | 4 | - | - | 4 |

Table 13. Length frequencies of blueback herring, *Alosa aestivalis*, larvae collected in upper Chesapeake Bay. 1963 and 1967 data combined to produce a general pattern.

| Size in millimeters | | | | | | | | |
|---------------------|-------|-------|-----|------|--------|-----------|---------|----------|
| | March | April | May | July | August | September | October | November |
| 120 | | | | | | | | |
| 115 | | | | | | | | |
| 110 | | | | | | | | |
| 105 | | | | | | | | |
| 100 | | | | | | | | |
| 95 | | | | | | | | |
| 90 | | | 1 | | | | | |
| 85 | | | | | | | | |
| 80 | | | 3 | | | | | |
| 75 | 1 | 2 | 1 | | | | 2 | 1 |
| 70 | | 1 | | | | | 2 | |
| 65 | | | | | | 1 | 3 | 1 |
| 60 | | | | | | 1 | 1 | 1 |
| 55 | | | | | | | 1 | |
| 50 | | | | | 3 | 4 | | |
| 45 | | | | 4 | 17 | 5 | | |
| 40 | | | | 4 | 9 | | | |
| 35 | | | 1 | 7 | | | | |
| 30 | | | | 1 | | | | |
| 25 | | | | | | | | |
| 20 | | | | | | | | |
| 15 | | | | | 1 | 1 | | |
| 10 | | | | | | | | |
| 5 | | | | | | | | |
| 0 | | | | | | | | |

Table 14. Length frequencies of the white perch, *Roccus americanus*, larvae collected in upper Chesapeake Bay. Data for several years (1963-67) combined.

| | | | | | | | | | | | | |
|-----|------|------|------|------|-----|------|------|------|-------|------|------|------|
| 100 | 3 | | | 1 | | | 1 | 1 | | 3 | | |
| 95 | | | 1 | | | | | | | | | |
| 90 | 1 | | | | | | | | 1 | 1 | | |
| 85 | 2 | | 1 | | 2 | | | | 2 | 6 | 1 | |
| 80 | 5 | | 7 | 1 | | | | 2 | | 8 | 2 | |
| 75 | 10 | | 3 | | 2 | | 1 | | 1 | 13 | 1 | |
| 70 | 8 | 1 | 7 | 2 | | | 1 | 2 | 1 | 7 | 3 | |
| 65 | 13 | | 5 | 8 | | | 3 | 1 | | 9 | 3 | |
| 60 | 8 | | 8 | 3 | 1 | 7 | 1 | | | 5 | 1 | |
| 55 | 8 | | 2 | 1 | | 2 | 1 | | | 1 | | |
| 50 | 2 | | 2 | | | 8 | 4 | | | | | |
| 45 | 1 | | | | 1 | 1 | 6 | | | | | |
| 40 | | | | | | 1 | 3 | | | | | |
| 35 | | | | | | 3 | | | | | | |
| 30 | | | | | | 10 | 5 | | | | | |
| 25 | | | | | | 16 | 13 | | | | | |
| 20 | | | | | 2 | 24 | 10 | | | | | |
| 15 | | | | | 7 | 42 | 18 | | | | | |
| 10 | | | | | | 42 | 10 | | | | | |
| 5 | | | | 28 | 40 | 26 | 13 | | | | | |
| 0 | | | 6 | 685 | 514 | 49 | 1 | | | | | |
| | Jan. | Feb. | Mar. | Apr. | May | June | July | Aug. | Sept. | Oct. | Nov. | Dec. |

Table 15. Monthly abundance of striped bass eggs from upper Chesapeake Bay, 1963-67. Data from the Patuxent River and Bay combined.

| Year | Months of Collections | | | | | |
|--------------|-----------------------|------|--------|------|-------|-----|
| | April | | May | | June | |
| | Total | % | Total | % | Total | % |
| 1963 | 664 | 32.9 | 1,344 | 66.6 | 12 | 0.6 |
| 1964 | 48 | 87.5 | 1,150 | 12.5 | - | - |
| 1965 | 3,826 | 99.5 | 18 | 0.5 | - | - |
| 1966 | 3,818 | 32.1 | 8,085 | 68.0 | - | - |
| 1967 | 5,792 | 32.3 | 10,955 | 61.1 | 1,169 | 6.5 |
| Grand Totals | 14,148 | | 21,552 | | 1,181 | |

Table 16. Vertical distribution of the eggs of the striped bass (*Morone saxatilis*) by date and location of collection in upper Chesapeake Bay during 1966 and 1967.

| Sampling Stations | | Sampling Dates | | | | | Total |
|-------------------|----------------|----------------|-------|-------|--------|-------|--------|
| | | 1966 | | 1967 | | June | |
| | | April | May | April | May | | |
| Havre de Grace | S ¹ | - | - | 2 | - | - | 2 |
| | B ² | - | - | 4 | - | 32 | 36 |
| Fishing Battery | S | - | - | - | - | - | - |
| | B | - | - | - | 6 | - | 6 |
| Chesa. City | S | 1,352 | 372 | 2,134 | 856 | 240 | 4,954 |
| | B | 908 | 1,216 | 190 | 1,352 | 368 | 4,034 |
| Old Courthouse | S | 296 | 1,580 | 756 | 690 | 4 | 3,326 |
| | B | 1,002 | 3,098 | 1,210 | 470 | 96 | 5,876 |
| I B | S | - | 3 | - | 2 | 8 | 13 |
| | B | - | 136 | - | - | - | 136 |
| I E | S | 71 | 197 | 292 | 19 | - | 579 |
| | B | 50 | 449 | 173 | 2,440 | 36 | 3,148 |
| II B | S | 71 | 2 | 450 | - | - | 523 |
| | B | 1,124 | 20 | 336 | - | 8 | 1,488 |
| II D | S | 134 | 163 | 35 | 62 | 32 | 426 |
| | B | 5 | 12 | 42 | 18 | 148 | 225 |
| III C | S | - | 348 | 39 | 610 | 20 | 1,017 |
| | B | - | 190 | 6 | 184 | 7 | 387 |
| III E | S | 37 | 344 | 2 | 121 | - | 504 |
| | B | - | - | 3 | 303 | 132 | 438 |
| IV B | S | 17 | 6 | 8 | 154 | - | 185 |
| | B | - | - | 2 | 688 | - | 690 |
| IV D | S | - | 440 | 94 | 2,864 | 2 | 3,400 |
| | B | - | 12 | - | 44 | 16 | 72 |
| V B | S | - | 2 | - | 26 | - | 28 |
| | B | - | - | - | 46 | - | 46 |
| V F | S | - | 3 | 14 | - | - | 17 |
| | B | - | - | - | - | - | - |
| Vert. | S | 1,978 | 3,460 | 3,824 | 5,404 | 306 | 14,974 |
| Dist. | B | 3,089 | 5,133 | 1,962 | 5,551 | 843 | 16,602 |
| Total | | 5,067 | 8,593 | 5,786 | 10,955 | 1,149 | 31,576 |

1 - S = Surface
2 - B = Bottom

Table 17. Length frequencies of yellow perch, Perca flavescens, larvae collected from upper Chesapeake Bay during 1966 and 1967. Data for both years combined.

| | | | | | | |
|-----|---------|-------|-------|-----|------|------|
| 101 | 1 | | | | | |
| 99 | 1 | | | | | |
| 98 | 1 | | | | | |
| 93 | | | | | | 1 |
| 83 | | 1 | | | | |
| 45 | | | | | | 1 |
| 36 | | | | | | 2 |
| 35 | | | | | | 1 |
| 34 | | | | | | 1 |
| 27 | | | | | | 1 |
| 22 | | | | | | |
| 21 | | | | | | 1 |
| 15 | | | | 1 | | |
| 14 | | | | 3 | | |
| 13 | | | | 2 | | |
| 12 | | | | 4 | | |
| 11 | | | | 11 | | |
| 10 | | | 12 | 32 | | |
| 9 | | | 24 | 25 | 1 | |
| 8 | | | 81 | 8 | | |
| 7 | | 1 | 168 | 1 | 1 | |
| 6 | | 1 | 248 | 3 | | |
| 5 | | | 30 | 2 | | |
| 4 | | | | 2 | | |
| | January | March | April | May | June | July |

Table 18. Monthly abundance of the eggs of the bay anchovy (Anchoa mitchilli) from the Chesapeake Bay area for the period 1963-67
Includes samples from Patuxent and Magothy rivers and Chesapeake Bay.

| Months | 1963 | | 1964 | | 1965 | | 1966 | | 1967 | | Total |
|-----------|-----------|------|---------|------|---------|------|---------|------|---------|------|-----------|
| | No. | % | No. | % | No. | % | No. | % | No. | % | |
| April | 2 | + | - | - | - | - | - | - | - | - | 2 |
| May | 35,127 | 1.9 | 56,014 | 13.9 | 36,273 | 10.6 | 1,220 | 0.9 | - | - | 128,634 |
| June | 212,202 | 11.5 | 74,224 | 18.4 | 58,922 | 17.2 | 75,184 | 65.2 | 170,924 | 28.5 | 591,456 |
| July | 1,478,203 | 85.4 | 110,653 | 27.4 | 204,862 | 59.7 | 38,747 | 33.6 | 423,090 | 70.6 | 2,255,555 |
| August | 23,243 | 1.3 | 163,331 | 40.4 | 42,933 | 12.5 | 23 | + | 5,011 | 0.8 | 234,541 |
| September | - | - | - | - | 60 | - | - | - | - | - | 60 |
| Totals | 1,748,777 | | 404,222 | | 343,050 | | 115,174 | | 599,025 | | 3,310,258 |

Table 19. Relative sizes of bay anchovy eggs collected at different salinities. Each range and mean was calculated from 30 individual measurements.

| Salinity | Major Axis | | Minor Axis | |
|----------|------------|-------|------------|-------|
| | Range | Mean | Range | Mean |
| 5 o/oo | 1.33 - .92 | 1.119 | 1.26 - .86 | 1.035 |
| 10 o/oo | 1.36 - .86 | 1.045 | 1.15 - .83 | .978 |
| 15 o/oo | 1.12 - .84 | .968 | 1.09 - .65 | .896 |
| 20 o/oo | 1.11 - .86 | .949 | .95 - .84 | .887 |

Table 20. Frequency distribution of the larvae of the bay anchovy, *Anchoa mitchilli*, by month of collection, water salinity, and vertical distribution in the Patuxent River during 1963.

| Salinity | o/oo | | | | | | | | | | Station | | |
|----------|------|------|-------|-------|--------|---------|--------|------|------|------|---------|---------|-------|
| | | Apr. | May | June | July | Aug. | Sept. | Oct. | Nov. | Dec. | Total | % | |
| 23 | Surf | - | - | - | - | - | - | - | - | - | - | - | - |
| | Bot | - | - | - | - | 26 | - | - | - | - | - | 26 | + |
| 22 | Surf | - | - | - | - | - | - | - | - | - | - | - | - |
| | Bot | - | - | - | - | - | - | - | - | - | - | - | - |
| 21 | Surf | - | - | - | - | - | - | - | - | - | - | - | - |
| | Bot | - | - | - | 151 | 5 | - | - | - | - | - | 156 | 0.1 |
| 20 | Surf | - | - | - | - | - | - | - | - | - | - | - | - |
| | Bot | - | - | 2 | 969 | - | - | - | - | - | - | 971 | 0.6 |
| 19 | Surf | - | - | - | - | - | - | - | - | - | - | - | - |
| | Bot | - | - | 12 | 34 | - | 1 | 1 | - | - | - | 48 | + |
| 18 | Surf | - | - | - | - | - | - | - | - | - | - | - | - |
| | Bot | - | - | - | 49 | - | - | 14 | - | 3 | - | 66 | + |
| 17 | Surf | - | - | - | - | - | - | - | - | - | - | - | - |
| | Bot | - | - | - | - | 9 | 9 | - | - | - | - | 18 | + |
| 16 | Surf | - | - | - | - | - | 1 | - | - | - | - | - | - |
| | Bot | - | 1 | 1 | 198 | 1 | 1 | - | - | 6 | - | 209 | 0.1 |
| 15 | Surf | - | - | - | 1,386 | 42 | - | - | - | - | - | - | - |
| | Bot | - | 4 | - | 91 | 210 | 5 | - | - | 6 | - | 1,744 | 1.0 |
| 14 | Surf | - | - | - | 757 | 34 | 54 | - | - | - | - | - | - |
| | Bot | 1 | - | 1 | 110 | 43 | 190 | 6 | - | 6 | - | 1,202 | 0.7 |
| 13 | Surf | - | 85 | 2 | 83 | 122 | 172 | - | - | - | - | - | - |
| | Bot | - | - | - | 52 | 1,671 | 596 | 4 | - | - | - | 2,787 | 1.6 |
| 12 | Surf | - | - | 2 | 1,935 | 2,948 | 2,680 | - | - | - | - | - | - |
| | Bot | 13 | - | 16 | 658 | 382 | 206 | 2 | - | - | - | 8,842 | 5.0 |
| 11 | Surf | 1 | - | 9 | 179 | 10,754 | - | - | - | - | - | - | - |
| | Bot | 8 | - | - | 658 | 382 | - | - | 2 | 4 | - | 11,997 | 7.0 |
| 10 | Surf | - | - | 159 | 648 | 3,798 | 1,738 | - | - | - | - | - | - |
| | Bot | - | - | 42 | - | 72 | 152 | 2 | 2 | - | - | 6,613 | 3.9 |
| 9 | Surf | 6 | 68 | 79 | 80 | 188 | - | - | - | - | - | - | - |
| | Bot | 8 | - | 22 | 24 | - | - | - | 4 | - | - | 479 | 0.3 |
| 8 | Surf | 6 | - | 826 | - | 284 | 4,024 | - | - | - | - | - | - |
| | Bot | - | 566 | - | - | - | - | - | - | - | - | 5,706 | 3.3 |
| 7 | Surf | - | 540 | 8 | 1,428 | 16,012 | 6,000 | - | - | - | - | - | - |
| | Bot | 62 | 2 | 28 | 24 | - | 30 | - | - | - | - | 24,134 | 14.2 |
| 6 | Surf | - | 434 | 230 | 314 | 37,304 | - | - | - | - | - | - | - |
| | Bot | - | - | 8 | 10 | 566 | - | - | - | 84 | - | 38,950 | 22.8 |
| 5 | Surf | - | 1,544 | 684 | 994 | 6,750 | 1,340 | - | - | - | - | - | - |
| | Bot | - | 42 | 424 | - | 2,304 | - | - | - | - | - | 14,082 | 8.3 |
| 4 | Surf | - | 82 | 186 | 19,208 | - | - | - | - | - | - | - | - |
| | Bot | - | - | - | 160 | - | 140 | - | - | - | - | 19,776 | 11.6 |
| 3 | Surf | 84 | - | 116 | 200 | 22,172 | 3,604 | - | - | - | - | - | - |
| | Bot | - | 110 | 10 | - | 240 | - | - | - | - | - | 26,536 | 15.6 |
| 2 | Surf | - | 538 | - | 4,508 | - | - | - | - | - | - | - | - |
| | Bot | - | - | 4 | 30 | - | - | - | - | - | - | 5,080 | 3.0 |
| 1 | Surf | 12 | 676 | 526 | - | - | - | - | - | - | - | - | - |
| | Bot | - | 12 | - | - | - | - | - | - | - | - | 1,226 | 0.2 |
| 0 | Surf | 24 | - | 128 | - | - | - | - | - | - | - | - | - |
| | Bot | - | - | - | - | - | - | - | - | - | - | 152 | 0.1 |
| | Surf | 133 | 3,967 | 2,955 | 31,720 | 100,408 | 19,613 | - | - | - | - | 158,794 | - |
| | Bot | 92 | 737 | 570 | 3,218 | 5,911 | 1,330 | 29 | 8 | 109 | - | 12,006 | - |
| Total | | 225 | 4,704 | 3,525 | 34,938 | 106,319 | 20,943 | 29 | 8 | 109 | - | 170,800 | 100.1 |
| % | | 0.1 | 2.8 | 2.1 | 20.3 | 62.4 | 12.3 | + | + | 0.1 | - | 0.1 | 99.9 |

Table 21. Length frequency of bay anchovy larvae and juveniles collected in upper Chesapeake Bay. Data for several years combined (1963-67).

| Size in millimeters | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
|---------------------|------|---|---|----|---|------|---|---|---|----|------|----|----|----|----|------|----|----|----|----|-----|----|----|----|----|------|----|----|----|----|----|
| | ** 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22 | 23 | 24 | 25 | 26 | 27 | 28 | 29 | | |
| | Jan. | | | | | Feb. | | | | | Mar. | | | | | Apr. | | | | | May | | | | | June | | | | | |
| 120 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 115 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 110 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 105 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 100 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| 95 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 90 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 85 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 80 | | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | 1 | 1 | 1 |
| 75 | | | | | | | | | | | | | | | | | | | 4 | 1 | 3 | | | | | 2 | | 2 | 6 | 3 | |
| 70 | | | | | | | | | | | | | | | | | | | 4 | 1 | 6 | | | | 1 | 2 | 2 | 2 | 4 | 4 | |
| 65 | | | | | | | | | | | | | | | | | | | 1 | 3 | 13 | 1 | | | 3 | 4 | 4 | 8 | 15 | 13 | |
| 60 | | | | | | | | | | | | | | | | | | | 7 | 2 | 13 | | | 6 | 9 | 8 | 15 | 11 | 12 | 1 | |
| 55 | | | | | | | | | | | | | | | | | | | 1 | 7 | 11 | 10 | | 1 | 12 | 19 | 9 | 12 | 5 | 1 | |
| 50 | 2 | 1 | | | | | | | | | | | | | | | | | 10 | 5 | 29 | 1 | 11 | 9 | 14 | 1 | | | 1 | | |
| 45 | | | | | | | | | 1 | | | | | | | | | | 1 | | 8 | | 6 | 5 | 4 | 1 | 1 | | | | |
| 40 | 4 | | | 3 | 1 | | | | | | | | | | | | | | | 2 | 1 | 1 | 2 | 1 | 1 | | | | | | |
| 35 | 9 | 1 | | 11 | 1 | | | | | 4 | | | | | | | | | | 2 | | | 1 | | | | | | | | |
| 30 | 8 | 3 | | 10 | | | | | 1 | | 1 | | | | | | | | 3 | 1 | | | | | | | | | | 1 | |
| 25 | 2 | 3 | | 3 | | | | | | | | | | | | | | | | | | | | | | | | | | | 2 |
| 20 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 9 |
| 15 | | | | | | | | | | | | | | | | | | | | | | | | | | | | 2 | 1 | 7 | 26 |
| 10 | | | | | | | | | | | | | | | | | | | | | | | | | | | | 2 | 44 | 12 | 48 |
| 5 | | | | | | | | | | | | | | | | | | | | | 5 | 1 | | | | | | 32 | 58 | 62 | 71 |
| *0 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | 2 | 11 | 18 |

* Categories of 5 mm: 0-4.9, 5-9.9, 10-14.9

** Each month divided into 5 six-day periods; 31st day added to last period

Table 21. (Continued).

| | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
|----|------|-----|----|------|----|----|----|-------|----|----|----|------|----|----|----|------|----|----|------|----|----|----|----|----|----|----|----|----|----|
| 1 | | | | | | | | | | | | | | 1 | 1 | | | | | | | | | | | | | | |
| | 1 | 1 | | 3 | | | | 1 | 1 | | | | 1 | | | | | | | | | | | | | | | | |
| | | 3 | | 3 | 2 | 2 | 1 | 1 | 1 | 1 | | 3 | 1 | | 1 | 1 | 1 | | | 1 | | | | | | | | | |
| | 3 | | 6 | 7 | 1 | 4 | | 2 | 1 | 2 | 2 | 3 | 1 | 4 | 1 | | | | | 2 | | | | | | | | | |
| 1 | 1 | 5 | 3 | 4 | 2 | 2 | 4 | 4 | 1 | 6 | 1 | 7 | 4 | 3 | 3 | | | | | 1 | | | | | | | | | |
| 2 | 3 | 4 | 1 | 7 | | | 2 | 2 | 10 | | 4 | 2 | 5 | 5 | 3 | | 1 | 1 | | 1 | | | | | | | | | |
| 3 | 1 | 4 | | 4 | 2 | 2 | | 1 | 2 | 2 | | 2 | 2 | 1 | 4 | | | | | 3 | | | | | | | | | |
| 1 | | | | | 2 | 3 | 2 | | 2 | 1 | 1 | 4 | 2 | 20 | 22 | 11 | 2 | | | 4 | | | | | | | | | |
| | 1 | | 3 | 6 | | | 3 | 2 | | 1 | 2 | 8 | 4 | 59 | 28 | 12 | 3 | | 5 | 2 | 1 | | | | | | | | |
| | | 1 | 3 | 2 | 9 | 4 | 3 | 5 | 4 | 11 | 4 | 5 | 7 | 89 | 35 | 13 | 3 | | 5 | | 4 | | | | | | | | |
| | 6 | 7 | 4 | 4 | 21 | 15 | 13 | 14 | 1 | 10 | 3 | 5 | 1 | 53 | 21 | 23 | 3 | 11 | | 4 | 1 | 5 | | | | | | | |
| | 1 | 15 | 5 | 5 | 4 | 35 | 21 | 16 | 15 | 1 | 5 | 4 | 2 | 4 | 62 | 33 | 32 | | 13 | 5 | 1 | 1 | 1 | 3 | | | | | |
| | 3 | 11 | 9 | 14 | 10 | 40 | 18 | 17 | 20 | | 10 | 3 | 15 | 5 | 60 | 29 | 16 | 7 | 6 | 10 | 10 | 5 | 8 | 3 | 7 | 5 | | | |
| | 4 | 31 | 11 | 23 | 7 | 48 | 9 | 29 | 35 | 5 | 16 | 4 | 13 | 7 | 28 | 25 | 8 | 4 | 4 | 13 | 2 | 3 | 6 | 1 | 5 | 2 | | | |
| 10 | 23 | 35 | 32 | 60 | 13 | 44 | 5 | 21 | 34 | 9 | 30 | 13 | 32 | 13 | 57 | 13 | 4 | | 1 | 4 | | | 1 | | | 5 | | | |
| 6 | 52 | 62 | 42 | 58 | 32 | 46 | 6 | 49 | 33 | 3 | 43 | 16 | 16 | 13 | 26 | 5 | | | | | | 3 | | 2 | | 1 | | | |
| | 73 | 90 | 75 | 141 | 44 | 70 | 9 | 50 | 56 | | 33 | 3 | 23 | | 5 | 5 | 2 | | | | | 2 | | | | | | | |
| 5 | 35 | 148 | 93 | 148 | 61 | 79 | | 45 | 12 | | 7 | | 6 | | | 3 | | | | | | | 1 | | | | | | |
| | 25 | 95 | 44 | 61 | 33 | 22 | | 18 | | | | | | | 1 | | | | | | | | | | | | | | |
| 30 | 31 | 32 | 33 | 34 | 35 | 36 | 37 | 38 | 39 | 40 | 41 | 42 | 43 | 44 | 45 | 46 | 47 | 48 | 49 | 50 | 51 | 52 | 53 | 54 | 55 | 56 | 57 | 58 | 59 |
| | July | | | Aug. | | | | Sept. | | | | Oct. | | | | Nov. | | | Dec. | | | | | | | | | | |

Table 22. Length frequencies of northern pipefish, *Syngnathus fuscus*, collected at Solomons, Maryland from 17 July 1961 to 25 July 1962.

| | | | | | | | | | | | | |
|-----|------|------|-------|------|------|------|------|------|------|-----|------|------|
| 110 | 2 | | | | | | | 3 | 5 | | | |
| 105 | | | | | | | | 1 | | 1 | | |
| 100 | 1 | | | | | | | | | | | |
| 95 | | | | | | | | | | 1 | | |
| 90 | | | 1 | | | | | 1 | | | | |
| 85 | 1 | | | | | | | | | 1 | 1 | |
| 80 | | | | | | 1 | 1 | | 1 | 1 | | |
| 75 | 2 | | | | | | | | | 1 | 1 | |
| 70 | 1 | | | | | | | | | 2 | | |
| 65 | 2 | | | | | | | | | 3 | | |
| 60 | 2 | | | | | | | | | 2 | 1 | |
| 55 | 7 | | 1 | | | | 1 | | | 2 | | |
| 50 | | | | 1 | 1 | | | | | 5 | | |
| 45 | 6 | 1 | | 3 | | | | | | 29 | | |
| 40 | 9 | 2 | | 6 | | | | | 3 | 64 | | |
| 35 | 4 | 1 | 2 | 3 | | 1 | | | 11 | 55 | | |
| 30 | 1 | | 4 | 5 | 1 | | | | 26 | 27 | 1 | |
| 25 | 1 | | 2 | 4 | | | | | 32 | 11 | | |
| 20 | | | 2 | 8 | | | | | 30 | 11 | 1 | |
| 15 | 1 | 1 | 2 | 13 | | | | | 48 | 18 | | |
| 10 | | 5 | 7 | 4 | | | | | 21 | 22 | 2 | |
| 5 | | 1 | | | | | | | | 3 | | |
| 0 | | | | | | | | | | | | |
| | July | Aug. | Sept. | Oct. | Nov. | Dec. | Jan. | Mar. | Apr. | May | June | July |
| | | | 1961 | | | | | | | | | 1962 |

Table 23. Length frequencies of northern pipefish, Syngnathus fuscus, collected in upper Chesapeake Bay. Data from the period 1963-67 combined.

| Size in millimeters | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
|---------------------|------|---|---|---|---|------|---|---|---|----|------|----|----|----|----|------|----|----|----|----|-----|----|----|----|----|------|----|----|----|--|
| | ** 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22 | 23 | 24 | 25 | 26 | 27 | 28 | 29 | |
| | Jan. | | | | | Feb. | | | | | Mar. | | | | | Apr. | | | | | May | | | | | June | | | | |
| 120 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 115 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 110 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 105 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 100 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 95 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 90 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 85 | 1 | | | | | | | | | 1 | | | | | | | | | | | | | | | | | | | | |
| 80 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 75 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 70 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 65 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 60 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 55 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 50 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 45 | 1 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 40 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 35 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 30 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 25 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 20 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 15 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 10 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 5 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| *0 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |

* Categories of 5 mm: 0-4.9, 5-9.9, 10-14.9.

** Each month divided into 5 six-day periods. 31st day added to last period.

Table 23. (Continued).

| | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
|-----|------|----|----|----|----|----|------|----|----|----|----|-------|----|----|----|----|------|----|----|----|------|----|----|----|------|----|----|----|----|----|---|--|--|--|---|---|
| 120 | 1 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 115 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 110 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 105 | 1 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 100 | 1 | | | | | | | | | | | 2 | | | | | | | | | | | | | | | | | | 1 | | | | | | |
| 95 | 1 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 90 | 1 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 85 | 1 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 80 | 4 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 75 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 70 | 2 | | | | | | | | | | | 1 | | | | | | | | | | | | | | | | | | | | | | | | |
| 65 | 1 | | | | | | | | | | | 1 | | | | | | | | | | | | | | | | | | | | | | | | |
| 60 | 1 1 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 55 | 1 | | | | | | | | | | | 5 | | | | | | | | | | | | | | | | | | 2 | 1 | | | | | |
| 50 | 3 | 1 | 4 | 1 | | | | | | | 1 | | | | | | | | | | | 2 | | 1 | | | | | | | | | | | 3 | 1 |
| 45 | 1 | 1 | 3 | 1 | 2 | 1 | 1 | | | | | | | | | | | 2 | 1 | 5 | | | | | | | | | | | 1 | | | | | |
| 40 | 2 | 1 | 9 | 3 | 1 | 3 | | | | | | | | | | | 2 | 4 | 2 | 1 | | | | | | | | | | | | | | | | |
| 35 | 2 | 1 | 1 | | | | | | | | | | | | 2 | | 5 | | | | | | | | | | | | | | | | | | | |
| 30 | 2 | | | 2 | | | | | | | | | | | 2 | 2 | 7 | 3 | 1 | | | | | | | | | | | | | | | | | |
| 25 | 2 | | | 3 | | | 1 | | | | | | | | | | | 1 | 1 | 1 | 2 | 15 | | | | | | | | | | | | | | |
| 20 | 2 | | | 1 | | | | | | | | | | | 2 | 1 | 2 | | | | | | | | | | | 17 | | | | | | | | |
| 15 | 1 | | | 2 | | | | | | | | | | | 1 | 4 | 1 | 12 | 16 | 2 | | | | | | | | | | | | | | | | |
| 10 | 1 | 1 | 4 | 1 | 3 | 1 | 5 | 19 | 2 | 2 | 13 | 17 | | | | | | | | | | | | | | | | | | | | | | | | |
| 5 | 1 | | | 2 | | | 3 | | | 2 | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 0 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| | 30 | 31 | 32 | 33 | 34 | 34 | 36 | 37 | 38 | 39 | 40 | 41 | 42 | 43 | 44 | 45 | 46 | 47 | 48 | 49 | 50 | 51 | 52 | 53 | 54 | 55 | 56 | 57 | 58 | 59 | | | | | | |
| | July | | | | | | Aug. | | | | | Sept. | | | | | Oct. | | | | Nov. | | | | Dec. | | | | | | | | | | | |

Table 24. Length frequencies of naked goby, Gobiosoma bosci, larvae collected from the Patuxent River during 1963.

| Size in Millimeters | May | June | July | August | September | October |
|---------------------|-----|------|------|--------|-----------|---------|
| | 15 | | | | | 1 |
| 14 | | 1 | 1 | | | |
| 13 | | 2 | 4 | | | |
| 12 | | 11 | 20 | 4 | | 2 |
| 11 | | 19 | 40 | 14 | 5 | 7 |
| 10 | | 14 | 54 | 37 | 31 | 5 |
| 9 | | 8 | 42 | 25 | 21 | 3 |
| 8 | | 14 | 39 | 22 | 14 | 1 |
| 7 | | 23 | 29 | 26 | 16 | 2 |
| 6 | | 30 | 36 | 36 | 18 | |
| 5 | | 52 | 45 | 45 | 12 | |
| 4 | | 66 | 46 | 17 | 1 | |
| 3 | | 27 | 56 | 16 | | |
| 2 | | 1 | 15 | 4 | | |
| 1 | | | | | | |
| 0 | | | | | | |

Table 25. Length frequencies of winter flounder, Pseudopleuronectes americanus, larvae collected in upper Chesapeake Bay. Data from several years (1963-1967) and idfferent locations combined.

| Size in millimeters | | | | |
|---------------------|----------|-------|-------|-----|
| | February | March | April | May |
| 23 | | | | |
| 22 | | | | |
| 21 | | | | |
| 20 | | | | |
| 19 | | | | |
| 18 | | | | |
| 17 | | | | |
| 16 | | | | 1 |
| 15 | | | | 1 |
| 14 | | | | |
| 13 | | | | |
| 12 | | | | |
| 11 | | | | |
| 10 | | | 1 | |
| 9 | | 2 | 32 | |
| 8 | | 8 | 173 | |
| 7 | | 23 | 161 | 2 |
| 6 | | 60 | 155 | |
| 5 | | 108 | 72 | |
| 4 | 1 | 166 | 19 | |
| 3 | 9 | 95 | 3 | |
| 2 | | 7 | | |
| 1 | | | | |
| 0 | | | | |
| | February | March | April | May |

Table 26. Length frequencies of menhaden, Brevoortia tyrannus, collected in upper Chesapeake Bay, 1963-67. Data compiled from several areas.

| Size in millimeters | | | | | | | | | | | | | | | | | | | | | | | | | |
|---------------------|------|------|----|----|----|----|-----|----|----|----|----|------|----|----|----|----|------|----|----|----|----|----|----|--|--|
| | Mar. | Apr. | | | | | May | | | | | June | | | | | Nov. | | | | | | | | |
| 120 | | | | | | | | | | | | | | | | | | | | | | | | | |
| 115 | | | | | | | | | | | | | | | | | | | | | | | | | |
| 110 | | | | | | | | | | | | | | | | | | | | | | | | | |
| 105 | | | | | | | | | | | | | | | | | | | | | | | | | |
| 100 | | | | | | | | | | | | | | | | | | | | | | | | | |
| 95 | | | | | | | | | | | | | | | | | | | | | | | | | |
| 90 | | | | | | | | | | | | | | | | | | | | | | | | | |
| 85 | | | | | | | | | | | | | | | | | | | | | | | | | |
| 80 | | | | | | | | | | | | | | | | | | | | | | | | | |
| 75 | | | | | | | | | | | | | | | | | | | | | | | | | |
| 70 | | | | | | | | | | | | | | | | | | | | | | | | | |
| 65 | | | | | | | | | | | | | | | | | | | | | | | | | |
| 60 | | | | | | | | | | | | | | | | | | | | | | | | | |
| 55 | | | | | | | | | | | | | | | | | | | | | | | | | |
| 50 | | | | | | | | | | | | | | | | | | | | | | | | | |
| 45 | | | | | | | | | | | | | | | | | | | | | | | | | |
| 40 | | | | | | | | | | | | | | | | | | | | | | | | | |
| 35 | 3 | | | 3 | 3 | 4 | 13 | 30 | 5 | 31 | 7 | 5 | 2 | 2 | | | | | | | | | | | |
| 30 | 1 | 4 | 7 | 6 | 9 | 6 | 27 | 37 | 6 | 15 | 4 | 3 | 1 | | | | | | | | | | | | |
| 25 | | 4 | 4 | 2 | 1 | 2 | 3 | | | | | | 2 | | | | | | | | | | | | |
| 20 | | | | | | | | | | | | | | | | | | | | | | | | | |
| 15 | | | | | | | | | | | | | | | | | | | | | | | | | |
| 10 | | | | | | | | | | | | | | | | | | | | | | | | | |
| 5 | | | | | | | | | | | | | | | | | | | | | | | | | |
| 0 | | | | | | | | | | | | | | | | | | | | | | | | | |
| | **13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22 | 23 | 24 | 25 | 26 | 27 | 28 | 29 | 30 | 51 | 52 | 53 | 54 | 55 | | |
| | Mar. | Apr. | | | | | May | | | | | June | | | | | Nov. | | | | | | | | |

* Categories of 5 mm: 0-4-9, 4-9.9; 10-14.9. 1/(4 of 24 measured).
 ** Each month divided into 5 six-day periods. 31st day added to last period.

Table 27. Cumulative percentages of fish larvae collected from different salinity ranges in Chesapeake Bay. Data compiled from the Patuxent, Magothy, and Susquehanna River gradients during the period 1963-1967.

| Salinity (ppt) | Percent | |
|----------------|---------|--------------------------|
| 0 | 11.4 | |
| 0-1 | 18.9 | |
| 0-2 | 35.2 | |
| 0-3 | 52.0 | |
| 0-4 | 59.2 | |
| 0-5 | 66.9 | |
| 0-6 | 74.1 | |
| 0-7 | 79.8 | |
| 0-8 | 82.6 | |
| 0-9 | 84.2 | Total number |
| 0-10 | 87.4 | of organisms = 5,148,596 |
| 0-11 | 95.0 | |
| ----- | | |
| | | Eggs = 3,879,478 |
| 0-12 | 97.7 | Larvae = 1,269,118 |
| 0-13 | 98.6 | |
| 0-14 | 99.0 | |
| 0-15 | 99.4 | |
| 0-16 | 99.5 | |
| 0-17 | 99.5+ | |
| 0-18 | 99.6 | |
| 0-19 | 99.6+ | |
| 0-20 | 99.7 | |
| 0-21 | 99.7+ | |
| 0-22 | 99.7+ | |
| 0-23 | 99.7+ | |

ACKNOWLEDGMENTS

During the period of this investigation, the services of many people have been used to make field collections and to process data. The dedication with which individuals performed their duties and the generosity of others not affiliated with the project have helped make this contribution possible.

The late Dr. Romeo J. Mansueti was responsible for initiating this project. His enthusiasm during the first three years of this project contributed substantially toward developing the momentum necessary to complete this work.

My appreciation is expressed to Dr. Cronin for making available the necessary facilities and time to continue this work at the Chesapeake Biological Laboratory. Financial support has been provided by the State of Maryland, Maryland Dept. of Water Resources, U. S. Army Corps of Engineers, Magothy River Association, Chesapeake Biological Laboratory, and the Natural Resources Institute of the University of Maryland.

I am especially thankful to Elgin Dunnington for his review of the entire manuscript.

My sincere thanks go to the following individuals:

BOAT OPERATORS:

Martin O'Berry, Captain R/Vs COBIA and ORION, Richard Younger,
Elgin Dunnington, David Cargo

FIELD ASSISTANTS:

Edward Newell, Chris Garbo, James R. Edmunds IV, Dr. Robert Biggs,
Dr. David Flemer

SAMPLE PROCESSORS:

Ellen Garner, Frances Younger, Lotte Dutcher, Minerva Hellerich, Gail
Oneyear, Peggy Windes, Alice Thompson, Elloleen Lippson

DATA PROCESSORS:

Vilma Garner, Benny Dutcher, Patsy Elliott, Arie de Kok

GRAPHICS:

Frances Younger



The Natural Resources Institute of the University of Maryland carries on an interwoven program of research, education, and training in the field of natural resources. It works under the direction of the University's Board of Regents and in cooperation with other public and private agencies and institutions. The Institute strives—through greater knowledge—to advance the best uses of Maryland's forests, fish, wildlife, and surface waters for the benefit of people, now and in the future.