

Seasonal forecasts of Chesapeake Bay hypoxia

Chesapeake Hypoxia Analysis and
Modeling Program (CHAMP) Meeting

October 30, 2019

Isabella Bertani¹ & Don Scavia²

¹ University of Maryland Center for Environmental Science

² University of Michigan

Seasonal forecasts of Chesapeake Bay hypoxia



Near-record dead zones forecast for Chesapeake Bay, Gulf of Mexico

Home » Maryland News » Chesapeake Bay's 'dead zone'...

Chesapeake Bay's 'dead zone' expected to get bigger

 John Aaron | @JohnAaronWTOP
June 13, 2019, 4:00 AM

ScienceDaily
Your source for the latest research news
Science News *from research organizations*

Large summer 'dead zone' forecast for Chesapeake Bay after wet winter and spring

Date: June 12, 2019
Source: University of Maryland Center for Environmental Science

Chesapeake Bay 'dead zone' could be largest in decades, scientists say

[Lucas Gonzalez](#), Salisbury Daily Times Published 6:00 a.m. ET June 20, 2019 | Updated 4:57 p.m. ET June 20, 2019

University of Michigan Chesapeake Bay hypoxia forecasting model


Streeter-Phelps Model

Biological Oxygen Demand (BOD):

$$\frac{dBOD}{dt} = -v * \frac{dBOD}{dx} - a * BOD$$

Dissolved Oxygen (DO):

$$\frac{dDO}{dt} = -v * \frac{dDO}{dx} + a * BOD - b * DO$$



t: time (d)


x: distance from source of *BOD* (km)

a: *BOD* decomposition rate (d^{-1})

b: *DO* re-aeration rate (d^{-1})

v: downstream advection ($km\ d^{-1}$)

University of Michigan Chesapeake Bay hypoxia forecasting model



Model driver: Jan-May average **TN load from Susquehanna**
at Conowingo



TN → **C** through Redfield Ratio (5.67 gC/gN)

F: fraction of **C** assumed to settle below the pycnocline

C → **BOD** through respiration ratio (2.4 gO₂/gC)

University of Michigan Chesapeake Bay hypoxia forecasting model

Driver:

Jan-May average

Susquehanna TN load



Calibration target:

Mean July hypoxic volume (HV)

([DO] < 2 mg/L)




Model output:

Average subpycnocline
[DO] as a function of
distance from TN source



Hypoxic length = sum
of all segments with
[DO] < 2 mg/L




Hypoxic length → **hypoxic volume**
through empirical V-L relationship

University of Michigan Chesapeake Bay hypoxia forecasting model

Seasonal forecast

The 2019 Forecast - Given the average January-May 2019 total nitrogen load of 309,403 kg/day, this summer's hypoxia volume forecast is 8.9 km³, the 4th largest in the past 20 years.




The average 2019 forecast is 8.9 km³.

There is a 95% probability that hypoxic volume will be between 7.5 and 10.2 km³.



University of Michigan Chesapeake Bay hypoxia forecasting model

Forecasting track record





Lessons learned from model track record and other analyses

Average July HV somewhat “arbitrary” metric and highly sensitive to transitory weather disruptions




Lessons learned from model track record and other analyses

Multiple estimates of HV now available, both from observations and 3D models – Opportunity to incorporate multiple sources of information during Bayesian calibration



Lessons learned from model track record and other analyses

Preliminary analyses from CHAMP group suggest that loading periods other than Jan-May might be relevant to total annual hypoxia



Lessons learned from model track record and other analyses

Susquehanna works as a reasonable proxy for total load, but including other sources may improve model performance

Relative contribution* of different geobasins to hypoxia as estimated by the CBP model

*Based on the effect of N and P loads from each basin on the 25th percentile of summer DO concentrations below the surface mixed layer

Geobasin	N	P	Total
JmsA	0.9%	0.4%	1.3%
PotA	16.3%	1.9%	18.2%
PxtA	0.5%	0.1%	0.6%
RapA	0.9%	0.2%	1.1%
Susq	45.0%	4.4%	49.4%
YrkA	0.3%	0.1%	0.4%
EshLow	3.3%	0.5%	3.8%
EshMid	1.8%	0.6%	2.4%
EshUpp	2.2%	0.5%	2.7%
EshVA	0.8%	0.1%	0.9%
JmsB	1.3%	0.3%	1.6%
PotB	6.7%	1.1%	7.8%
PxtB	0.9%	0.2%	1.1%
RapB	1.2%	0.2%	1.3%
Wsh	5.4%	1.2%	6.5%
YrkB	0.6%	0.1%	0.7%
Total	88.1%	11.9%	100.0%

Planned short-term revisions to the University of Michigan Chesapeake Bay hypoxia forecasting model – before 2020 forecast

Re-calibrate model to different sets of HV estimates, HV metrics, loading periods and load sources

HV estimates:

- HV estimated through interpolation of cruise data
- Simulated HV from 3D models (e.g., VIMS, UMCES)


HV metrics: average July, average summer, total annual, monthly

Load sources: consider major load sources other than Susquehanna (e.g., Potomac, Rappahannock, cumulative point sources)

Compare model skill and track record (e.g. through blind forecasting) and uncertainty across different calibration versions


University of Michigan Gulf of Mexico hypoxia forecasting model

Forecasting track record



University of Michigan Gulf of Mexico hypoxia forecasting model

Management application – Ensemble of four models



Planned short-term revisions to the University of
Michigan Chesapeake Bay hypoxia forecasting
model – before 2020 forecast

Thank You

Feedback and suggestions from the CHAMP
group welcome!

Isabella Bertani: ibertani@chesapeakebay.net

Don Scavia: scavia@umich.edu