

Using the Tributary Summaries: *Case Studies of the Potomac and York Tributaries*

Modeling Workgroup Quarterly Meeting
January 5, 2022

Presented by: Breck Sullivan (USGS)

Presentation Materials from:
Rebecca Murphy (UMCES/CBP)
Vanessa Van Note (EPA)
Jeni Keisman (USGS)

Rappahannock Tributary Summary:

A summary of trends in tidal water quality and associated factors, 1985-2018.

June 7, 2021

Prepared for the Chesapeake Bay Program (CBP) Partnership by the CBP Integrated Trends Analysis Team (ITAT)




This tributary summary is a living document in draft form and has not gone through a formal peer review process. We are grateful for contributions to the development of these materials from the following individuals: Jeni Keisman, Rebecca Murphy, Olivia Devereux, Jimmy Webber, Qian Zhang, Meghan Petenbrink, Tom Butler, Zhaoying Wei, Jon Harcum, Renee Karrh, Mike Lane, and Elgin Perry.

Overview of Presentation

1. What are the Tributary Summaries?
2. Looking at two Tributary Summaries
 - *The Potomac*
 - *The York River*
3. Potomac Story Map
4. Next Steps for the Tributary Summaries

13 Tributary Trend Summaries




- **Maryland Mainstem** (*The 5 Chesapeake Bay mainstem segments within the MD state boundary. Drainage basins include the Susquehanna River and upper Chesapeake shorelines*)
- **Maryland Upper Eastern Shore** (*The Northeast, Bohemia, Elk, Back Creek, Sassafras, and Chester Rivers, the C&D Canal, and Eastern Bay*)
- **Choptank** (*the Choptank, Little Choptank, and Honga*)
- **Maryland Upper Western Shore** (*Bush, Gunpowder, Middle Rivers*)
- **Maryland Lower Western Shore** (*Magothy, Severn, South, Rhode, and West*)
- **Patapsco & Back Rivers**
- **Patuxent** (*includes the Western Branch tributary*)
- **Potomac**
- **Rappahannock** (*includes the Corrotoman tributary*)
- **York** (*includes the Mattaponi and Pamunkey tributaries*)
- **James** (*includes the Appomattox, Chickahominy, and Elizabeth tributaries*)
- **Lower E. Shore** (*includes the Nanticoke, Manokin, Wicomico, Big Annemessex, and Pocomoke rivers & Tangier Sound*)
- **Virginia Mainstem** (*no summary but Appendices are provided*)

What are the Tributary Summaries?



A compilation of information by tributary or region on:

- Tidal water quality and trends



Status	Type of trend
Not meeting criterion	Significant ($p < 0.05$)
Meeting criterion	Degrading
Not applicable	Improving
	Unlikely ($p > 0.25$)


Figure 5. Pass-fail DO criterion status for 30-day OW summer averages in Rappahannock segments along with long-term trends in Chesapeake Bay Program, www.chesapeakebay.net, North






What are the Tributary Summaries?

A compilation of information by tributary or region on:

- Tidal water quality and trends,
- **Watershed characteristics and changes**



SGS Station Name	Trend start water year	Percent change in FN load, through water year 2018		
		TN	TP	SS
RAPPAHANNOCK RIVER AT REMINGTON, VA	1985	24.4	-	-
RAPPAHANNOCK RIVER AT REMINGTON, VA	2009	15.4	-	-
RAPPIDAN RIVER NEAR RUCKERSVILLE, VA	2009	-5.1	-	-
ROANOKE RIVER NEAR LOCUST DALE, VA	1985	2.5	-	-
ROANOKE RIVER NEAR LOCUST DALE, VA	2009	3.5	-	-
ROANOKE RIVER NEAR CULPEPER, VA	2009	-8.9	-6.8	-7.1
01668000 RAPPAHANNOCK RIVER NEAR FREDERICKSBURG, VA	1985	-12.7	52.5	79.9
01668000 RAPPAHANNOCK RIVER NEAR FREDERICKSBURG, VA	2009	6.3	27.9	28.3



What are the Tributary Summaries?

A compilation of information by tributary or region on:

- Tidal water quality and trends,
- Watershed characteristics and changes,
- **Landscape drivers.**






Figure 2. Distribution of land uses in the Rappahannock watershed. Percentages are the percent change from 1985 for each source sector.



Where can I access the tributary summaries?

Chesapeake Assessment Scenario Tool

HOME PUBLIC REPORTS LEARNING ABOUT CONTACT US

New to CAST?
Rapidly develop scenarios for reducing nitrogen, phosphorus, and sediment with varying best management practices to streamline environmental planning.
Register for increased functionality and to stay updated.
Register Where To Start

RESOURCES

- DEVELOP A PLAN
Get answers to your questions about how to use CAST to develop a plan.
Develop A Plan
- SOURCE DATA
Download data tables including information on load sources and agencies, BMPs, animals, geographic references and delivery factors.
View Source Data
- BMPS
View information on best management practices (BMPs) including calculations, a quick reference guide, and protocol and expert panel reports.
Learn More
- MAP TOOLS & SPATIAL DATA
View geographical information and shapefiles.
Learn More
- COSTS
Download BMP costs data and view cost profiles for each state and Chesapeake Bay Watershed.
Learn More
- TRACK PROGRESS**
View helpful information on verification, river trends, how to submit progress data via NEIEN, and modeling Federal facilities.
Track Progress

Chesapeake Bay Program Office Software Release: 6.10.1

Chesapeake Assessment Scenario

HOME PUBLIC REPORTS LEARNING ABOUT CONTACT US

The following information is available below:

- Phase 3 WIP BMP Information
- Trends Over Time
 - BMPs implemented
 - Loads delivered to the streams and Bay
 - Wastewater
 - Nutrients applied to the land
 - Animal numbers
 - Septic systems
 - Estuary Summaries
- Tributary Summaries
- River Trends
- Progress Reporting
 - Phase 6 NEIEN Appendix
 - Codes List and Tables
 - NEIEN Submission Instructions
 - Document Exchange Template
 - NEIEN Schema
 - CAST data update frequency

[CAST - TMDL Tracking \(chesapeakebay.net\)](https://chesapeakebay.net)




Tributary Summaries

The Chesapeake Bay Program and its partners compiled tributary basin summaries for 12 major tributaries or tributary groups in the Chesapeake Bay Watershed. These documents summarize the following in one place: 1) How tidal water quality changes over time; 2) How factors that drive those changes change over time; and, 3) Current state of the science on connecting change in aquatic conditions to its drivers.

- Choptank (includes the Choptank, Little Choptank, and Honga) [Summary, Appendix](#)
- Potomac: [Summary, Appendices, Story Map](#)
- Maryland Mainstem (includes the five Chesapeake Bay mainstem segments within the Maryland state boundary. Drainage basins include the Susquehanna River and upper Chesapeake Bay shorelines) [Summary, Appendix](#)
- Maryland Upper Eastern Shore (includes the Northeast, Bohemia, Elk, Back Creek, Sassafras, and Chester Rivers, the Chesapeake & Delaware Canal, and Eastern Bay) [Summary, Appendix](#)
- Maryland Upper Western Shore (includes the Bush, Gunpowder, and Middle rivers) [Summary, Appendix](#)

Where can I access the tributary summaries?

https://www.chesapeakebay.net/who/group/integrated_trends_analysis_team



The screenshot shows the top portion of the Chesapeake Bay Program website. At the top left is the logo, which features a stylized bird in flight over a sun and water, with the text "Chesapeake Bay Program" and the tagline "Science. Restoration. Partnership." below it. To the right of the logo is a navigation menu with five items: "Discover the Chesapeake", "Learn the Issues", "State of the Chesapeake", "Take Action", and "In the Future". Below the navigation menu is a breadcrumb trail: "WHO WE ARE > HOW WE'RE ORGANIZED > INTEGRATED TRENDS ANALYSIS TEAM". The main heading of the page is "Integrated Trends Analysis Team" in a large, bold, black font. Below the heading are three social media icons: Facebook, Twitter, and Email. At the bottom of the screenshot, the beginning of a section titled "Upcoming Meeting" is visible, with a partial view of a meeting card for January 10:00.

Projects and Resources

Tributary Summaries

The Chesapeake Bay Program and its partners produce tributary basin summary reports for the Bay's 12 major tributaries using tidal monitoring data from more than 130 monitoring stations throughout the mainstem and tidal portions of the Bay. These reports use water quality sample data to summarize 1) How tidal water quality (TN, TP, DO, Chlorophyll a, Secchi Depth) has changed over time, 2) How and which factors may influence water quality change over time, and 3) Recent research connecting observed changes in aquatic conditions to its drivers.

These documents can be found here: <https://cast.chesapeakebay.net/Home/TMDLTracking#tributaryRptsSection>

How do we use this information?

- To answer questions such as:
 - *Have water quality indicators in my river been improving or degrading over time?*
 - *How have landscape factors that drive water quality change in my watershed changed over time?*
 - *What clues do they provide that might explain observed water quality change (or lack of change)?*
 - *What should I target to turn a degrading trend around or maintain improvements for future water quality and living resource conditions?*
 - *What should scientists focus our analyses on to provide better answers in the future?*
- As readily-available **background** for change over time observed with monitoring data.
- **Model Evaluation** for MTM

Case Studies today:


- 1) Potomac Tributary
Summary
- 2) York Tributary Summary

Case Study 1: Potomac Tributary Summary

- Completed Dec, 2020.
- Uses data from 1985-2018.

Keisman, J., Murphy, R. R., Devereux, O.H., Harcum, J., Karrh, R., Lane, M., Perry, E., Webber, J., Wei, Z., Zhang, Q., Petenbrink, M. 2020. Potomac Tributary Report: A summary of trends in tidal water quality and associated factors. Chesapeake Bay Program, Annapolis MD.

- Story Map produced by USGS:
<https://wim.usgs.gov/geonarrative/potomactrib/>






Case Study 1: Estimated Loads

Estimated loads to tidal portions of the tributary from USGS RIM Stations at the tidal-nontidal interface.


- True condition loads are highly variable due to freshwater flow:
 - TN has an overall decline that is significant due to substantial efforts to reduce Nitrogen loads from WWTPs and the introduction of the Clean Air Act..
 - TP has an overall increase that is not significant.
 - SS has an overall decline that is not significant.
 - Point source loads have decreased for TN and TP.
- Note that “flow-normalized” loads are mostly decreasing in the Potomac

Total estimate of observed loads to tidal Potomac




Case Study 1: Chlorophyll *a*

- Long Term vs Short Term Trends:
 - Long term observed change
 - Long term flow-adjusted change (i.e., if flow had been average)
 - Recent 10-year observed change
 - 10-year flow-adjusted




Case Study 1: Chlorophyll *a*

- Trends for chlorophyll *a* are **split into spring and summer** to analyze chlorophyll *a* during the two seasons when phytoplankton blooms are commonly observed.
- **Mixed Trends:**
 - Long Term mostly degrading or showing no trend.
 - Short term trends also mixed, with a few more improvements





Case Study 1: Secchi

- A measure of visibility through the water
- Shows mostly degradation or no trend.
- Fairly consistent with chlorophyll *a*.



Case Study 1: Bottom DO

- Summer (June-Sept) bottom DO is improving at many stations overall.
- Possible improvements over the short-term at the deepest stations are a good sign too (and consistent with other deep areas of the Bay).



Case Study 1: WQ Status

- Tracking Open Water, Deep Water and Deep Channel DO Criteria.
- We include a record of the evaluation results indicating whether different Potomac segments have met or not met specific WQ criteria for DO.



Open Water Summer Criteria Status													
time period	ANATF_DC	ANATF_MD	PISTF	MATTF	POTTF_DC	POTTF_MD	POTTF_VA	POTOH1_MD	POTOH2_MD	POTOH3_MD	POTOH_VA	POTMH_MD	POTMH_VA
1985-1987							ND		ND	ND	ND		ND
1986-1988							ND		ND	ND	ND		ND
1987-1989							ND		ND	ND	ND		ND
1988-1990							ND		ND	ND	ND		ND
1989-1991							ND		ND	ND	ND		ND
1990-1992							ND		ND	ND	ND		ND
1991-1993							ND		ND	ND	ND		ND
1992-1994							ND		ND	ND	ND		ND
1993-1995							ND		ND	ND	ND		ND
1994-1996							ND		ND	ND	ND		ND
1995-1997							ND		ND	ND	ND		ND
1996-1998							ND		ND	ND	ND		ND
1997-1999							ND		ND	ND	ND		ND
1998-2000							ND		ND	ND	ND		ND
1999-2001							ND		ND	ND	ND		ND
2000-2002							ND		ND	ND	ND		ND
2001-2003							ND		ND	ND	ND		ND
2002-2004									ND	ND			
2003-2005									ND	ND			
2004-2006													
2005-2007													
2006-2008													
2007-2009													
2008-2010													
2009-2011									ND	ND			
2010-2012									ND	ND			
2011-2013									ND	ND			
2012-2014									ND	ND			
2013-2015									ND	ND			
2014-2016									ND	ND			
2015-2017									ND	ND			
2016-2018									ND	ND			

Case Study 1: WQ Status

- Tracking Open Water, Deep Water and Deep Channel DO Criteria.
- We include a record of the evaluation results indicating whether different Potomac segments have met or not met specific WQ criteria for DO.




Deep Water and Channel Status

time period	Deep Water		Deep Channel	
	POTM H_MD	POTM H_VA	POTM H_MD	POTM H_VA
1985-1987		ND		ND
1986-1988		ND		ND
1987-1989		ND		ND
1988-1990		ND		ND
1989-1991		ND		ND
1990-1992		ND		ND
1991-1993		ND		ND
1992-1994		ND		ND
1993-1995		ND		ND
1994-1996		ND		ND
1995-1997		ND		ND
1996-1998		ND		ND
1997-1999		ND		ND
1998-2000		ND		ND
1999-2001		ND		ND
2000-2002		ND		ND
2001-2003		ND		ND
2002-2004				ND
2003-2005				ND
2004-2006				
2005-2007				
2006-2008				
2007-2009				
2008-2010				
2009-2011				
2010-2012		ND		ND
2011-2013				ND
2012-2014				ND
2013-2015				ND
2014-2016				ND
2015-2017				ND
2016-2018				ND

Case Study 1: WQ Status

- Comparing trends in station-level DO concentrations to the computed DO criterion status for a recent assessment period can reveal valuable information:
 - Whether **progress is being made towards attainment in a segment that is not meeting the water quality criteria,**
 - or conversely the possibility that **conditions are degrading even if the criteria are currently being met.**



Case Study 1: Potomac Tributary Report

- The Potomac Tributary Report is the only finished summary *meaning*,
 - The report contains an “**Insights on Changes**” section, which pulls in additional research to provide further context for the WQ trends and changes in the watershed.

To answer questions like:


*→ **How do tidal waters respond to actions in the watershed?** (Actions may include WWTP upgrades, implementation of agricultural best management practices to reduce nutrient pollution, etc.)*

Two important findings from the Potomac Tributary Report:

1. Local response to large nutrient reductions happens and is clearly shown with the data.
2. Long-term response to watershed-wide nutrient reductions is happening in the tidal waters.


Tidal water response: 1) *Local response to large nutrient reductions happens*

Mattawoman Creek: Very large WW load reductions



SAV coverage (ha), water clarity (Secchi disk depth), and algal biomass (chlorophyll *a* concentration) in Mattawoman Creek. From Boynton *et al.* (2014).


Gunston Cove: Very large WW load reduction




Algal biomass (as chlorophyll *a*), Secchi depth, and SAV acreage for the period 1994 – 2016 in Gunston Cove. From Jones *et al.* (2017).

Tidal water response: 1) *Local response to large nutrient reductions happens*

Mattawoman Creek: Very large WW load reductions



Gunston Cove: Very large WW load reduction




What this tells us: This data clearly shows that investment in large-scale nutrient reductions is successful for improving water quality dramatically in local systems.

Tidal water response: 2) Long-term response to watershed changes is happening

- Over the long-term, nutrient loads have decreased across the Potomac watershed.
- Tidal nutrient concentrations have decreased at almost all tidal stations.


Table 3. Trends (2009 – 2018) in flow normalized total nitrogen (TN), total phosphorus (TP), and suspended sediment (SS) for nontidal network monitoring locations in the Potomac River watershed.

Parameter	No. of stations	Value	Trend direction		
			degrading	improving	no trend
TN	28	n	7	14	7
		median %	15.4%	-5.8%	1.1%
TP	18	n	0	12	6
		median %	-	-28.9%	8.5%
SSC	18	n	5	5	8
		median %	23.7%	-24.4%	5.2%




Tidal water response: 2) Long-term response to watershed changes is happening

- These tidal trends are **not just local response**, but have been shown to be impacted by loads from many types of sources.





Mean annual change in the percent contribution of nitrate from wastewater, fertilizer, atmospheric deposition, and nitrification, based on an isotope mixing model, with distance down-estuary from wastewater treatment plant output. Adapted from Pennino *et al.* (2016).




Tidal water response: 2) Large-scale, long-term response is happening

- Other water quality responses are not as clear
- But research shows there may be a reason: **Nutrients have improved, but still need to be lower to limit phytoplankton growth in most places.**



Tidal water response: 2) *Large-scale, long-term response is happening*

- Other water quality responses are not as clear
- But research shows there is a reason: Nutrients have improved, but still need to be lower to limit phytoplankton growth in most places.





What this tells us: The data shows that watershed-wide nutrient reductions have improved nutrients in the Potomac. The science supports the conclusion that with more reductions, improvements will continue.

Case Study 2: York River

- Watershed stations: Mostly increasing flow-normalized nutrient loads
- Tidal: Long-term TN and TP trends are mixed, but more increasing than decreasing

→ *Patterns are relatively consistent watershed-to-estuary*





Case Study 2: Example York River

- Increasing development and fairly consistent agricultural land use in the last decade
- Expected long-term loads have plateaued, or increased

→ *This and similar information can help understand why nutrients are not decreasing and help target actions*

Distribution of land uses in the York watershed. Percentages are the percent change from 1985 for each source sector.



Expected long-term average loads to the tidal York, as obtained from the Chesapeake Assessment Scenario Tool (CAST-19). Data shown are time-average, steady-state (Not true conditions).


Tributary Summaries

How can the Tributary Summaries support the Modeling Workgroup?

- The summaries provide a comprehensive overview of water quality changes in the tidal watershed, isolating the factors influencing changes in water quality.
- Modelers can use tributary summaries to better inform and guide their tributary models and phase 7 development.
 - Help answer questions like:
 - What has changed in the tidal trends that the model isn't capturing?
 - Are we accurately reflecting the long-term trends with our models?
- Aligning Tributary Summary updates with the MTM workplan

Potomac Tributary Story Map

- Story Map produced by USGS:
<https://wim.usgs.gov/geonarrative/potomactrib/>



Next Steps for the Tributary Summaries

- Discussing priority for updating tributary summaries
 - Rappahannock, James River, York River, Eastern Shore Tributary, Patapsco/Back River
- Introducing “Insights on Changes” section to other tributary summaries.
- Considering addressing climate change in the reports.
 - Ex. Historical trends of rainfall duration and intensity
 - Move Water Temperature Tidal Trends from Appendix to main report

Please Check out the 2020 Tidal Trends!

Long-Term and Short-Term Changes on the ITAT Webpage for:

- TN
- TP
- TSS
- Chlorophyll-a
- Secchi Depth
- DO
- Water Temperature

Maps of 2020 Tidal Water Quality Change

1. Long-Term Change

Observed change in water quality by station from the beginning of the period to 2020.

Surface Total Nitrogen, Annual 1985-2020 (462.47 KB)

Surface Total Phosphorus, Annual 1985-2020 (470.11 KB)

Surface Chlorophyll-a, Spring 1985-2020 (466.4 KB)

Surface Chlorophyll-a, Summer 1985-2020 (488.69 KB)

Secchi Depth, Annual 1985-2020 (449.67 KB)

Surface Total Suspended Solids, Annual 1999-2020 (444.54 KB)

Surface Water Temperature, Annual 1985-2020 (482.07 KB)

Bottom Dissolved Oxygen, Summer 1985-2020 (467.25 KB)

Overview of Findings: https://www.chesapeakebay.net/channel_files/44102/2020_tidal_trends_-_itat_11-19-21.pdf

New Manuscript: “Nutrient Improvements in Chesapeake Bay: Direct Effect of Load Reductions and Implications for Coastal Management”

<https://pubs.acs.org/doi/10.1021/acs.est.1c05388>

Contact Information

- **ITAT Co-coordinator:** Breck Sullivan, USGS:
bsullivan@chesapeakebay.net
- **ITAT Co-coordinator:** Vanessa Van Note, EPA:
VanNote.Vanessa@epa.gov
- **ITAT Staffer:** Alex Gunnerson, CRC/CBP
agunnerson@chesapeakebay.net
- Rebecca Murphy, UMCES/CBP
rmurphy@chesapeakebay.net

Links and References

CAST/Tributary Summaries: <https://cast.chesapeakebay.net/Home/TMDLTracking#tributaryRptsSection>

Potomac Story Map: <https://wim.usgs.gov/geonarrative/potomactrib/>

References:

Boynton, W. R., C. L. S. Hodgkins, C. A. O’Leary, E. M. Bailey, A. R. Bayard and L. A. Wainger, 2014. Multi-decade responses of a tidal creek system to nutrient load reductions: Mattawoman Creek, Maryland USA. *Estuaries Coasts* 37:111-127, DOI: 10.1007/s12237-013-9690-4.

Jones, R. C., K. Mutsert and A. Fowler, 2017. An ecological study of Gunston Cove: Final report. Provided to the Department of Public Works and Environmental Services, Fairfax County, VA, p. 181.
https://www.fairfaxcounty.gov/publicworks/sites/publicworks/files/assets/documents/gunston-cove_2.pdf.

Keisman, J., Murphy, R. R., Devereux, O.H., Harcum, J., Karrh, R., Lane, M., Perry, E., Webber, J., Wei, Z., Zhang, Q., Petenbrink, M. 2020. Potomac Tributary Report: A summary of trends in tidal water quality and associated factors. Chesapeake Bay Program, Annapolis MD. <https://cast.chesapeakebay.net/Home/TMDLTracking#tributaryRptsSection>

Pennino, M. J., S. S. Kaushal, S. N. Murthy, J. D. Blomquist, J. C. Cornwell and L. A. Harris, 2016. Sources and transformations of anthropogenic nitrogen along an urban river–estuarine continuum. *Biogeosciences* 13:6211-6228, DOI: 10.5194/bg-13-6211-2016.